



BEQUEATHED

to the


Devonshire Hospital, Buxton,

by the late

George Lorimer, A.M., M.D.,
etc.

(House Surgeon from 1873 to 1884).

Honorary Medical Officer, 1902 to 1907.



Digitized by the Internet Archive
in 2016

<https://archive.org/details/b2814871x>

W. Brewster M.D.

A TREATISE
ON
THE NATURE AND CURE OF
GOUT AND RHEUMATISM,
&c. &c.

A TREATISE
ON THE
NATURE AND CURE
OF
GOUT AND RHEUMATISM,

INCLUDING
GENERAL CONSIDERATIONS
ON
Morbid States of the Digestive Organs;
SOME
REMARKS ON REGIMEN;
AND
PRACTICAL OBSERVATIONS ON GRAVEL.

~~~~~  
BY CHARLES SCUDAMORE, M.D.

*Member of the Royal College of Physicians,  
Of the Medical and Chirurgical Society of London, &c. &c.*

~~~~~  
Principiis obsta ; sero medicina paratur,
Cum mala per longas invaluere moras. OVID.

=====

THE THIRD EDITION,
REVISED AND MATERIALLY ENLARGED.

=====

LONDON:

PRINTED FOR THE AUTHOR,

By Joseph Mallett, 59, Wardour-street, Soho;

AND SOLD BY

LONGMAN, HURST, REES, ORME, AND BROWN, PATERNOSTER ROW.

—♦—
1819.

HEBERDEN
SOCIETY
LIBRARY

TO

MATTHEW BAILLIE, M.D.

FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS,

F.R.S. of Lond. & Edin. &c. &c. &c.

DEAR SIR,

As a testimony of that esteem and respect, which your professional and private character equally inspire, permit me the honour of inscribing to you the following pages.

The former editions of this work having met with a flattering reception, and a new one being now demanded, I am again afforded the satisfaction of expressing towards you those sentiments which I must ever entertain; and of exerting my best endeavours to improve the principles of the healing art, as connected with the important subjects of the present Treatise.

That you may long continue to promote the cause of Science and Humanity, by your valuable labours and bright example, is the constant and sincere wish of,

DEAR SIR,

Your obliged

and faithful Servant,

CHARLES SCUDAMORE.

6, Wimpole Street,

August 7th, 1819.

CONTENTS.

<i>Dedication</i>	v
<i>Preface to the First Edition</i>	xi
———— <i>Second Edition</i>	xxi
———— <i>Third Edition</i>	xxv

GOUT.

Dr. Cullen's Nosological Arrangement examined, 1.—
Observations on Sauvages and other Authors, 10.—
Own Arrangement, 13.

ACUTE GOUT.

*History of, comprehending an Account of the Premo-
nitory Symptoms; the Paroxysm; its subdivisions
into the Symptoms of a first Fit; the subsequent
Attacks; the Sequelæ; 15.—The remote Causes; and
first, of the Predisposing, 52.—A Morbid State of
the Digestive Organs, generally considered, 78.—Of*

RHEUMATISM.

Introduction and general Observations, 639.—Textures affected by Rheumatism, 642.—Symptoms of Acute Rheumatism, 646. — Sequelæ, 649. — Predisposing Causes, 650. — The Exciting Cause; — Proximate Cause, 653;—Ratio Symptomatum, 656. — Diagnosis, 656.—Constitutional Treatment, the remedies being separately considered, 657.—Local Treatment, 667. — Treatment in the Convalescence, 669.—Prophylaxis. ibid.

CHRONIC RHEUMATISM.

General Description, 671.—Textures affected, 673.—Sequelæ.—Remote Causes, 676.—Diagnosis, 676.—Treatment generally considered, 679.—Cases, various, in illustration, in which Sciatica is included, 692.—Conclusion, 712.

PREFACE

TO THE FIRST EDITION.



ALTHOUGH the subject of Gout has engaged the attention of Authors from the earliest records of Medicine, there are few diseases which have remained more enveloped in doubtful theory ; or in which, the practice has been so unsettled and involved in prejudice.

It may indeed be affirmed, that the Gout has at all times been a favourite object for the exercise of Empiricism ; and every day's observation demonstrates, that, whilst the impositions of quackery are received by gouty persons with the fondest credulity, the art of regular medicine is treated

with indifference, or even rejected: so powerfully does an unknown agent affect the imagination, and indispose the mind to the exercise of the more sober faculty of judgment.

How irrational, in principle, the employment of empirical remedies really is, may in a few words be pointed out. Even the occasional success of a nostrum is injurious in its consequences, from the numerous misapplications to which such success gives rise; and, in this way, the more popular its character, the more extensive is the evil.

In a medicine even of known composition and properties, when administered as a remedy for the same disease in different persons, we see the action so much modified by individual constitution, that in one instance it aggravates the symptoms, as much as in another it gives relief. How much more indefinite and hazardous must be the application of a medicine, which is offered as a remedy for almost every kind of disease, and indiscriminately adopted in every kind of constitution. If any active powers belong to such a medicine, its inappropriate employment must produce occasional

injury ; if it be inert in its properties, it becomes a strong negative source of harm, by excluding other means which might be beneficial.

It is now to be asked, in what degree has the reproach which has fallen on the Medical Art, with regard to Gout, been really deserved ?

A large class of gouty subjects surrender themselves wholly to empirical treatment. Another portion, and probably not the smallest, prefer their *seeming* security in exercising the difficult philosophy of *patience and flannel*. They do this, either because they distrust the efficacy of remedies ; or because they labour under the infatuation, that the gout is a certain remedy for other evils ; and should rather be invited than forbidden, in its approach and stay. The observation of the late experienced Dr. Heberden is still applicable to the present question. “ But as the supposed hazard of curing the gout is now the general belief, it seems reasonable that it should not be made the opprobrium of the art of healing, till the patients will conquer their fears, and allow that it ought to be cured. Happy, however, would it be for man-

kind, if the difficulty of curing the gout were to become as little as the danger of it*.”

The greater number of authors upon Gout, from the earliest period to the present, have adopted a species of empiricism, in founding their whole rule of practice on some imaginary hypothesis of a *proximate cause*. Nor can the regular physician be entirely acquitted of having lent his sanction to empirical remedies. In proof of this, I may advert to the medical patronage at first so freely bestowed on the *Eau Medicinale*. Its magic powers of ease were very naturally hailed with delight by the suffering patient; and Gout no longer appeared a disease of difficult management, or a source of terror. The sequel, however, has shewn the fallacy of the charm; and has served to illustrate the fact, that the best remedies are not those which hastily produce a palliative and transient relief; but, on the contrary, such as are administered upon sound general principles; are carefully adapted to the variations of every particular case and consti-

* Commentaries, p. 47. — “It is not a sacred disease: There will be no profaneness in handling it freely.”—*Rush*.

tution; and are pursued with steadiness and perseverance.

It must, I fear, be allowed, that the regular treatment of Gout has been considerably neglected by the Profession. It appears to have been less an object of attention than any other disease of equal importance; and it is usually pronounced by all parties to be intractable in its nature, and but little understood. In this reflection, however, I do not overlook the several modes of active practice which have at different times been proposed within these few years. Yet it has always appeared to me, that the views which have been formed of the nature and cure of Gout, have been too partial; and deficient in attention to that code of general principles, which is essential to the true understanding and treatment of every disease.

Upon this conviction I was led to the present undertaking. I am well aware how imperfectly I have supplied the deficiency of which I complain; but I shall have performed no useless task, if it should appear that I have entered into the right path of investigation; and have succeeded in mak-

ing some advances towards a clear and comprehensive illustration of the subject.

In the tabular arrangement which is annexed to this Treatise, I have detailed the *analytical* method of examination, which I have adopted in the study of the cases which have come under my care, and within my immediate information. From the method itself I have derived great facility of arranging my facts, and establishing my conclusions; and it is after no inconsiderable exercise of observation and reflection, that I venture to present these pages to the public.

On these grounds I feel myself entitled to advance the following general positions:

That the Gout is a disease not only injurious to the constitution, but destructive of the organization of the particular textures which it affects; and, by such united influence, tends both to shorten and embitter life:

That it is as completely within the useful influence of Medicine as any other severe disease:

That the fit may be immediately relieved in its painful symptoms, and materially shortened in its duration :

That most of its natural bad consequences may, by timely care, be prevented ; and, finally,

That all these advantages may be afforded by means, which, in removing the disease, tend at the same time to restore the constitution.

The security of the patient from future attack, is a question of separate consideration. It is true that it depends much on the Medical Treatment which is pursued in the paroxysm, and during the period of convalescence ; but it is still more connected with the patient's own peculiar care. The gouty diathesis being established in the constitution, it is excited into action by many remote causes ; and, certainly, of the chronic diseases, this appears to be influenced by a greater variety of hurtful agents than any other. Hence, without a corresponding observance of regimen, and general care, the gout, ere long, returns. In this case it will probably happen that the treatment which

was successful in the paroxysm, however judicious it may have been, is regarded as of little value ; or is even accused as being the source of disappointment.


The *prophylactic* means, indeed, deserve the most serious and attentive consideration in every individual case. In no disease, however, can the skill of the Physician be proof against a want of care in the general habits of the patient ; and those who are not their own Physician in this respect, have no right to censure the advice which they but imperfectly follow.

Upon the subject of *Rheumatism*, I have, on the present occasion, confined myself within narrow limits ; and the practical consideration of the chronic species of this disease, I have entirely reserved as matter for a future volume.

I have long conceived that some useful light might be thrown on the phenomena of Rheumatism, both in relation to its theory and treatment, by attempting a more accurate principle of classification than has yet been laid down by authors ;

and by employing that analytical method of inquiry, which unfolds to our view all the various causes and phenomena, both local and constitutional, which can be discovered in any way to influence, or to be connected with, the disease.

The illustrious Sydenham remarks, “ The improvement of Physic, in my opinion, depends upon collecting as genuine and natural a description, or history of all diseases, as can be procured ; and laying down a fixed and complete method of cure.”



PREFACE

TO THE SECOND EDITION.



AN early demand for the first impression of a work, is one of the most satisfactory proofs which an author can receive, that his labours have neither been altogether devoid of interest, nor exerted in vain.

Under the influence of this consideration, I have carefully endeavoured to render the present edition more copious and useful than the former. For this purpose, I have attentively revised the opinions before advanced ; I have again studied the cases from which I had drawn my conclusions ; and have also added a detail of all those results and observations, which a progressive experience and more mature reflection have since afforded.

Convinced of the real and extensive benefits, which are to be derived from a scientific and patient

cultivation of the healing art, I have strenuously opposed every innovation of empiricism and empirical principles ; and have endeavoured to shew, that it is only by a steady and unremitting observation of natural phenomena, that the philosophy of medicine can be established on a firm basis ; and the profession itself be elevated to its proper dignity and rank.

Between the diseases *Gout*, *Gravel*, and *Rheumatism*, there are so many points of natural affinity, that they have appeared to me to form a very suitable class of objects to be embraced in the design of the same work.

In the Treatise upon Rheumatism, which I have still offered in a limited manner, as compared with the extent of the subject, I have, in my introductory observations, entered into an apology for its brevity, which I hope will be found satisfactory.

In advancing many critical objections to the opinions and experiments of other authors, I trust that I shall not be found guilty of injustice or want of candour. I have used only that freedom towards them, which I desire and invite towards myself.

In the formation of every science, the collision of ideas, and comparative observations, invariably tend to elicit information and establish truth.

In an art like that of practical medicine, which, under the most favourable circumstances, is difficult, and often deceptive, the distrust of received opinions and high authorities is calculated to lead to the discovery of new and useful facts ; and to render the researches of genius and wisdom more firm in their foundation. A passive contentment with what has been already accomplished, would at once tend to narrow the avenues of science ; to arrest the progress of improvement ; and to present a strong barrier to the advancement of human knowledge.

The mere desire of novelty is an unworthy passion ; but that spirit of inquiring scepticism, which urges the mind to examine with strict scrutiny the present records of science, I hold to be the very spring of action towards all improvement.

How, unless in this manner, could the chaff be winnowed from the grain ? How could the illu-

sions of false or mistaken philosophy be dispersed, and made to yield their place to the substantial forms of truth?

Impressed with these ideas, and at the same time desirous to cherish a due respect for the opinions of others, I must agree in the sentiments of Pliny, when he says,

“ Quamvis enim cedere auctoritati debeam, rectius tamen arbitror, in tanta re, ratione quam auctoritate superari.”

Lib. i. Ep. 20.

PREFACE

TO THE THIRD EDITION.



THE additions to the present Treatise are so numerous, and so much interspersed throughout the work, that I have found it impossible to embody them into an Appendix, as I should have been gratified in doing, for the convenience of those who may have purchased either of the former editions. But should another at a future period be required, I fully purpose to publish the work in two volumes, confining the subject of Gout entirely to the first ; and to print a certain number of additional copies of the second volume, for the accommodation which I have stated.

The edition now offered to the public, will be found to contain nearly one third proportion of matter more than the last. The additions consist almost wholly of practical observations and cases ; together with a Series of Experiments, chiefly relating to the examination of those Gout Nostrums

which have so long engaged the curiosity of the public, and still give occasion to doubt and discussion, - both as to the nature of their properties, and the propriety of their administration. Upon the subjects of Gravel and Rheumatism, I have enlarged as far as my limits would permit; and have endeavoured to furnish some useful materials, although professedly very incomplete.

With respect to the Experiments, notwithstanding that I have already introduced them with some prefatory observations, I shall beg leave in this place also to say a few words. It was at first my intention to have printed the details separately from the Treatise, and to have offered them to the sole attention of the Medical Reader; but, on mature reflection, I considered that they might prove both interesting and useful to the gouty patient, as furnishing some important hints respecting the immediate treatment of his disorder. The recital of animal sufferings, inflicted in the way of experiment, certainly in itself requires some apology; and I should willingly have considered the delicacy and feelings of the general reader, in suppressing many particulars; but as constantly as

I attempted this, I found that I materially diminished the information which I had to offer; and that I could not so satisfactorily shew the grounds of my own conviction, as to the hurtful consequences of employing these medicines, and the importance of deterring gouty persons from resorting to them.

The results of these experiments have proved a great source of instruction to myself; and I will appeal to every one, whether it is not allowable that a medicine of doubtful character should be made to shew its deleterious agency on animals; if, from such means, any decided advantage can be derived to the health and comfort of our fellow creatures.

The high sanction which the use of these pretended specifics has so long received, demanded from me every collateral evidence to support the opposite opinions which I have ventured to maintain. The point for which I most earnestly contend, is, that the administration of gout nostrums, in addition to any hurtful agency which they possess, involves in the treatment completely false principles. It consti-

tutes a practice essentially unsound. If, therefore, my views be just, and I gain the concurrence of the rational and well-informed; — if I assist in establishing a proper and regular mode of treatment for the gout, and thus lessen the amount of human suffering;—and, finally, if I oppose with success the unworthy claims of empiricism, and gain new adherents to the cause of scientific medicine; I shall be contented with my labours, and reflect with satisfaction on their result.

Should the doctrines and opinions which I have advanced be founded in truth, they will stand the test of time and opposition;—if, on the other hand, they originate in error, they must, when submitted to the same ordeal, as inevitably fall.

The reader is requested to correct the following typographical errors.

- Page 31, line 22, *for* distented, *read* distended.
- 46, — 11, *for* inspiration, *read* inspissation.
- 53, — 23, *for* 124 to 91, *read* 84 to 105.
- 62, last line, *for* twenty-three, *read* twenty-five.
- 92, line 27, *for* flushings, *read* flutterings.
- 94, — 10, *for* afflict, *read* affect.
- 558, note, line 1, *for* magneisa, *read* magnesia.

A TREATISE,

§c. §c. §c.

A CLEAR arrangement and exposition of the principles of a science, essentially conduce to establish correct views with regard to its theory, and to facilitate its practical applications. This is more especially the case with regard to pathology and the practice of medicine; and I trust, therefore, that no apology will be necessary for the extent of preliminary matter which the following pages may embrace.

I feel it, in the first place, incumbent on me to state the grounds on which I have ventured to propose a different nosological arrangement in my description of Gout, from that which was laid down by Dr. Cullen.

This author, who, on the subject of nosology, continues to be the standard authority in the British schools of medicine, has drawn his general characters and varieties of this disease as follows :

“ *Podagra*.—*Morbus hæreditarius, oriens sine causa externa evidente, sed præeunte plerumque ventriculi affectione insolita; pyrexia; dolor ad articulum, et plerumque pedis pollici, certe pedum et manuum juncturis, potissimum infestus; per intervalla revertens, et sæpe*

cum ventriculi, vel aliarum internarum partium, affectionibus alternans.

“ *Varitas 1^{ma}*. Podagra (*regularis*) cum inflammatione artuum satis vehementi, per aliquot dies persistente, et paulatim, cum tumore, pruritu, et desquamatione partis, recedente.

“ *Var. 2^{da}*. Podagra (*atonica*) cum ventriculi, vel alius partis internæ atonia, et vel sine expectata aut solita artuum inflammatione, vel cum doloribus artuum lenibus tantum et fugacibus, et cum dyspepsia vel aliis atoniæ symptomatis, subito sæpe alternantibus.

“ *Var. 3^{ta}*. Podagra (*retrograda*) cum inflammatione artuum subito recedente, et ventriculi vel alius partis internæ, atonia mox insecuta.

“ *Var. 4^{ta}*. Podagra (*aberrans*) cum partis internæ inflammatione, vel non prægressa, vel prægressa et subito recedente, inflammatione artuum.”

I shall offer my comments briefly in the order of the quotations (which will be translated from the preceding original text of the author), passing over those parts, the correctness of which, as general statements, I admit; and reserving further and more extended observations for the respective heads of the subject in the course of this work.

An hereditary disease.—As the gout is so frequently found to occur where no hereditary influence can be referred*, this circumstance is erroneously stated as a definite character of the complaint. Also in many of the

* I formerly stated that gout without any hereditary influence was the most frequent; but my later series of cases does not show this to be the fact. I have therefore varied my critical remark accordingly.

worst examples of gout which have come under my observation, there has not been a trace of the disease in the family.

Arising without any evident cause.—This is obviously expressed by Dr. Cullen as a distinguishing character from rheumatism, which he begins with defining, “a disease from an external and often an evident cause.” Although the gout is a disease arising out of the internal faults of the system, and does not, like rheumatism, originate as a *local disease* of structure externally excited, yet its two-fold character in this respect must not be overlooked.

The gout is often called into action, even in the first fit, by a cause equally external and evident as that which leads to rheumatism: namely, vicissitude of temperature; a sprain; contusion; or any kind of local injury, which, in a person not disposed to gout, would produce only common inflammation, or some other kind of inflammation or diseased action, according to the specific constitutional disposition of the individual.

Fever.—As a general position, fever is correctly stated to be a character in the gouty paroxysm; although it is not invariably an attendant; and indeed may be said to exist only when the action of the heart sympathises with local inflammation and pain, or when some remote cause has induced the inflammatory diathesis.

Pain at a joint, and, for the most part, the great toe, certainly most strongly affecting the joints of the feet and hands.—This implies too strongly that a joint is the only seat of true gout. I admit that, if the foot be the part attacked, the description is, commonly, just; and, also that,

in the majority of examples, the gout chooses the foot in its first attack, and that the ball or first joint of the great toe is the particular part affected. It is in the commencement of the disease that this circumstance is most remarkable; but even in a first attack, it sometimes happens that the inflammation fixes on the tendinous parts of the foot and hand; and in subsequent attacks especially, so many other situations remote from the feet and hands, and not only the articular structure, but tendons and bursæ also, are affected in an equally characteristic manner;—I mean equally serving to constitute the disease as gout—that the definition, putting false limits to the external indications, is calculated to mislead, and therefore is exceptionable. Hence also, podagra, (from *πῆξ*, a foot, and *ἀγῖα*, a seizure) which Cullen*, in imitation of Boerhaave, chose as the title of the disease, is too confined in its expression. Arthritis (from *ἄρθρον*, a joint), for the reason just now stated, is not an accurate term, but deserves to be retained in preference; for the gout, if not situated at a joint, chiefly affects those textures which are more or less subservient in function to the joints.

And often alternating with affections of the stomach, or of other internal parts.—I presume that this passage expresses, that there is frequently much mutual sympathy between the stomach, or some other internal part, and the external seat of complaint; so that when the stomach, for example, is affected with pain, spasm, coldness, nausea, or other uneasy sensation, the occurrence

* Arthritis nomen, utpote apud medicos ambiguum, rejeci, et Podagra nomine, ntpote typum morbi præcipuum notante cum celebri Boërhaavio usus sum.—*Synop. Nos. Method. Cullen.*

of external inflammation and pains, often will produce internal relief; and, in a certain degree, the converse of this may take place: but if the idea be extended to the signification of *alternation* in the inflammatory action, it denotes a change of action which must tend to danger and death; and therefore is an expression not appropriate to a general description of gout.

The sympathetic action which Dr. Cullen appears to describe, relates rather to the passive form of gout (which hereafter I shall denominate *chronic*), in which, the external inflammation is slight and wandering, and external pains reciprocate with many internal sympathies. This definition accords in part with his *atonic variety*.

OF THE VARIETIES.

Variety 1. Gout (regular). — The fundamental distinctions which Dr. Cullen adopts, of *regular* and *irregular* gout, convey, on the one hand, more precision of definition, than the various modifications which the disease assumes, seem to admit; and, on the other, allow more looseness and latitude of application, than is compatible with good practice.

An attack of gout is not essentially the less regular, because it seizes some other part not immediately belonging to a joint; or appears in some other situation instead of the foot and hand; or because it proves of long duration, and changes frequently its seat. The following statement also is deficient in accuracy, and appears to me exceptionable as a definition. *With a sufficiently strong inflammation continuing for several days, and gradually receding with tumor, itching, and desquamation of the part.*

The time of duration here mentioned belongs almost only to the first fit; and in regard even to this, much exception prevails. The *swelling* is not deferred till the inflammation has gradually receded; but usually takes place in the course of twenty-four hours, and is rarely delayed beyond forty-eight. The *itching*, as being the last and least remaining irritation of nerve, and also incidental to a scurfy state of the skin, is an ordinary occurrence; but the desquamation of the part is by no means sufficiently frequent to serve as a strictly characteristic description.

Variety 2. Atonic.—*With an atony of the stomach or other internal part, and either without the expected or usual inflammation of the joints, or with only slight and transient pains in the joints, and often suddenly alternating with dyspepsia or other symptoms of atony.* The *first* part of this definition is descriptive only of the dyspepsia, or other internal derangements, which may take place in the gouty individual as a common occurrence, and without being necessarily dependant on that state of the system which tends to gout. The latter part describes the passive or chronic form of the disease.

Variety 3. Retrocedent.—*With inflammation of the joints suddenly receding, and quickly followed by atony of the stomach, or some other internal part.* I am fully convinced, that inflammation, having even a fatal tendency, is a more frequent consequence of retrocedent gout than atony; and this definition therefore is of great practical importance. In two cases which have come under my own treatment, and which were produced by exposure to wet and cold, very painful inflammatory symptoms affecting the abdominal viscera took place, requiring the prompt and free employment of the lancet.

I do not deny that in debilitated persons, in whom the circulation is weak, the local gouty action may suddenly, and from slight causes, be exchanged for an internal attack of a spasmodic nature; but such a case, I should consider as an exception to the more usual effects of true retrocedent gout; and I am persuaded that we should be very cautious in drawing the conclusion, that *atony* is the internal affection, succeeding to the suspension of that which is external.

Variety 4. Misplaced.—*With inflammation of some internal part, inflammation of the joints either having not preceded, or having preceded, and quickly disappeared.* This definition begins with direct reference to the occurrence of any of the phlegmasiæ in a gouty individual, whether the disposition to the paroxysm exist at the time or not. It concludes with a repetition of what was before said of the retrocedent variety, with the difference, that the new internal action is here unequivocally designated as that of inflammation.

To the term and import of *irregular* gout, I have already adverted, considering it to be like Musgrave's distinction of *anomalous*, uncertain and without limits in its application, and only calculated to preclude the possibility of accuracy either in theory or practice.

Dr. Cullen observes (Par. 518), "Whatever symptoms we can perceive to be connected with, or to depend upon the disposition, which produces the inflammatory affection of the joints, but without its taking place, or being present at the same time, we name the *irregular* gout."

I grant, and even contend, that the disease possesses a complex character; and that it consists of an external local action, which is the offspring of constitutional causes; so that, if it were true that we could point out

any one particular state of the system, as the invariable circumstance either antecedent to the local action which we call the gout; or, co-existing with it, in the clear relation of cause and effect; we might, on the occurrence of such a state of the system, without its being followed by the usual external characters, very justly assert it to be an attack of irregular or anomalous gout. We cannot however boast, that our knowledge of the intimate nature of the disease is sufficient to authorise such conclusions. It should be considered, that gouty persons are not exempted from other diseases; nor are the various morbid sympathies which they suffer internally, of a peculiar or specific nature. It is undoubtedly true, that the familiar employment of these terms, *irregular* and *anomalous*, gives an unbounded latitude to call every disease*, and every morbid sympathy occurring in a gouty individual, a *disguised gout*.

This consideration is of more importance than may at first sight appear. I do not take up the argument for the sake of verbal dispute. A name being so readily found for an obscure disease, the practitioner considers himself as excused from the difficult task of nicer discrimination. He probably directs his treatment according to some prejudiced views of the nature of gout, instead of taking his indications from an attentive investigation, founded on anatomical knowledge, physiological reasoning, and sound pathology. In other words, he treats the disease according to what *it is not*, rather than according to what *it is*. The *patient*, influenced only by the suggestions of his fears, anxiously desires to fix the wander-

* In proof of this opinion, I refer the reader to Musgrave de Arthridite Anomale, and to Par. 518, 519, 520, Cullen's First Lines, Also, to the French authors on Gout.

ing gout, as he pronounces it, in his extremities; and with this view, drinks brandy or madeira. From the depression which he occasionally feels, he seeks the false relief procured by stimulating cordials; and produces a state of intoxicating excitement, at the expence of positive injury to his constitution.

The circulation, when excited by heating cordials, may be urged to relieve itself by that kind of inflammation to which there is the principal tendency in the constitution; and, what may be regarded as a fortunate event under such a mode of proceeding, the gout may take place; but in the mean time, apoplexy, or some other evil much worse than the gout, so earnestly invited, may be produced.

I acknowledge that dyspepsia, and any other chronic affection, and even acute diseases, may receive considerable modifications of character from the influence of a gouty habit, or even from the remote tendency to gout; and that some correspondent modification of treatment may then be required; but this admission is rather general than particular, and forms a part of this medical rule, that our principles and practice should always be applied and varied, in correspondence with the character of the individual constitution and habit.

Finally, in regard to any disease in a gouty person, whether simple or anomalous, I should adhere to the ordinary nomenclature (if a name were desired); taking into consideration any probable influence, which, the arthritic temperament and constitution might exert.

Dr. Cullen has combated the doctrine of a morbid matter with much ability and ingenuity; but he is not himself very satisfactory in the following piece of pathology, Par. 533: "In some persons there is a certain vigorous and plethoric state of the system, which, at a

certain period of life, is liable to a loss of tone in the extremities. This is, in some measure, communicated to the whole system, but appears more especially in the functions of the stomach. When this loss of tone occurs while the energy of the brain still retains its vigour, the *vis medicatrix naturæ* is excited to restore the tone of the parts, and accomplishes it by exciting an inflammatory affection in some part of the extremities. When this has subsisted for some days, the tone of the extremities and of the whole system is restored, and the patient returns to his ordinary state of health."

The loss of tone in the extremities, which is here described as the first link in the chain of the phenomena, is rather fancifully conceived. The system, though in a plethoric state, can scarcely be said to be vigorous. The reflection of the loss of tone from the extremities to the internal functions; the impulse given to the *vis medicatrix naturæ*; and the consequent restoration of the general and local tone by means of the inflammation excited; compose a mass of doctrine, in which, a variety of hypotheses are confusedly blended together, and are calculated to perplex and mislead the judgment.

From the foregoing theory, Dr. Cullen appears to have deduced all his practical views.

Sauvages has given a very complicated arrangement on the subject of gout, creating many distinct species from the modifications which the disease occasionally assumes, by combination with other diseases, or by the influence of the season of the year*. The Greeks distinguished the varieties of gout according to situation; as, podagra in the feet; chiragra in the hands; pechya-

* Arthritis asthmatica, rheumatica, æstiva, hyemalis, &c. &c.
—*Sauvages' Nosol. Method.*

gra in the elbow; gouagra in the knee; dentagra in the teeth; cleisagra in the articulations of the clavicles; onagra in the articulations of the humerus; rachisagra in the spine of the back; and tenontagra in the large tendons*. These terms descended to the Latins, and were indiscriminately applied to gout and rheumatism. Indeed when we consider, that until the time of Ballonius†, rheumatism does not appear to have been considered as a distinct disease from gout, we should surely, on all occasions, receive the authority of the ancients upon the treatment of gout with much circumspection.

Boerhaave is said not to have introduced the mention of rheumatism in his first course of lectures; but he was afterwards but too well made acquainted with it, from becoming himself the subject of the disease.

Sydenham, in 1683, may probably be considered as the first author who wrote with much perspicuity on gout and rheumatism; but I cannot help thinking of this eminent man, that, although his pen had the merit of great excellence, both in style and matter; and he himself appears to have been the very model of a virtuous physician; his doctrines upon gout, which were conceived in the full force of the humoral pathology, have had, even to the present day, a most injurious influence on practice.

The antiquity of the notion concerning the humoral nature of gout, is illustrated by the etymology of its name—*gutta*, a drop‡; signifying, it is said, that the dis-

* Cælius Aurelianus, Lib. v. cap. 2.

† De Rheumatismo et Pleuritide Dorsali. Paris, 1642.

‡ “Gutta dicitur, ex absurda veterum hypothesisi, qua ab humoris cujusdam stillicidio, in articulos facti, cum dolorem priri putabant.”—*Sauvages' Nosol. Method*, vol. ii. p. 19.

ease was caused by the dropping of some humour into the joints.

It may be remarked of the nomenclature of the ancients, that it was always expressive, though often erroneous. The examination of medical etymology through its whole extent, would be a curious and interesting branch of inquiry. I shall not, however, pursue this digression any further, but proceed to the more immediate consideration of my subject.

It appears to me, that in nosological divisions, a degree of refinement is sometimes introduced, which rather perplexes than affords light to the practitioner. In medicine, more than in any other art, it is dangerous to frame distinctions without a difference. With this conviction, and with a persuasion that nicer separations are as useless as they are difficult, I shall be contented to divide gout into *acute*, *chronic**, and *retrocedent*; considering the acute form of the disease, without regard to particular situation, as the first variety; the chronic, as the second; and the retrocedent, as the third. In an attempt to mark the general characters of gout, I feel the

* Dr. Latham, in speaking of the usual denomination of the varieties of gout, thinks that "they might more properly be distinguished like rheumatism, into the acute and chronic."—*Letter on Rheumatism and Gout*, 1796.

Dr. Hamilton makes the following observation: "Perhaps it may be more simple to divide this disease into two kinds only, the *acute* and the *chronic* gout; because, agreeably to the different phenomena, the several gradations, if I may so term them, between the highly inflammatory acute state of it and the lowest debilitated chronic state, the two extremes of the disease, involving all the irregularities of it, may be readily comprehended."—*Letters on Gout*, 1806, p. 70.

necessity of entering into a description rather in detail, instead of confining myself to a brief definition; because, the disease is too complex in its nature, to be distinguished with certainty by a few signs alone.

GOUT.

A constitutional disease, producing an external local inflammation of a specific kind; the susceptibility to it often depending on hereditary bodily conformation and constitution, but with great frequency wholly acquired; almost never occurring before the age of puberty, not frequently under the age of five-and-twenty, and most commonly between the ages of twenty-five and forty; affecting chiefly the male sex; and particularly, persons of capacious chest and plethoric habit; in the first attack, invading usually one foot only, and most frequently at the first joint of the great toe; but in its returns, affecting both feet, or other situations, as the hands, knees, and elbows; and not only in the articular structure, but also in the other textures belonging to the moving powers, different parts being affected together, or in succession; often accompanied with sympathetic inflammatory fever, which is usually marked by nocturnal exacerbations and morning remissions; much disposed to return at periodical intervals, and for the most part preceded by some premonitory symptoms.

ACUTE GOUT.

Inflammation and pain of the articular, tendinous, or bursal structure, usually attacking one part only at the

same time; but in succession of attack, affecting different parts together; with preternatural fulness of the adjacent veins, and, in certain situations, with œdematous swelling of the integuments occurring in twenty-four or forty-eight hours from the invasion of the fit; vivid redness of surface, which is sometimes shining; entire disability of the affected part, with peculiar sensations of burning, throbbing, cutting and pricking, and weight: the action sometimes readily changing situations spontaneously, or from slight causes; terminating almost invariably without suppuration, and usually with some critical indications of the event.

CHRONIC GOUT.

Inflammation and pain more slight, irregular, and wandering, than in the acute; faint redness of surface; much permanent distension of parts, or continued œdema, and impaired moving power; without critical indications of its terminating; commonly associated with a morbid state of the digestive organs, a languid or oppressed circulation, and much nervous irritation in the system.

RETROCEDENT GOUT.

Metastasis, or transference of the gouty action in the paroxysm, from the external part, to some internal organ.

HISTORY OF ACUTE GOUT.

IN some individuals I have traced that a gradual increase of abdominal corpulency, attended with an inert state of bowels, a scanty secretion of urine, and uncomfortable sensations of a fulness of the general habit, has been imperceptibly introductory to the attack. On the other hand it occasionally happens, and especially in the first fit, that the immediate invasion of the disease is not preceded by any warning; but the patient, after going to bed with the ordinary feelings of health, is surprised in the night with the first symptoms of the disease; yet, except in the first fit, we see much more commonly, that some preceding symptoms intimate for several days, or give even a longer notice, that the gout is at hand. Of these *premonitory* symptoms, as I have found them occur in different individuals, more or less connected with a certain attack of the disease, I may enumerate the following:—

Depression of spirits*, with drowsiness and frequent yawning, night-mare, and restless sleep; heart-burn, acidity of stomach, and sometimes to the degree of rejecting acid matter; flatulence; irregular appetite, with oppression after a meal; a frequent sense of coldness and soreness at the epigastric region; general itching of the skin; costive bowels; or, more rarely an irritable state of the canal, amounting to diarrhoea; scanty and deep-

* One gentleman who is usually admonished of a fit by a remarkable depression of spirits, states that sometimes, on the contrary, his mind is surprisingly elated. An excessive appetite for meat, two or three days before the attack, is his most common precursory symptom.

coloured urine, becoming turbid on cooling; or, in opposition to this, it is sometimes copious and pale; pricking and numbness in the lower extremities; muscular twitchings in the day, and catchings or cramps in the upper or lower limbs, more particularly in the one which is about to be attacked, and occurring chiefly in the first attempts to sleep at night; much coldness of the legs and feet, and occasional general chillness; or, general rigors which are even urgent. Some complain of universal feverish sensations, having with cold extremities, heat of the head with flying pains, vertigo, and frequent flushings. In the disturbed functions of the stomach, an irritability which amounts to occasional retchings sometimes takes place. An excessive appetite for one or more days before the fit is not unusual; but this is attended with occasional heart-burn and nausea. One patient informs me, that for a week before the attack his appetite cannot be satisfied. To use his own words, "he could eat all day long." Another also describes this precursory symptom in the same degree, with the addition of having an excessive secretion of saliva. I have seen a gouty person suffer this latter symptom to the degree of active ptyalism, as if under the strong influence of mercury, increasing with the symptoms of the paroxysm; and he states that this always happens with a fit, in its first stage.

The nervous system is often apprised of the approaching gout by previous general lassitude, with much agitation of mind, palpitation of the heart* or of the aorta in

* The morbid sympathies are variously referred, and often without any apparent connection with the cause by which they are supported. I am informed by Dr. Baillie, of a gentleman who suffered palpitation at the heart for six months without relief from medicine; but a fit of the gout coming on, it sud-

some part of its course, but especially in the epigastric region; tremors, and internal flutterings. Sometimes the patient going to bed with no expectation of the approaching evil, sleeps through the night, but is distressed with some horrid dream; and in the morning finds that the demon of his vision has been the gout. A gentleman, who has for many years been afflicted with gout, for a day or two before his attack, has alternate chills and flushings, is extremely nervous, and he says, without the power of restraint, sheds an abundance of tears.

A deaf patient complains that at the appearance of the fit, his infirmity is so much increased, that he cannot possibly be made to hear.

In one person, a heat of the eyes with slight membranous inflammation, is one of the usual premonitory symptoms.

A cough, with much mucous secretion in the tracheal membrane, sometimes precedes the fit; yielding when the gout becomes fixed, or occasionally continuing with the other symptoms.

This description of cough is to be distinguished entirely from the recent catarrhal cough and symptoms, which sometimes usher in a paroxysm, when exposure to wet and cold has been the exciting cause. The mucous membranes of parts which have been once affected with disease, or which are prone to disease, are much disposed to acquire a state of unhealthy action; in apparent sympathy, either with that condition of the constitution which borders on a fit of the gout; or in which, the disease is threatening its return, but does not develope itself.

denly and entirely left him. I have met with a similar case, and which I shall more particularly describe, when treating of chronic gout.

A remarkable irritability of the bladder and urethra, with increased secretion in the mucous membrane of these parts, prevails with some persons shortly before the fit; and when the urethra is affected with stricture, these symptoms almost with certainty take place.

X A gentleman who has a slight stricture, informs me that during a few days before one of his fits of gout, he was troubled with so much ardor urinæ, dysuria, and discharge, that he persuaded himself of the reality of gonorrhœa; and sought advice accordingly. He was judiciously instructed to wait the use of any treatment for such a complaint. When the gout appeared in one foot, all the other symptoms just mentioned almost instantly left him. Sir Everard Home, in his work on Strictures, observes, "So much is the urethra in its natural state under the influence of gout, that it is sometimes affected by it on the coming on of every attack, with all the symptoms of inflammation, as pain in making water, and a purulent discharge; and as soon as the gout fixes itself in the foot, they entirely disappear." He adds, "this natural susceptibility of the urethra to be influenced by gout appears to be much increased, when that canal is in a diseased state, so as to increase all its symptoms, and when they have gone off, to produce a recurrence of them, and prevent the disease to which it is liable, from being completely removed." He offers cases in illustration of these remarks, noticing however, that some gouty persons even with stricture are exempted from this peculiar irritation in connexion with the paroxysm.

I have not witnessed many examples with these complete symptoms described, unless that a stricture has existed; but with the exception of the discharge, I have seen persons affected, in a very remarkable degree, with

inflammatory and spasmodic irritation of the urethra, shortly before the fit; and with them, this occurrence has been a sure presage of the gout which was to follow.

Sometimes the symptoms come on with the paroxysm, and in either case, whether in this manner, or precursory, most usually continue in a greater or less degree, for a few days during its height. In some instances, however, as already mentioned, the whole complaint yields at once to the gout. In other examples the complaint exists in a chronic state, proving very distressful. I shall hereafter relate two urgent cases of this description.

In addition to the local premonitory symptoms which have been mentioned, the following also occur. Unusual weakness and tenderness of joints, and more particularly in the limb threatened with immediate gout, accompanied with shootings, numbness, prickings and transient spasms; a fulness of the veins of the limb, and dark colour of the skin in the part about to be affected with inflammation. Although some stiffness and inability in the moving powers of the joints general precede the fit, yet I am acquainted with one example, in which, the patient expects a long confinement, if he has felt increased energy and alertness of limbs a day or two before his attack: so curiously are diseases modified by idiosyncrasy of constitution.

A swelling of the feet after slight exercise, with uneasy stiffness of the joints, and a dry state of the skin, with more or less of heat in the soles of the feet, may also be mentioned as precursory. One person informs me that he is generally warned of the approaching attack, by finding, a few days before, "that his shoes pinch him."

A sudden cessation of the usual perspiration of the feet, is also noticed by some gouty persons, just before the paroxysm.

A female patient of delicate constitution describes, that a day or two before the fit, she experiences much perspiration of the limbs, great restlessness, cramps and startings of the legs in the early part of the night, followed by composure about five in the morning;—and she notices the veins of the extremities to be remarkably distended.

I should add as a general statement, that the most severe of the local signs which I have just enumerated, affect those only who have experienced frequent returns of the disease; and in whose limbs, more or less of disorganization of structure has been produced.

Those gouty persons who are affected with concretions (chalk-stones), experience for a short time before the fit, pricking pains in the parts where they are situated. This is described even by those, who have minute points of concretion in the lobes of the ears, and in no other part of the body. When these concretions have given rise to ulcerations in the hands or feet, the sores are unusually tender, aching, and sometimes very painful, as the fit is making its approach.

On the subject of precursors of the fit, it deserves to be remarked, that in a general inflammatory diathesis of the system, and usually from partial exposure to cold acting as the exciting cause, an internal part is sometimes *first affected* with inflammation (constituting what Dr. Cullen denominates the *misplaced* variety), either for a short time, or for some continuance; but this at once suddenly subsiding, the *gout* succeeds; apparently concentrating in itself all the inflammatory disposition of the system.

In two cases of pneumonia, which came under my observation, every symptom affecting the lungs immediately disappeared, on the supervention of the gout, that occurred within a few days.

In another instance, the inflammation of the lungs proceeded to its full extent, in union with the gout in both feet.

Morgagni mentions of himself, that he had suffered from an ophthalmia in each eye, without relief from the usual remedies. At length, a mild attack of gout took place in one foot, doubtless solicited (though not expected, it being his first fit) by the pediluvium and friction, of which he had made a free employment. He adds*, "*Oculorum inflammationem statim inivit, ac diebus insequentibus sustulit.*"

Such exchanges of disease have been called *conversions*; and the subject has been illustrated by many authors†.

OF THE PAROXYSM.

THE description of *Sydenham*, drawn from his own sufferings in this disease with the hand of a master, is given with much spirit, and has been generally quoted for its accuracy; but as, in some parts, it is much obscured by the doctrines of the humoral pathology, and is altogether rather circumscribed, I shall venture to delineate its history from the results of my own inquiry and careful observation.

* *Morgagni*, Epist. lvii. p. 221.

† *Hoffman de Morbis Mutandis*. Dr. Ferriar, "on the Conversion of Diseases." Dr. Parry, in his late Work on Pathology, has presented us with much interesting disquisition on the subject.

Of the Symptoms of a First Fit.—Whether any premonitory symptoms have occurred, or that the patient is seized with gout, while in the apparent possession of health, the most frequent time of the active invasion of the fit, is between twelve and three in the morning; the exact period being liable to some variation, according to the peculiarities of constitution in the individual, his habits, and his hour of rest. He is suddenly awoke* with pain in the affected part, which is most commonly the ball of the great toe of one foot only; and immediately experiences, in a slight degree, sensations of heat, stiffness and weight; which soon increase to burning and throbbing. He is restless, feverish, and watchful, till about five or six in the morning, when, under favourable circumstances, an abatement of the symptoms, with gentle perspiration, permits some tolerable sleep†. Usually on the first morning, the surrounding integuments are swollen, the skin is slightly red, in some instances universally shining as if varnished, and the veins of the foot, in a direction from the inflamed part, appear remarkably

* In those cases in which the symptoms have come on with rapidity before bed-time, the patient, according to my observation, has been indulgent in the luxuries of the table, and has had a quick, although imperfect, digestion.

† I have met with an instance of a first fit, in which the patient slept with more than usual profoundness through the night, and was astonished in the morning by finding that the gout was severely fixed in one foot, which he discovered only on attempting to leave his bed. In two other cases also, the sleep was merely disturbed by uneasy dreams; and when the hour of rising came, the gout had to a great degree taken its station, with one patient in the ankle; with the other, at the ball of the great toe, wholly unexpected.

full. In a severe attack, there is scarcely any remission of the symptoms for two or three days ; but more usually, they are suspended or much abated in the day, and return with violence late in the evening, or before midnight, lasting till about five in the morning. Even on the first morning of the disease, the integuments pit slightly on pressure, and on the second morning this effect is very distinct. The surface is now of a vivid scarlet red. The patient complains of shooting pain, throbbing, heat and weight, with intense sensation of gnawing as if from the biting of an animal. The least compression of the joints is scarcely to be borne. The pyrexia and disorder of the natural functions, as dependant on the first fit, appear to be truly sympathetic ; and are proportioned to the local inflammation and pain. In very slight attacks, the sensible influence of the disease upon the system is so moderate, that sometimes the patient thinks he has received a strain, or some kind of local injury, and treats it accordingly*.

In the first fit, the attacks of this kind usually prove only of two or three days' continuance ; but in contradiction to this general statement, I know a case, in which, one foot was first attacked, and it being considered as an inflammation from a supposed strain, was treated with an oatmeal and vinegar poultice. The other foot shortly became affected, and three months elapsed before this patient regained his health. In another case of first attack, the ball of the toe being the part affected, the pa-

* In one instance of unexpected gout in the winter season, a gentleman first persuaded himself that a severe chilblain had seized the foot ; but rubbing the part freely with a stimulating embrocation, he violently aggravated the inflammation, and then discovered his error.

tient, conceiving that he had suffered a strain, applied a stimulating poultice. This fit was more tedious and painful than any which he has since suffered, although he has been gouty for many years.

The œdema which has been mentioned, continues a short time after the disappearance of inflammation; and sometimes the cuticle of the affected part desquamates, with much itching. The duration of a first fit is seldom less than five, or more than ten days. It now and then happens, that the other foot also becomes affected in succession, giving rise to similar phenomena, but with greater consequent disorder of the system; and such fits also are sometimes very tediously protracted, if left to their natural course. I am informed by a lady, that in her first fit, each foot being affected at first in succession, afterwards together, next appearing to recover, and then again alternately relapsing, five months of suffering were passed over before she felt any degree of convalescence. Another lady also relates to me, that her series of sufferings in the first fit was continued beyond four months; in the course of which time, one foot, each knee, wrist, elbow, and shoulder, were affected in as severe a degree as I have ever heard described*..

A gentleman of generous but not intemperate habits, in whose family gout was quite unknown, after a long walk, felt a remarkable stiffness in the tendo achillis, and concluded that he had received a slight strain. The sheath of the tendon became distended, and the integuments

* This case, which from the outline of my statement may appear to belong to rheumatism, was certainly an example of acute and chronic gout. Both from the communication of the medical attendant, and from my own observations since this lady has been under my care, I am convinced of the fact.

rather œdematous, but not until a few days had elapsed, did he discover the real nature of the disease. In the evening, rigors and head-ache took place, and in the morning, the great toe was stamped with the true characters of gout.

I may state as a general fact, that the gout is more disposed to be mild and regular in its first attack in men, than in women.

In one hundred and ninety-eight cases of gout, I have made the following comparison of the parts affected in the first fit:

	<i>Cases.</i>
In the great toe of one foot only.....	130
In the great toe of each foot.....	10
In the little toe and the one adjoining.....	1
In the great toe and instep.....	3
In the great toe and instep of each foot.....	2
In the great toe of one foot, and instep of the other..	1
In the great toe, instep, and ankle of one foot.....	1
In each great toe, and each hand.....	1
In the external side of the foot, and the great toe ...	1
In the external side of one foot, and the great toe of the other	1
In the heel, and the great toe	1
In the ankle of one foot, and the great toe of the other.	1
In the tendo achillis, and the great toe of the same limb.	2
In the outer side of one foot.	4
In the outer side of each foot.....	1
In the inner part of the sole of the foot.....	1
In the heel of one foot.....	1
In the heel of each foot.	2
In the heel and instep of the same foot.....	1
In the heel of each foot, in each hand, and each elbow	1

	<i>Cases.</i>
In the instep of one foot.....	5
In the instep of each foot.	3
In one instep first, afterwards each knee, each wrist, each elbow, and each shoulder.....	1
In one ankle.....	10
In each ankle.	1
In the ankle of one foot, and the toe of the other.....	1
In the ankle and instep of one foot.	4
In the tendo achillis. ...	1
In the tendo achillis, and ankle.....	1
In the thumb, knee, ankle, and great toe, on one side, and, lastly, the great toe of the other foot.	1
In the right knee and left hand at the same time.....	1
In the back of one hand.....	2
In the wrist.	1

From this statement it appears, that although podagra, as descriptive of the part affected in the first attack, contains the more general truth, yet so much exception prevails, as to render it too limited a term even to mark the first fit, as a positive designation.

I find that in the examples of hereditary gout, the great toe has been for the most part the situation first chosen, and that the most remarkable exceptions have been in those persons who have wholly acquired the disease.

The frequency of the returns of gout after the first fit, is in proportion to the constitutional tendency of the disease, and to the unfavourable mode of life of the individual. If the toe of either foot was the affected part in the first fit, the same toe is most frequently the part invaded in the second fit; but it seldom happens that the other foot escapes. The phenomena which arise, are similar in their character to those I have already de-

scribed, but are more marked in their degree, are of longer duration, and from the connexion which the local symptoms are discovered to have with the general system, the constitutional nature of the disease becomes gradually more and more manifest.

As the disposition of the constitution to gout increases, the intervals become shorter, the fits are of longer duration, and the parts affected more numerous. The gout, more than any of the other phlegmasiæ, is disposed to return at periodical intervals, and sometimes, in its annual visit, it is punctual almost to a day. One patient informs me, that for three years in succession, his attack returned on the 12th of April, and several others have assured me of a similar occurrence, almost in the same degree.

The early part of spring and the latter end of autumn, are periods during which it is most prevalent; for in these seasons, changeable weather with its attendant vicissitudes of heat and cold (the strongest of all the exciting causes of gout), is the most productive source of injury to the human frame. There is no part of the year which gives absolute exemption from the disease, when it is much established in the constitution; and summer alone is the period of expected security.

I have met with some cases in which the first fit has taken place in the middle of summer. One patient declares that his worst attacks occur in summer; and another states, that all his fits have occurred in this season, except in one year, when a second afflicted him in winter. In our English climate, this account still involves in its explanation, the exciting cause of *vicissitude of temperature*.

I must add, however, that continued excessive heat of weather is very unfavourable to gouty persons, and of

this fact I have had numerous proofs in the recent remarkable summer of 1818. Some of my gouty patients of relaxed constitution affirm, that they find themselves altogether more indisposed, in the summer, than the winter season.

The general circumstances of attack are such as have been already mentioned ; but the premonitory symptoms, and subsequent sympathies in the paroxysm, usually increase in strength with each returning fit. Yet of its notice of invasion, such is the capricious nature of this disease, that even confirmed gouty subjects are sometimes attacked at the very moment, when they most congratulate themselves on the possession of health and strength ; and I have witnessed some fits thus sudden and unexpected, to be very tedious and severe.

In a few individuals, the gout never chooses any other situation than the feet ; but in much the greater number of examples, in the course of its progress, many parts become affected in the same paroxysm ; and although the inflammation establishes itself in its different seats, in succession, it often occupies many parts at the same time ; either raging with pain in all its situations, simultaneously ; or affecting the different parts successively, with alternate violence. The feet, knees, hands, and elbows, become indiscriminately affected* ; and together with the ligaments of the joints, the bursæ mucosæ, the sheaths of tendons, and the muscular aponeurosis, in various situations, are the frequent seats of the disease.

* It is however worthy of notice, that on many occasions, the disease has a remarkable disposition to observe a certain order ; attacking, for instance, in a regular series, the right foot, the left foot, the right knee, the left knee, and so on ; or, the right foot, the right knee, the left foot, the left knee.

I have more than once seen the whole of the tendinous portion of the gastrocnemius muscle severely affected with gout. In some martyrs to the disease, not only the shoulders, but even the hips themselves are now and then affected with all the characteristic pains of the disorder, in the course of a long fit. The rapid and indeed instantaneous manner with which the inflammation sometimes quits one part, and transfers itself to some other part of the same limb, is not the least curious among the various characters of gout. As in the early attacks of the disease, so in all its returns, the night is the most common time of its invasion, and the period of the greatest suffering; yet instances are not unfrequent, in which the patient is surprised in the day by this unwelcome visitor, and often in a manner almost incredibly sudden. With some few, the pain prevails most during the day, and night brings a little soothing influence of sleep; but to others, neither day nor night scarcely gives any intermission of torture, for the first few days of an inveterate attack. One gentleman informs me, that, in a late attack, the agony which he suffered from the gout in both feet, subsided only in a slight degree in the day, for three days and nights. Another, that during thirteen days, he did not procure more than thirteen hours sleep. A third much afflicted patient describes, that a whole fortnight of the paroxysm gave him continued misery and torture, almost without the least abatement of suffering in the day; and that during the whole time he could scarcely endure the weight of the sheet upon his limbs. The gout, in this case, was situated in both feet and both knees.

But even this account is a feeble and inadequate description of the extreme distress which an urgent fit of gout affecting the upper and lower limbs, in every part, produces. Witness the disabled patient, lying in agony,

in one continued posture, unable to move hand or foot, requiring even to be fed, and reduced to every humiliating circumstance of helpless suffering!

The external appearances of the disease vary considerably, according to the situation and particular texture of the part which is affected. The redness of surface, together with œdematous swelling, are most remarkable on the great toe, on the foot, the back of the hand, the fore arm, and at the elbow; while at the ankle, knee, and wrist, the increased bulk is produced chiefly by the distension of the bursæ, and of the sheaths of the tendons, and takes place, often with little change in the natural colour of the skin. If there be redness, it appears, in these parts, chiefly in patches. In the situations before mentioned, the colour, which continues for some time to be of a scarlet hue, is diffused over a considerable extent of surface, and occasionally assumes the aspect of spreading erysipelatous efflorescence; and this sometimes in so great a degree, as to imitate erysipelas itself. When the cellular parts have been for some time swollen and tense, the blood which has stagnated in the loaded and obstructed vessels, ceases to give the vivid blush of red, and changes to the different shades of purple.

In some few instances, under my observation, the patient being corpulent and of a full habit, with a temperament partaking strongly of the sanguinous, the capillary vessels have partially given way to the force of the circulation, and the skin has been here and there, in spots, discoloured with the effused blood.

When the bursæ mucosæ are affected with gouty inflammation, they become distended, are exquisitely tender and painful, and often increase to a great size very suddenly. Sydenham observes, "Sometimes the morbid matter is thrown upon the elbows, and occasions

a whitish swelling almost as large as an egg, which becomes gradually inflamed and red." This swelling is the enlarged bursæ. I have seen it formed in the course of a night, and sometimes even more suddenly, to a great size, in the ham, near the glutæi muscles, and at the elbow as above mentioned. Sometimes these bursal distensions subside suddenly; but more commonly they continue permanent, tedious, and often untractable. This happens more particularly at the knee joint, in which situation the bursæ become exceedingly distended, occasionally without any discolouration, and bearing free pressure; but in other instances the skin exhibits much heat, redness, and tenderness to the touch, with great fulness in the surrounding vessels; and, in each case, the patient is either quite disabled from motion, or rendered very lame. At the great toe, the bursal swelling often assumes all the threatening appearances of an abscess. In a gouty hand, I have seen an old ganglion put on a similar character, and producing acute throbbing and excessive pain. The affected tendons, when carefully examined, appear to be much thickened; and sometimes seem not only large, distended, and rigid, but also inseparably matted together.

The ligament of the patella, when much affected with gouty inflammation, produces peculiar lameness, so that the patient can scarcely make the least extension of the limb without insupportable pain. A slight swelling of the part may be discovered, with great tenderness; but there is seldom more than pale redness in the skin. This appears in a small patch, and, very commonly, the integuments possess their natural hue.

In urgent and continued gout, the veins of the whole limb are preternaturally distended with blood, and, when contrasted with the healthy limb, present the appearance

of universal fulness. This state of the veins is most remarkable in the leg, but in the arm it is also very distinct. Near the part inflamed, the venous branches appear very numerous, diverging in their course, and almost bursting from fulness. The congestion which is here represented, becomes in some degree diminished, when abundant effusion into the cellular membrane has taken place at the gouty part; but even then the symptoms soon become aggravated by any attempt to keep the limb in a depending position; and if the patient endeavour to raise it even in a small degree, he feels it restrained, as if by a heavy and irresistible weight.

The disabled appearance of parts which have been long subject to the ravages of gouty inflammation, an appearance which is sometimes so distinctly marked as to resemble paralysis, is most strikingly exhibited in the hands and individual fingers.

The pain which is inflicted by gout, is much more severe than that from almost any other kind of inflammation; and the complicated sensations that result from it, are really of a nature peculiar to this disease.

The pain which the patient describes, seems to be considerably modified by the particular texture and situation of the part affected. Observation has taught me, that the sense of weight and total loss of power, are most severely felt, when the whole of the anterior part of the foot is the seat of disease; that inflammation in the first joint of the great toe produces the strongest throbbing; and that the sense of tightness is most urgent, when the elbow joint and the tendons at the wrist are the parts affected. I gather from a careful comparison of different cases, that the two last situations are, of all others, the most afflictively painful. One gentleman, who has had very sad experience in the sufferings of the gout, considers

the ham-strings as the most painful part; another, the ligament of the patella. In a fit in which the gout fixes in various situations, both old and new, any part which is for the first time affected with the inflammation, is usually represented to be the severest source of suffering. Accordingly as the affected parts are tendinous in structure, the difficulty of motion is increased, and the most careful attempts are followed by agonising pain, and sudden excruciating cramp. This occurs even when, from external appearances, a slight degree only of gout is manifested. I have often witnessed these effects in a very striking manner at the close of the fit; and in one instance most remarkably, where gout was lingering in the tendo achillis at its insertion.

Sydenham described the sense of heat experienced in the height of the paroxysm, *as that of water just warm being poured upon the membranes of the affected part* (“cum sensu quasi aquæ tantum non frigidæ, partis affectæ membranis affusæ”). This is a very inadequate account of the sensation of temperature produced by the local action of gout; which rather conveys to the patient’s mind the idea of *boiling* water; or, in the strong language of some patients, even of boiling lead. The quantity of heat which is evolved from the inflamed surface, is found by the thermometer to be always, more or less, at a higher temperature than the natural standard. The strong metaphorical terms, by which gouty persons endeavour to convey an idea of their sensations during the paroxysm, are evidences of the extreme severity of the sufferings which this disease inflicts.

In order to exemplify in a familiar and striking manner this particular point, I shall state my quotations, abridged from several cases, of the language of patients them-

selves, while describing to me their feelings during the height of the paroxysm.

J. L. aged forty-six, had *acquired* gout, first at thirty-five. In the last fit, different parts became affected in the following order: the right hand, right elbow, right toe, left knee, right ankle, left toe, left ankle. The inflammation affected several of these parts at the same time, but in general, with alternation in the severity of pain. He felt as if wedges were forced into the joints, and were effecting their separation; and a throbbing tightness, as if from a strong ligature, a burning heat, and oppressive weight, were present in an exquisite degree. Upon the smallest movement of the fingers, instant pain and spasm seized the fore arm and elbow, and when the toes were moved, the whole foot and ankle became similarly affected.

M. K. aged fifty-five; first had *acquired* gout at twenty-eight. Describes the pain, "as if a saw were at work in the joints;" has spasms on his first attempts to sleep at night, followed by agonizing sensations of cutting, pricking, and excessive throbbing; and feels the heat, as if from scalding water poured between the affected parts.

T. L. aged fifty-one; first had *acquired* gout at thirty. He compares his pain to the severe gnawing of a dog acting upon the joint, accompanied with cuttings and prickings; and with heat sometimes as from hot lead; and the sense of weight is intense. He also describes the pain at the ball of the great toe, as sometimes resembling that of a severely aching tooth.

A. G. aged forty-two; first had *acquired* gout at thirty-five. His legs, in the paroxysm, are often cold, while the affected feet burn, as if on hot coals; with great

throbbing, and piercing shootings, sometimes as if a knife were thrust in; and he can scarcely describe the intolerable weight.

J. P. aged forty-two; first had *acquired* gout at thirty-six. Expresses the feeling of heat, as if the parts were in a furnace; with pulling and throbbing; and has a sense of weight, as if a hundred weight were placed on the foot.

D. R. aged fifty-four; first had *acquired* gout at fifty. He is usually attacked about two in the morning, with violent shooting and gnawing pains in the great toe, which he compares to the sensations he felt some years ago, when the small bone of his leg was re-uniting after a fracture.

E. S. aged fifty; first had *hereditary* gout (from his father) at forty-seven. The burning pain of the affected parts, is compared to the heat of a red-hot iron; the oppressive sense of weight is as if they were covered with a mill-stone; to which are superadded distressing throbbings, with quiverings of tendons, and spasms of the muscles; and the skin feels as if girt with a strong ligature.

J. S. aged forty-one; first had *hereditary* gout at thirty-four. In his last attack in the feet, conceived that some one was boring into the joint with a gimblet; then that an abscess was forming; and at the height of his suffering, thought that he could not have suffered more even from *amputation* of the part. Cramps in the muscles of the legs and the toes, and also in the intercostal muscles, were severely distressing.

The strong expression of wrenching and tearing of the parts, as if the ligaments were undergoing forcible separation, is very commonly used by the gonty sufferer.

A lady, remarkably of the nervous temperament, de-

scribes that while the inflamed part of the foot occasions intensely burning sensations, the other parts are in the contrary state, as if resting upon cold marble.

In addition to these symptoms, the consequence of gout alone, the patient is sometimes afflicted with accompanying rheumatism, which affects either the neck, shoulder, loins, or sciatic nerve; and in some urgent cases, in which continued exposure to wet and cold had been the exciting cause, I have seen two or more of these situations under its influence, at the same time that gout has been raging in other parts. If any of the other phlegmasiæ has either preceded, or immediately accompanied the attack of gout, it most commonly soon yields to the latter disease, as I have before mentioned; but the symptoms of rheumatism, when just precursory, or co-existing, are sometimes more obstinate and fixed; although, even this complaint occasionally yields immediately to the superior influence of the gout.

The *constitutional* symptoms of the paroxysm are now to be further noticed.

In very slight attacks of the gout, it sometimes happens that the secretions do not exhibit to the eye any morbid appearance; but in severe cases, the digestive functions are evidently much affected, as is shewn by the following indications:

The tongue is furred; there is thirst and loss of appetite. The stomach is affected with flatulency, occasional spasms, and many uneasy sensations. In conjunction with nausea, and sour eructations, a watery fluid is sometimes rejected, which is very acrid and acid. It is either colourless, or of a grass-green appearance; and, in a long fit, this occurrence happens from time to time. The bowels are for the most part torpid; and, being excited by medicine, the fæces appear unusually foul and

offensive, are dark in colour, blackish, or of an olive green; or in some cases of a clay-like appearance, and often are remarkably loaded with vitiated mucus. The urine is of deeper colour than natural, is secreted scantily with relation to the quantity of the patient's drink; and, on cooling, deposits a pink or brick-dust sediment, with much mucus. Its specific gravity is much increased beyond the healthy standard.

During the most urgent symptoms of the paroxysm, it is usually passed with considerable irritation, both as to frequency and sense of heat. The pink or lateritious sediment appears, more or less, in every portion of the urine, during the inflammatory symptoms*. When these have entirely subsided, and the state of the liver (on the condition of which, the symptom in question principally depends) is still remaining unhealthy, the sediment of the urine often assumes a whitish colour, and is compared by the patient to the appearance of magnesia. This and the pink sediment frequently alternate, the one or the other appearing, as inflammatory or nervous action most prevails.

* In the London Medical Repository, the Reviewer (who has done me the honour to notice this Treatise at great length) is in error in the following criticism: "There is some degree of incorrectness in regarding the deposition of the pink sediment as one of the symptoms occurring during the continuance of the paroxysm, as it does not begin to appear until it is fairly over, or the crisis is passed." No. for Oct. 1816, p. 294. I can confidently state that the deposition of the pink or lateritious sediment *begins* with the paroxysm; provided that the paroxysm itself commences with urgent symptoms; upon which, it is the attendant.

The sensibility of the nervous system, as is shewn by what has been related of the sensations of the inflamed parts, is in a high state of morbid excitement.

Sydenham describes his pain in the beginning of the fit, with all the force of figurative expression; so unequal are the terms of ordinary language to give a picture of the sufferings of the gout. He represents the pain as that of "a dislocated bone," and sometimes as "from the gnawing of a dog." He adds, that "the membranes of the parts affected become so exquisitely painful, as not to endure the weight of the clothes, nor the shaking of the room from a person's walking briskly therein. And hence the night is not only passed in pain, but likewise with a restless removal of the part affected from one place to another, and a continual change of its posture. Nor does the perpetual restlessness of the whole body, which always accompanies the fit, and especially in the beginning, fall short of the agitation and pain of the gouty limb."

Some patients inform me, that in their most painful fits, they suffer still more distress from nervousness and general irritation, than from the pain itself.

In examining the symptoms of one hundred and twenty cases of gout, which I have recorded in minute detail, I find that in ninety of the whole number, *cramps* are mentioned, more or less strongly, among the sufferings incidental to the gouty constitution; and, almost without exception, are described as being urgent either just before the paroxysm, or during its height, or just at the close; and with some persons, at each of these periods.

The muscles situated on the thigh and leg are the most frequently affected; but those which move the toes,

and fingers; the diaphragm; the muscles of the chest, the abdomen, and even of the ribs, are not spared in this painful affection. Almost any change of posture serves to produce the spasmodic action, when the disposition to it prevails. I have seen a patient instantly seized with cramp on the most careful attempt to extend the leg, when the gout was present; the gastrocnemii muscles becoming afterwards visibly trembling and convulsed. One person expressed, that, with every spasm, he felt as if painfully electrified over the whole frame. Some describe the agony of the cramp as being the most painful suffering which they experience, and declare that they think the duration of it beyond the one or two minutes to which it does continue, would be wholly insupportable. A jerking of the limbs, and excessive restlessness, in the early part of the night, may be added to the distressing list of symptoms.

A general febrile action usually accompanies the local inflammation. This febrile action is symptomatic, and, together with the pain and increased disorder of the affected parts, returns at night, and remits when the local symptoms become relieved in the morning; but when the local symptoms do not remit, the febrile irritation is also constant. In the evening, and sometimes in the day, the patient complains of transient chills, affecting the frame generally; or, in a more partial manner, they run in the course of the spine alone. Some suffer this symptom, which occasionally amounts to a strong rigor, together with much lowness of spirits, on first going into bed: and it is followed by an irregularity in the distribution of the animal heat; one part having the sensation of dry and burning heat, another of *marble* coldness. When a strong inflammatory diathesis is present, or after an excessive application of some of the exciting causes,

the action of the heart and arteries is violently and permanently increased; and, universal heat of skin with the usual symptoms of pyrexia are proportionally urgent.

Painful hæmorrhoids, with occasional discharge of blood, sometimes accompany the gout. The symptoms which mark the favourable decline of the paroxysm, will be described under the head of *Prognosis*.

The same internal causes which support a tedious and harassing continuance of the fit; or, as Sydenham has well expressed it, "the chain of fits," also often induce a severe relapse, at a moment when the patient flatters himself with the expectation of recovery. In this condition of the constitution, a relapse may follow a slight application of any of the remote causes, till weeks, or even months, have elapsed.

The restoration of general health and vigour in some instances precedes the recovery of the limbs; but in others, is delayed till long afterwards: and, where proper treatment has been neglected, in old and violent gout, the lameness and tenderness of the affected parts continue very permanent; the patient cautiously chooses the most even ground for his walking exercise; and still, with every care, his foot sometimes turns suddenly as if it were dislocated; producing exquisite pain.

With respect to the length of the interval between the paroxysms, Sydenham has observed, "According as the immediately preceding fit has been more or less severe, the following fit comes on in a longer or shorter space of time; for, if the last fit proves violent, the next will not attack the patient till the same season of the year returns again."

This observation is not without its share of truth; but it may be remarked, that the benefit of a longer respite is in this manner dearly purchased; nor is the ad-

vantage certain. A long and violent fit is often succeeded by another of equal magnitude, in the same year. Some relate that when they have enjoyed an unusual interval from a fit, any slight cause will serve to excite an attack.

Finally, it may with little exception be stated of the gout, that it acquires strength with each returning fit, both as to the number of parts which it attacks, and as to the duration and degree of suffering; and that it does not, like some chronic diseases, wear itself out by repetition, and yield to the friendly power of time. Both constitutionally and locally, also, the susceptibility to the disease increases. A premature old age comes on; and, together with crippled and painful limbs, the nervous system is so enfeebled, that both mind and body grow less equal to sustain the conflict.

Such is the melancholy but faithful portrait of the disease, when it is permitted to pursue its *natural course*; —such is the sad and certain tyranny of *neglected and encouraged* gout!

SEQUELÆ OF GOUT.

The sequelæ, or occasional consequences of acute gout, may be divided into those which are *constitutional*, and those which are *local*.

The chronic form of the disease is of frequent occurrence.

The same state of system from which gout has been much produced, sometimes give rise to apoplexy, and paralytic affections; and very usually induces habitual

cramps, and much nervous irritation and wandering pain.

A diseased condition of liver, sometimes proceeding even to a material change of structure, belongs to the train of gouty evils; for seldom indeed is the action of this organ healthy, either before or during the paroxysm. In two very gouty females, I have met with chronic splenitis. Such, at least, was my view of the disease. For the reasons already stated, hypochondriasis is a common consequence of frequent gout.

The stomach often becomes permanently weakened; or even where it continues in due vigour, as far as is judged of by the appetite, it is found that an impaired state of the digestive functions, with irregular action, and generally a sluggish condition of the bowels, are almost the certain consequences which follow a long series of gouty attacks.

Irritation of the urinary organs, and gravel, occur rather before and during the paroxysm, than in the interval; and according to my experience, calculus of the bladder is a very unfrequent complaint amongst gouty persons. The contrary opinion appears, however, to prevail. *Sydenham*, after enumerating the effects of the disease, "in the pain, lameness, inability to motion of the parts affected, the sickness, and other symptoms," adds, "the gout breeds the stone in the kidneys in many subjects; either because the patient is obliged to lie long on his back, or because the secretory organs have ceased performing their proper functions; or else because the stone is formed from a part of the same morbid matter; which, however, I do not pretend to determine. But from what cause soever this disease proceeds, the patient is sometimes at a loss to know whether the *stone* or the *gout* be most severe."

*Morgagni** relates the case of an arthritic patient, who had also *nephritis calculosa*. He died apoplectic. On examination after death, it is stated, "The kidneys were found larger than they naturally are, but the right was the largest; so that, together with the fat whereby it was covered, it was almost equal to the size of a man's head. Therein were eleven stones; most of them large and ramifying. In the left was one, and that also was ramified and small. They resembled, in colour and branches, black coral." Morgagni, in another part of the same Epistle, observes, "The calculi of the kidneys are often joined with the gout."

In investigating the question of the occurrence of the calculus in the bladder (the disease which we call the stone), in those subject to gout, I find that among two hundred and thirty-one persons, five only were so afflicted. Of the gravel, I shall again speak more particularly.

The local changes produced by the gouty inflammation are various. The cartilages, in aggravated cases of gout undergo erosion, and consequent loss of function in the joint, more or less complete; or, in the worst instances, they become entirely removed by the subsequent action of the absorbents. When this has happened, a vacancy is discovered at the joint, very visible in the examination of the living subject. In the hand and feet these changes most occur; and then the fingers and toes are distorted in different directions, quite at the disposal of the muscles.

The synovial secretion frequently fails in its healthy quantity and quality after deep-seated gouty inflamma-

* Ep. lvii.

tion; whence arises stiffness and lameness of the joint, with snapping, and an indefinable sensation of dryness in the parts, upon motion.

A gentleman, who suffered gouty inflammation in the ankle, six months ago, says, that in addition to much weakness of the part, he has often the sensation as if there were gravel within the joint.

The ligaments become thickened, shortened, much deprived of their elasticity, and habitually tender.

The bursæ mucosæ acquire a state of permanent distention, and are sometimes enlarged to a great size; being soft and yielding to the touch. Of the small bursæ, the contents occasionally become so much inspissated, as to constitute hard, solid tumours, which, in some instances, are with difficulty distinguished from bone itself. The bursal disease over the first joint of the great toe, which is familiarly known by the name of *bunion*, is a very common complaint with gouty persons. The deep-seated and immediate textures of the joints, which have now been described, in some lamentable instances are so completely disorganized, that the most confirmed and irreparable lameness ensues. The parts appear entirely consolidated together, and are as immovably fixed as in bony ankylosis itself. I have witnessed one remarkable case in which the ankle joint in each limb was distorted to such a degree, as to give the appearance of club feet, with still greater lameness from the tenderness of parts.

The secretion from the sheaths of the tendons is also vitiated in a similar manner; and hence arise the thickened knotty feel of gouty tendons; their hardness, contraction, and rigidity.

The muscular aponeurosis sometimes becomes thickened and contracted from gouty inflammation; and the muscles themselves apparently become shortened from

the influence of frequent spasm. This morbid change in the aponeurotic texture, in a great degree restrains the moving powers beneath it, from their necessary freedom either of flexion or extension. In one patient, who has suffered repeated severe gouty inflammation in the gastrocnemius muscle of one leg, I find such a state of *board-like* hardness, as to conclude, that adhesion has taken place, between the aponeurotic texture and tendinous fibres of the muscle, and the covering integuments.

A varicose state of the veins of the legs causing habitual sensations of achings, fulness, and heat, not unfrequently occurs in old gouty persons; and is occasionally accompanied with purple cutaneous spots (blotches); and, though more rarely, with ulcerations of the skin. In the legs of three gouty men, advanced in years, I have seen *icthyosis* existing in a very great degree.

Dr. Munro, jun. observes*, "The extremities of the bones of the feet and hands have been said to be occasionally converted into a white substance like chalk, in persons who have long been afflicted by gout." From my own examination of the parts in the living subject, and in morbid preparations, I am induced to think that the bones do not suffer any morbid change of structure from the influence of gout; nor that this happens even to the periosteum. It has not appeared to me, that the gout fixes so deep as this texture.

In the interesting case of dissection of a gouty person by Mr. Watson, the following account is related: "On the middle of the right tibia, there appeared an oblong tumour resembling a node; over which, the integuments were very thin and ready to burst. It was a mere dc-

* Outlines of Anatomy, vol. i. p. 154.

position of chalk-like matter, between the skin and periosteum, and, though thick and large, had not as yet done any injury to the bone." The author also states, "One of the great toes was found to be much enlarged; and, upon dissection, the first joint of it was found to be enclosed in a bed of chalky matter like a fossil shell; but the bone itself was neither increased in size, nor altered in its texture*."

The gouty concretions (commonly called *chalk stones*) occur only in a few individuals of particular gouty idiosyncrasy. They result from an inspersion of the peculiar morbid secretion which constitutes their composition, and are found in various situations from within the synovial membrane of the joint, even to the layers of the cutis. I have found them in the living subject, filling the bursæ, and condensed to great hardness; in the sheaths of tendons, feeling almost stony; in the cellular membrane, either in hard or soft lumps; and under the cuticle, pressing for escape.

In one gouty person who comes under my frequent observation, the concretions near the surface have caused numerous ulcerations both in the hands and feet, and the chalk-like matter is constantly secreted. I have met with many similar cases of less extent.

In three persons who have gouty concretions both in the hands and feet, I have found a remarkable state of hardness in the nails of the toes and fingers; possessing an extraordinary brittleness; fragile, and scarcely capable of being cut.

In very confirmed cases of concretions in the hands and feet, the fingers and toes present a sad spectacle of

* Medical Communications, vol. i.

disorganization. When the deposit is external to the joint and contained within the bursæ, or the thecæ of the tendons, the finger or toe is stiff almost to the degree of a perfect ankylosis. When within the capsular ligament, ulcerative absorption takes place; the cartilage becomes removed; and one or more of the phalanges receive a distorted position from the action of muscles. I have seen from these causes, a finger and a toe drawn across the others in a completely transverse manner; and distortions in a less degree are always found in cases of this description.

In the Hunterian Museum at Glasgow, I examined several preparations, shewing the changes of structure caused by the concretions of this kind; and I here offer the written account of them*, which I copied.

“ L. L. No. 26. s. A finger from a gouty hand; a joint opened, and bent upon itself, to shew the cartilages eroded a little.

“ 27. Ditto very much eroded, the joint full of chalk-like substance.

“ 27. a. s. Ditto, the joint invested every where with a thin layer of chalk.

“ 28. s. Thumb from the same hand, shewing the same effect.

“ 29. s. Another thumb, shewing the same.

“ 29. a. s. Ditto.

“ 29. b. s. Another thumb; the chalk accumulated about the joint in considerable quantity.

“ 30. The account omitted.”

* The statements are chiefly in the hand-writing of Dr. Baillie.

I am favoured by my friend Mr. Brodie, with the following clear and interesting particulars of the dissection of a gouty person.

“*External Appearances.*—Several joints of the fingers were ankylosed, and the fingers variously distorted. The middle finger of the left hand was shorter than the rest, and the skin over it was loose. The bone of the second phalanx appeared to have been nearly absorbed, so that there were scarcely any remains of it; and there was only a small quantity of soft substance in its place.

“The right wrist and elbow were ankylosed, as were also several of the joints of the toes. The knees admitted of incomplete flexion and extension, and the motion of the joints was attended with grating.

“In various parts of the body there were orifices in the skin, communicating with membranous cysts, situated in the adipose substance, and discharging a chalky fluid of the consistence of cream.

“*Dissection.*—The pleura pulmonalis and costalis were universally adhering. The stomach, spleen, liver, and gall bladder, adhered universally to each other, and to the contiguous parts. The gall bladder contained two large gall stones. There were no other preternatural appearances in the thorax or abdomen.

“There were no remains of the cartilages of the left knee. The corresponding parts of the patella and condyles of the femur were worn into grooves and ridges, evidently from friction on each other: presenting, however, a compact surface, and not exposing the cancellous structure, as would have been the case, if friction had taken place in the dead body. A thin layer of white chalky matter was deposited on the bones, where the cartilages had disappeared, in several places. On the

margin of the articulating surfaces were several small exostoses. The ligaments and the synovial membrane were in a natural state, with the exception of the thin layer of the latter which is extended over the cartilages, and which had disappeared like the cartilages themselves.

“ In the right wrist, the first row of the carpal bones were united by long ankylosis to each other, and to the radius.

“ The other joints were not examined.”

“ The patient was an elderly female, who had been subject to severe attacks of the gout for many years, and who was supposed to have died of a cancerous affection of the stomach.”

Of the active influence produced on the absorbent vessels by the stimulus of gouty concretions when deep seated, the following account of a case, which has been obligingly sent to me by Mr. Howship, further serves as a demonstration.

“ A female, aged 43, had for many years suffered severely from gout. The attacks of the disease, at first confined to the first joint of the great toe, subsequently affected the articulations of the fingers, the wrists, and the heels; ultimately the hands were almost covered with tumours, the action of the joints of the fingers being all either much impeded, or quite destroyed, by the repeated effusions of chalky matter.

“ For the space of three or four years the disorder manifested itself principally by very extensive, painful, and tedious inflammation of the heel, which ulcerated, and threw out a copious and long continued secretion of chalky as well as purulent matter; and no sooner was the one heel recovered, than the other became affected in the same way.

“ In 1815, this patient, still confined to her bed, and

totally helpless from the inflamed and ulcerated state of the heel of the left leg, experienced a very smart attack of erysipelatous inflammation upon the right leg; and when this began to give way, violent inflammation took place lower down at the ball of the great toe. The integuments covering the joint soon sloughed out, and exposed a deep ulcerating surface. A little dry lint covered with a warm poultice, was the only application made to the parts, which were examined daily. About a fortnight after the separation of the slough, some white matter made its appearance at the bottom of the ulcer, and was removed with the dressings. A few days after this, a mass of soft chalky substance as large as a pea was seen, and was turned out with the end of a probe. In the course of a week, I observed that the same appearance had again taken place, and that the chalky secretion was deposited in exactly the same part of the ulcer as before; and it seemed to be deeper seated than the surface of the head of the metatarsal bone. I therefore made a particular examination with a probe, and thus ascertained that part of the surface of the bone was removed; the margin of the opening being readily felt all round among the granulations. Within this space, there was a depression answering to the form of each of the small masses of chalk that had been previously removed. The ulcer, even within the cancellated texture of the bone, looked tolerably well; it was covered with granulations, and secreted healthy pus.

“ This peculiar secretion continued for some time; a fresh deposit of compact chalky substance being protruded, and removed from the ulcerated space within the bone, every few days, for several weeks. At the termination of this period, the action lost its specific character, and the wound healed.

“ This unfortunate patient, whose case I have already adverted to, in my *Practical Observations in Surgery*, as exemplifying the symptoms produced by gouty affection of the brain, eventually fell a victim to the disease.

“ The above case affords an instance, in which, the capillary arteries within the cancellated structure of bone evidently took up the specific action of gout, by secreting the chalky matter peculiar to that disease. It appears to me, that the irritation from the inflamed surface, had, in the first instance, considerably disturbed the circulation upon the membranous expansions lining the cancellated texture of the affected part of the bone, increasing the activity of its powers, both in secreting and absorbing. In the papers published in the *Medico-Chirurgical Transactions*, on the Formation, Structure, and Diseases of Bone, I have already stated, that the first effect of a certain degree of irritation falling upon the membranous sheaths and expansions within bone, is a change in the state of the membranes, which, from being thin and transparent, become granulated and opaque, from increased vascularity; and that absorption of bone, more or less extensive, is one of the earliest consequences of this change.

“ The circumstance of the bone having been in the present instance partially removed by absorption, argues no new rule for the treatment of such cases; but the secretion of the urate of soda by the capillary arteries within the bone, affords a very curious, and at the same time a very interesting proof, that the circulation within the ossific parts of our fabric, is not governed by laws differing from those that regulate the soft parts of the body; being equally subject to feel the effects of any transient impression, dependant on accidental surrounding circumstances.”

With respect to the ossific changes which were pro-

duced in these cases, and which, in the first, were so numerous and remarkable, it appears to me necessary to consider them as wholly secondary; and, as being an effect of ulcerative absorption, occasioned by the stimulus of the uric compound confined within the capsular ligaments, and acting as a foreign body. It would seem to be more easy for Nature to remove by absorption the bones pressed upon and resisting this deposition from the morbid action of the extreme vessels, than to exert her *vis medicatrix*, in putting a stop to this action of vessels, when there should be too little space for the bone and the secreted matter, both to be contained. On the subject of these concretions, I shall have occasion to remark further, at a future page.

THE REMOTE CAUSES OF GOUT, PREDISPOSING AND EXCITING.

ALTHOUGH I shall treat of these different causes separately, it is obvious that the distinction is in some degree artificial. The accidental and stronger application of some of the predisposing causes, as, for example, diet, strong liquors, &c. is sufficient to render them exciting causes; and the effect which is thus produced, will be in relation to the state of the system; and will be proportional to its readiness to take on the gouty action.

PREDISPOSING CAUSES.

Hereditary Predisposition.—By Dr. Cullen, and by writers in general, gout has been too expressly defined an *hereditary* disease. In conformity with this character,

the offspring of the gouty parent should seldom escape from having the disease; and still less, should we find, that in the list of its subjects, the examples of acquired gout, *not* hereditary, form the most numerous class. With a view of obtaining accurate information on this subject, I instituted the following comparison:

Of two hundred and thirteen patients, the number of those in whom the disease was

Hereditary from the father, was.	62
————— mother,	29
————— father and mother,	14
Of those whose grandfather only had gout,	14
————— grandmother only had gout,	1
————— grandfather and grandmother ..	1
————— uncle only in the family had gout, ..	7
————— mother's uncle,	1
————— aunt only in the family had gout, ..	2
Not known either on the father's or mother's side,	84

215

From this statement it appears, that the cases of acquired gout, in which no family reference could be traced, were to the rest, as 84 to 131; and in the examples contrasted with those immediately hereditary, as 124 to 91.

Morgagni mentions his own gout as original. Heberden observes, "I have known a female who suffered by the gout, to the degree of having numerous sores from chalk stones, though it had never been heard of among any of her relations." One patient, a butcher, who cannot trace gout to any former generation in his family, informs me, that he and his three brothers suffer the dis-

ease in a very severe degree. He has two sisters, one of whom is much afflicted with gravel.

Cadogan asserted of the gout, that "it is not hereditary." In the fair acceptance of the expression, this is inaccurate; but still it comes nearer to the truth than the opposite and more prevailing doctrine. I speak of hereditary disease, as one depending on similarity of organization between the parent and child, although it is of too delicate a nature for the observation of our senses, and for direct detection. In the same manner as the natural functions of the body depend on particular structure, so we must conclude to be the case with regard to many diseased actions, which are of a specific nature; such as scrofula, cancer, idiopathic phthisis, or *hereditary gout*.

In my former observations on this part of my subject, I too hastily concluded, that the tendency of the gout to appear early in life, and severe, was much the most strongly manifested in those who possessed the hereditary disposition to the disease. I have now put this question to the test, in examining the cases of one hundred and eighteen persons under my own care, who have had gout in a very severe degree; leaving out of the comparison, all the other cases which I have seen, of a mild description. From this estimate, the following result appears:

In Men, Acquired.

At the age of 18,	1
Beginning between the 20th & 30th year,	15
_____ 30th & 40th year,	19
_____ 40th & 50th year,	9
_____ 50th & 60th year,	1
	<hr/>
	45
	<hr/>

In Men, Hereditary both from Father and Mother.

Before the 20th year,	2
At the 32d year,	1

In Men, Hereditary either from the Father or Mother.

Beginning between the 20th & 30th year,	12
————— 30th & 40th year,	23
————— 40th & 50th year,	7
At the 62d year,	1
	<hr/>
	43
	<hr/>

In Men, the Gout having existed more remotely in the Family.

At the 22d year,	1
—— 24th year,	1
—— 25th year,	1
—— 32d year,	1
—— 34th year,	1
—— 40th year,	1
—— 43d year,	1
—— 48th year,	1

In Women, Acquired.

At the 18th year,	1
—— 20th year,	1
—— 26th year,	1
—— 34th year,	1
—— 40th year,	2
—— 46th year,	1
—— 57th year,	1
—— 60th year,	1

In Women, Hereditary either from the Father or Mother.

At the 36th year,	1
—— 37th year,	1
—— 39th year,	1
—— 47th year,	1
—— 52d year,	1
—— 64th year,	1

In Women, the Gout having existed more remotely in the Family.

At the 40th year,	1
-----------------------------	---

In reading over my details of the cases of men, here alluded to, I find the examples of highly inveterate symptoms to have been on the side of those whose parents on either side never had gout, contrasted with the hereditary, as 36 to 28. Of the female patients, the comparison affords six on the hereditary list, four on the acquired.

The proportionate number of cases of *extreme severity*, in women, is, in this comparison, on the side of hereditary gout. This part of the question appears to me very fairly determined, as it respects male gouty patients; but a larger series may be thought necessary, as it relates to females. I must retract the following observation which I formerly advanced: "It very seldom happens that a female has the gout, whose parents have been free from it." The present comparative view is in contradiction of this position. Of the rare occurrence of the disease among women, as contrasted with its frequency among men, I shall make a separate consideration.

Dr. Adams (whose loss we have now to lament), in his late Treatise*, has drawn a distinction, which appears

* On the supposed Hereditary Properties of Diseases, &c.

to me not very well founded, between the disposition and the predisposition to a disease. He attaches the strongest signification to the former of these expressions (for they are only expressions); a signification, which as the word is compounded, is surely rather due to the latter. The epithets of strong and slight, in connection with either expression, would, I conceive, make the distinction sufficiently clear and marked. The author, after having defined his terms, observes, "If it were true in all as it is in most cases, that the habits of the sedentary and healthy are necessary to induce the gouty action, there could be no question, that it is only hereditary in predisposition; but in some, the susceptibility to gout is so strong as to require no other stimuli for inducing the action, than such as seem absolutely necessary for the support of ordinary health. In *gout*, therefore, we must admit the two degrees of susceptibility, *disposition* and *predisposition*; nor will it be often difficult to fix their exact limits." In his fundamental arrangement of the subject, the following view is offered: "Diseases either appear at birth, in which case they are called *congenital* or *connate*; or they arise afterwards. The first only can with propriety be called *hereditary* or *family diseases*; all others we should consider as *hereditary* or *family susceptibilities* to certain diseases." This distinction seems both judicious and necessary; but the reference in each case, must, I apprehend, be made to *structure*. For myself, I confess, that I cannot form any satisfactory notion of hereditary quality, either healthy or morbid, which is not founded on structure. With respect to gout, we see in numerous instances, that the parents, either father or mother, of one or more children who when adult have sooner or later become gouty, had not themselves the disease until many years after the birth of such child

or children. The disease is not the less hereditary in the legitimate meaning of the term, because the susceptibility which is transmitted from the parent to the child, appears at an advanced period of life, instead of being *connate*, or instead of occurring before the age of puberty; and with regard to gout, it seems, that at this early period of growth, the actions of the system oppose its production. By way of illustration, though the analogy be but faint, I may make reference to the personal resemblance which a child acquires as he grows up, to his father or mother, or more distant relations, and which is not manifested at the infantile age. A gradual developement of function, in connexion with the constitutional tendency to particular diseases, is matter of daily and familiar observation; and equally implies, as it appears to me, a similarity of structure, brought to maturity by years.

With respect to *family* disposition to gout, and its dependance on hereditary influence, or upon other causes only, the comparison, which I have made, presents the following results. I have found as many of the individuals of a family affected with the disease, and with as much severity, where it has not been known either on the father's or mother's side, as in the opposite instances, in which, either one or other of the parents has been gouty. In one family, three brothers and a sister, out of six children, have been severely affected; in another, also consisting of six, four brothers have suffered from very aggravated gout, and the two sisters only have escaped. In these examples, the disorder was not known in the two preceding generations. A female, whose aunt only in the family had gout, states that with herself her three brothers had the disorder severely; her sister was free from it. On the other hand, in one family, the father and his two sons and a daughter, out of five children, have had

severe gout; in another, the father, mother, and all the sons and daughters, four in number, have been afflicted. Its confinement to one or two only in the number of a large family, is also to be noticed. Thus, out of fourteen children, but one brother and one sister had gout; and this was acquired without any share of hereditary influence. In another family, of ten children, the father had gout, and one son only was affected; but he very severely. Again, among nine children, six sons and three daughters, the father being gouty, one son only was affected. Such is the variation which prevails; and in the endeavour to discover a general rule of conclusion, I can only say, that, *cæteris paribus*, those individuals of a family, whether either parent may have had the gout or not, will be the most liable to the disease, whose bodily conformation, constitution, and temperament, and particular habits of living, shall be most conducive to that particular state of the system, out of which I consider the gout to arise, and of which I shall treat in regular order. But, before I take leave of the subject, I must remark, that in the case of one or two only out of a large family of children being affected, although one or other of the parents has had the disease, I have in two or three instances distinctly found the personal resemblance to the gouty parent, to be much stronger in the gouty child than in any of the other children. When *both* parents have gout, we may expect the disposition to the disease to be manifested in *several* of the children. I have not yet met with an exception to the fact. Much, therefore, is certainly to be assigned to the influence of hereditary claim; but authors have hitherto greatly mis-stated the question.

The Adult Age.—The exemption of youth from the gout, is a striking character of the disease. As some explanation of this phenomenon, it appears to me, that

during the rapid growth of the body in the early years of life, that plethoric state of the system, which is connected with more or less of congestion in the system of the vena portarum, and upon which, I consider, that a first fit of gout in particular, so much depends, does not take place. The diseases which occur below the adult age, affect more particularly the general arterial system. It is true that in youth, there is an absence of many of the strong predisposing causes of gout, but not to the extent of serving as a sufficient explanation of the point in question. The gout does not usually take place, until many years after the full and continued operation of all the indiscretions and irregularities, which belong to the early life of the adult.

I am persuaded that the commonly-asserted cases, which represent the existence of the gout in very early youth, are really examples of rheumatism. Some gouty persons have given me confident assurances, that their first fit took place at fifteen; and one gentleman declares, at seven years of age. Another gentleman, whose mother had gout, narrates, with such precision as to demand my assent to the fact, that he was attacked in the great toe at eight years of age. He underwent the operation of lithotomy at twelve. I am sceptical as to the accuracy of most of such statements, at the same time that I admit the occasional exception to the general rule, in the occurrence of the gout between the fifteenth and twentieth year; but at an earlier period, it must be held as a singular phenomenon.

Sydenham remarks, "I have not hitherto found children, or very young persons, affected with the true gout."

Hippocrates has an aphorism on this point, which may be thus translated: *Puer non laborat podagra, ante veneris usum.* Aph. xxx. sect. 6.

Heberden says, "I never knew a certain instance of the gout beginning before the age of puberty*. Sydenham, in speaking of the gout in females, is erroneous in stating that it "attacks only aged women." The disease does not certainly make its invasion so early in the female sex as in men; but it seldom waits the period of very advanced years; and I know one female who was attacked with gout at the ball of the great toe, at the age of eighteen. The fit was of tedious duration, and irregular in its progress. The disease was not known in her family; but in her structure, temperament, and some of her habits, there was abundant illustration of several of the causes of predisposition, of which I am now treating.

The following tabular view exhibits the period of the first attack in two hundred and nine cases:

At the age of eight	1
————— seventeen	1
————— eighteen	2
Between twenty and twenty-five } years of age. }	25
———— twenty-five and thirty ..	38
———— thirty and thirty-five	41
———— thirty-five and forty	37
———— forty and forty-five	18
———— forty-five and fifty	25
———— fifty and fifty-five.	11
———— fifty-five and sixty	5
———— sixty and sixty-five.	3
At the age of sixty-six	2

* Commentaries, p. 33.

I have not myself witnessed more than one example of a first attack before twenty; nor any after sixty-six.

Particular Bodily Conformation.—It may be observed of gouty persons, that, for the most part, they are formed with a capacious and circular chest; and that they have large full veins, and loose solids. Even in those examples, in which the veins are small, they usually appear very much distended, when the gout is approaching. In relation to stature and bulk, I have made a comparison in two hundred and fifty-four examples, the results of which are as follow:

	<i>Males.</i>	<i>Females.</i>
Tall and corpulent.	64	9
Short and corpulent	25	8
Middle height and corpulent.	41	4
Middle height and slight	9	0
Middle stature and bulk	28	1
Tall and middle bulk	14	1
Short and middle bulk	21	1
Tall and slight	8	0
Short and slight.	16	4
	<hr/> 226	<hr/> 28

This statement confirms the observation of Sydenham, that the gout chiefly attacks the gross and corpulent; but his opinion, “that such as are liable to this disease have large heads;” and Cullen’s, that it attacks especially “men whose skins are covered with a thicker rete mucosum, which gives a coarser surface,” I cannot, from attentive investigation, discover to be founded in truth.

Of the twenty-three persons classed in the above cal-

culature as short and slight, or of middle height and slight, seventeen had *hereditary* gout.

Constitution and Temperament.—I believe it may with truth be affirmed, that the gouty in general possess good constitutions, *abused* by habits of indulgence; and hence the familiar adage, “Gout is the disease of those who will have it.” A corpulent state* of body usually precedes the invasion of the disease, and in many persons increases with its progress; but when its miseries are frequent and severe, loss of flesh, either partial or general, becomes a certain consequence; and this sometimes proceeds even to emaciation. On the *temperament* of gouty persons, Dr. Cullen remarks, “If, with the ancients, we might ascertain, by certain terms, the temperament of men, I would say, that the gout attacks especially men of a *choleric-sanguine temperament*, and that it very seldom attacks the purely sanguine or melancholic. It is, however, very difficult to treat this matter with due precision.” I entirely accord with the author in the concluding part of this quotation. We should, however, aim at greater precision in our terms, in relation to the subject of temperament, than is met with in the medical nomenclature even of the present day. The notions of the ancients on this point, together with their general opinions on disease, were so much obscured by the extreme doctrines of the humoral pathology, and so much blended with fiction and wild hypothesis, that in our practical language we necessarily cease to employ their terms; yet they are still retained in books, and without any method of improvement being attempted.

* Corpulence is often such an excess of health, that it is really the beginning of disease.

The term temperament, as an expression for the particular character and bias of the living actions, marked in the original stamp of the constitution, and more or less evinced through life, is convenient in use and forcible in meaning; and at the same time that we reject a few of the ancient epithets, which were applied to the supposed morbid states of the blood and secreted fluids*, we may preserve some arrangement on the subject with every advantage.

My present limits permit me only to be brief; but I would be tempted to propose a simple distinction into two kinds of temperament; the sanguineous, as designating that peculiar to persons in whom common inflammation is easily produced; and the *nervous*, as applying to those constitutions in which inflammatory action is excited with difficulty, but in which a morbid sensibility of nerve distinctly predominates; while the combination of the two, in different proportions, may be denoted by the expressions, sanguineo-nervous, or nervo-sanguineous. Particular morbid tendencies I would express by the ancient term *diathesis* (διαθεσις), signifying disposition, as gouty diathesis, bilious diathesis, dropsical diathesis, &c. The term *melancholic*, though physically to be referred to the structure of the nervous system, may be applied with propriety to the mental disposition; and to this term may be contrasted that of *lively*, that is, possessed of vivacity. The terms *irritable* and *nervous*, as applied to the nervous system at large, are very synonymous in

* Hippocrates conceived that there were four humours in the body; blood, bile, black bile, and phlegm; and hence, accordingly as one or other of these humours was considered to abound in an individual, he was said to be of a sanguineous, bilious, melancholic, or phlegmatic temperament.

their usual acceptation; but probably it would be correct to apply the term *irritable*, to the morbid sensibility of nerve which is associated with quickness and vehemence of temper, and energy of mental action; and that of *nervous*, to the association of a *similar* state of nerve, with irregularity of spirits and more timidity of mind.

Gouty persons exhibit in the full and dilated state of their veins, strong marks of a stimulated circulation connected with weak vessels: and I concur with Dr. Cullen in the opinion, that they possess a mixed temperament. Even in the acute state of the disease, they can employ powerful stimulants (although such stimulants are always improper), without the same degree of injury, or the same certainty of producing it, as would follow in most kinds of inflammation. They are usually highly irritable and nervous, not only from being so rendered by the sufferings of the gout; but for the most part, from the influence of their original constitution.

The Male Sex.—The comparative infrequency of the gout in *females*, naturally suggests the inquiry—Whence does the male sex derive its peculiar predisposition to the disease, if its greater frequency in them really arise from a stronger predisposition? But if not, what other causes for this fact can be assigned? The more common occurrence of this disease in men than in women, must, without doubt, be principally referred to the chief remote cause, excess in living, and especially excess in wine, being applied in a greater degree by the former. But in addition to this circumstance, the superior delicacy of the female structure and habit, puts some restraint on the acquirement of the inflammatory and plethoric state of vessels which appertains to gout. The actions of the uterus are not without effect in counteracting a general

redundancy of blood*. A gout of imperfect development, or of a chronic form, is more common in women than in men. It is of unfrequent occurrence in the former, unless with the concurring influence of strong hereditary predisposition. In the few exceptions which do occur to this general rule, we meet with the circular chest, large full veins, relaxed solids, and tendency to corpulency, which have been already described as prevailing in gouty men.

Station of Life, and Occupation.—Sydenham, speaking of the calamity of the gout, says, with equal modesty and beauty of expression, “But what is a consolation to me, and may be so to other *gouty* persons of small fortune and slender abilities, is, that kings, princes, generals, admirals, philosophers, and several other great men, have thus lived and died. In short, it may in a more especial

* Hippocrates gives an aphorism on this subject (Aph. xxix. sect. vi.), which is usually thus rendered. *Mulier non laborat podagra, si non menses ipsi defecerint.* In the Medical and Physical Journal, No. 214, the Reviewer has called in question the accuracy of my interpretation of this aphorism. He attaches to the word *defecerint*, very improperly as I think, the literal signification of *deficient*, considering it as a comparative expression. To me it appears clear, that the Greek word in the aphorism *εχλιπη* signifies *dereliquerint*, or having left entirely. Those females who suffer gout are such as are very liable to *mænorrhagia*, and have the gout either at the early part or about the middle of life; or those who have been so liable, and have the gout come on a few years, sooner or later, after the final cessation. It is at the latter period that the gout most occurs with females; and to this, I think, the aphorism is intended to apply.

manner be affirmed of this disease, that it destroys more rich than poor persons, more wise men than fools; which seems to demonstrate the justice and strict impartiality of Providence, who abundantly supplies those that want some of the conveniencies of life with other advantages, and tempers its profusion to others with equal mixture of evil; so that it appears to be universally and absolutely decreed that no man shall enjoy unmixed happiness or misery, but experience both: and this mixture of good and evil, so adapted to our weakness and perishable condition, is perhaps admirably suited to the present state*.”

Since the period at which these sentiments were entertained, luxury has so much increased among the whole community, that the gout sometimes finds entrance even into humble dwellings. In London, amongst the inferior classes, I have observed butchers, innkeepers, butlers, and porters in wealthy families, to be very subject to gout. It is also frequent amongst coachmen, chiefly such as live in families; for, together with much excitement from liquors, they are constantly exposed to the variations of the weather. In short, such stations and occupations as most induce repletion and inactivity, or full living with only passive exercise, lead to gout; and even in some constitutions, in which there is great tendency to plethora and corpulency, moderate indulgence in diet, notwithstanding active exercise, seems to implant the disposition to the disease.

* In a similar spirit of moral reflection, the Poet happily expresses himself:—

“ O fortunatos nimium, sua si bona norint

“ Agricolas!—

Georg. II. 458.

State of Mind.—The influence of the passions on the bodily health we know to be at all times powerful; but as far as respects the gout, I think that the depressing passions have the strongest effect. Grief and anxiety impair the healthy energy of the brain, primarily; and affect the digestive functions, the circulation and secretion of the liver, and the actions of the intestinal canal, in a secondary manner; thereby, in a gouty diathesis, predisposing to the disease.

Severe Study.—This cause, comprising not only want of exercise, lateness and irregularity in the hours of rest and sleep, but also its consequences, weakness of stomach, and inactive bowels; by its effects on the nervous system, from the over-action of the brain, produces that form of irritative debility, which increases the susceptibility of the frame to disease, and consequently to gout, if such be the predisposition of the individual*. Sydenham mentions that his immoderate application to the composition of his Treatise, occasioned him the severest fit of the gout which he ever had†.

One gouty gentleman, of plethoric habit, informs me that if he allows himself to read with any intensity of attention, or for a long continuance, he experiences the same kind of giddiness which he suffers from stooping.

Animal Food.—This, when taken in excess of quantity, not only weakens the stomach by distention, but stimulates and oppresses the whole digestive process, beyond its healthy power. In the form of high-seasoned

* The late Mr. Pitt, and his father, had gout at a very early period of life. The father was never a votary of Bacchus, and neither of *Venus* (as we are told); but both were *ardent students*.

† Dedicatory Epistle to Dr. Short.

dishes, it proves a cause of morbid excitement, and induces such a false degree of appetite, that a quantity sufficient for several meals is often comprehended in one alone. An improper amount of quantity in the twenty-four hours, however divided, serves very materially to lay the foundation of a plethoric state; but in order that the gout may be excited, the influence of fermented liquors must be superadded.

Strong Liquors.—Of these, in relation to the gout, wine, if in excess, is certainly the most injurious. The use of raw spirits destroys the appetite, weakens the tone of the stomach in a permanent manner, and even leads to disease of structure; so that although an inflammatory condition of vessels is produced, the *plethora ad molem*, as it has been called, is counteracted. Wine contains so much more of alcohol in a given bulk, than malt liquors possess, that it is probably from this cause, and at the same time from its not breaking down the energy of the stomach as spirits do, that it proves so active an agent in introducing the gout. Much of this effect also depends on the kind of wine which is taken, as the gouty know by constant experience. Champaign, bad claret, and new port, will predispose to the gout much more strongly, than equal, or even larger quantities of madeira and sherry; because, in addition to their equal or greater heating effects, they give rise to more acidity; from which, the stomach and alimentary canal suffer certain irritation. In a system, otherwise fitted for gout, or in the settled gouty habit, the limbs quickly suffer by sympathy. The light wines and aceseent liquors made from our own fruits, contribute to the gravel rather than the gout; and raw spirits more particularly induce severe dyspepsia, obstructed and hardened liver, and dropsy.

In comparing the influence of different liquors, it must be considered, that wine-drinkers also partake much more largely of the solid luxuries of the table, than drinkers of spirits and malt liquors can do, and therefore produce more powerfully the combined effects of heating excitement and redundant assimilation. Dr. Rush, contrasting the diseases and remedies amongst the Indians of North America, with those of civilized nations, makes the following observations: "I have heard of two or three cases of gout among the Indians, but it was only among those who learned the use of rum from the white people. A question naturally occurs here, and that is, why does not the gout appear more frequently among that class of people, who consume the greatest quantity of rum among ourselves? To this I answer, that the effects of this liquor upon those enfeebled people, are too sudden and violent, to admit of their being thrown upon the extremities, as we know them to be among the Indians. They appear only in visceral obstructions, and a complicated train of chronic diseases. Thus putrid miasmata are sometimes too strong to bring on a fever, but produce instant debility or death. The gout is seldom heard of in Russia, Denmark, or Poland. Is this occasioned by the vigour of constitution peculiar to the inhabitants of those northern countries? or is it caused by their excessive use of spirituous liquors, which produce the same chronic complaints among them, which we said were common among the lower class of people in this country? The similarity of their diseases makes the last of these suppositions the most probable. The effects of wine, like tyranny in a well-formed government, are felt first in the extremities; while spirits, like a bold invader, seize at once upon the vitals of the constitution."

I saw lately a corpulent and robust man, aged thirty-

four, labouring under very severe gout, for the fourth time. Both feet were affected. It was entirely acquired, and began at the age of twenty-nine. In the earlier part of his life, he had been constantly on the sea coast as a smuggler, and at the same time that he was exposed to the hardships of that irregular life, he drank two or three pints of hollands a day, during the long period of four or five years; with no other effect than being rendered very nervous, and occasionally dyspeptic. Afterwards, he became a bricklayer; and being possessed of property, indulged himself both in wine and porter, in addition to a slight use also of spirits. His exercise was much less active than formerly, and after about two years of this new mode of living, the gout came as the result.

In another instance also of a very similar kind, the patient, until he was twenty-eight years of age, led a country life, in a situation where he drank hollands freely and constantly. He then took the place of a butler in London, and entered upon the usual inactive habits of that situation, in concurrence with indulgence in wine and strong malt liquor, and abundant eating. His present age is fifty-nine; and during the last twenty-four years, he has not, in any one year, escaped a severe attack of gout. In this case, as in the former, the gout was unknown in the family; and each patient was of the make and temperament, which I describe, as the most predisposing to the disease.

Van Swieten states*, that the Dutch knew little of this disease till they changed their favourite beverage, beer, for wine. Linnæus informs us, that the Laplanders and many country people in Sweden, who drink malt spirits, but never wine, are strangers to the gout†.

* Commentaries, 1255.

† Mr. Good, in his profound and interesting work on Nosology,

I am disposed to think, that in this country, and particularly in the metropolis, gout is much increased in frequency among the lower stations of life, since the very general and free use of porter. This is a very nutritious fluid, and in conjunction with spirits, even with a moderate quantity of solid food, may be viewed as inducing the plethoric inflammatory state, and as a consequent introduction to gout.

In Scotland, gout is much more rare than in England. In Edinburgh, where the habits of the people approach the nearest to those of London, it is found most; but it is scarcely ever known among the inferior classes. In two thousand two hundred cases of disease admitted into the Royal Infirmary, as clinical patients under the care of Dr. Gregory, there were only *two* examples of gout. I also learn that Dr. Hamilton, who has been one of the physicians of the Infirmary nearly thirty years, and must in that course of time have admitted many thousand patients, has not seen more than two cases of gout, out of the total number of diseases under his care in that Institution. In the London hospitals, on the contrary, according to my recent inquiry, gout is by no means so uncommon; although, as it is not usual to admit cases of gout, its frequency in the list of hospital patients, must not be expected*.

observes, "Gout is one of the maladies which seems to have been common to England in its earliest ages of barbarism. It is frequently noticed by the Anglo-Saxon historians, and the name assigned to it is *foet-adl* (foot-ail).

* In the first edition of this Treatise, I committed a remarkable mistake with respect to gout, in my reference to the valuable paper in the Medico-Chirurgical Transactions, "On the Comparative Prevalence, Mortality, and Treatment of dif-

In Glasgow the gout is very rare, even amongst the higher classes; and, in connexion with this fact, I may mention the following circumstances. Among a population of about 110,000 persons, which Glasgow and its environs comprehend, there are not, as far as I could learn, twenty private carriages in use. The demand for hackney coaches is so small, that the proprietor has found it necessary to discontinue the regular stand, and the whole number of them does not exceed four or five. Sedan chairs are in common use with the ladies, but the gentlemen prefer the wholesome exercise of walking; and, as this city does not rank inferior to any other in liberal hospitality, the infrequency of gout must, in great measure, be imputed to the greater comparative activity

ferent Diseases; by Sir Gilbert Blane, Bart." The cases of gout, which I incorrectly stated as being in the list of the hospital patients, occurred only in the private practice of this physician; as the following quotation shews. "The diseases which stand most prominent for their prevalence among the upper classes of society, are gout, disorders of the stomach, and liver complaints. With regard to gout, there is not a single case of it to be found in the hospital list, whereas there are in the private list one hundred and thirty, constituting about a twenty-sixth part of the whole. No disease affords so strong a proof of the power of habits of life over health."

This calculation of Sir Gilbert's hospital practice, refers to the years from 1784 till 1794, and comprehends the number of 2406 patients. For the reason which I have suggested, it must not be concluded, that gout does not occur amongst that class of persons, who are admitted into the London hospitals with other diseases. At the same time, I grant entirely, that the prevalence of gout in the respective classes of society, proceeds in an exact ratio with the freedom used in animal food, strong liquors, and general indulgencies. The remainder of my argument I have already offered.

of the people, and better regulation of the general habits, than belong to London. In Glasgow, also, punch is a more general beverage at the best tables than wine. Hence I often heard it facetiously remarked, "that punch keeps off the gout." It may certainly be said, that weak acidulated punch is much more diuretic than wine, and less heating in its effects. But although the argument may be maintained, that punch does not induce a predisposition to the gout, no one would deny that a free indulgence in this liquor must impair the healthy powers of the stomach, and lead to some serious disease. The lower orders in Scotland, who drink whiskey freely, and live chiefly on farinaceous and vegetable food, acquire severe dyspepsia, liver disease, and dropsy, or diabetes*; but seldom, or never, the gout.

Indolence powerfully assists the predisposition to gout. It promotes the tendency of full diet to produce plethora, by its restraining the various excreting functions; and also prevents that regular distribution, and due expenditure of the nervous energy, which the healthy action of the brain, and the several secreting organs, demand. It may be said to cause a morbid accumulation of nervous excitability.

Of facts, directly to illustrate the influence of the predisposing cause now under consideration, I might offer an ample list; but the following examples will be sufficiently in proof.

* I am led to believe, from attentive observation, that *Diabetes* is considerably more frequent in Scotland than England; and that the opposite habits of the people in these two countries, such as I have described, tend, in a sort of respective ratio, to produce gout with the one course of life, and diabetes or dropsy with the other.

A gentleman, whose aunt only in his family had gout, of strong constitutional predisposition to the complaint, was first attacked at the age of thirty-five, when he was filling the situation of school-master in the country. His usual habit was to take animal food freely both at dinner and supper, with malt liquor, and either a small portion of wine, or a glass of mixed liquor. He described himself as carefully moderate in the use of liquor. The gout returned severely at short intervals. Soon after this he had occasion to travel into Ireland, during a period of nine months, and drank both wine and whiskey very freely; but his exercise was exceedingly active, and he was free from gout. Resuming again his sedentary life, although with more care than formerly in his habits, the gout soon came back with violence; and in the course of its progress, has affected the feet, knees, ham-strings, hands, and elbows. A similar example has lately come before me. A school-master, extremely zealous in the discharge of his duties, of large frame, circular chest, very plethoric, having invariably a good appetite, and indulging to the extent of eating heartily of animal food at dinner, and supper; temperate in vinous and spirituous liquors, but every day partaking of one or the other in some degree, and rather freely of grateful malt liquor; so sedentary as to pass days and weeks without exercise, even when well; became gouty at the age of twenty-four, the disease being unknown in his family. I have seldom seen a greater martyr. His fits, and the distressing consequences, have sometimes been of six months' continuance.

An officer, whose grandfather only in the family had gout, had been subject to severe attacks twice every year. He was employed on very active service in Ægypt, for nine months, and often lay out on the sands

protected only by carpets, the nights being cool and the days hot. He drank both wine and spirituous liquors very freely, and much more so than before or since; but his bodily exertions were excessive, and his mind was enthusiastic in his duties. He was quite free from gout. Leaving this station, he went to Gibraltar, and there becoming very inactive, in the course of three months, the disease returned with all its violence; and has ever since affected him severely, twice a year, as before.

Plethora.—A redundancy of blood in the general system, is certain to be a result of the combined influence of the three preceding remote causes; and appears, as far as I have observed, to be the condition most preparatory to the operation of the exciting causes. That form of plethora which is connected with congestion in the circulation of the liver, may certainly be considered as peculiarly concerned in the production of gout.

Nimia Venus.—From this cause, the nervous energy becomes unduly exhausted; much irritative debility is produced; and the digestive functions, together with the general powers of the system, are disordered. Habits of indiscretion, in this respect, are so constantly joined with the gratifications of the table*, that indirectly it may be considered a predisposing cause of gout. Syden-

* The gout was saluted by the Greek poets as the daughter of Bacchus and Venus:

“Αυσιμελου Βάκχου, και λυσιμελους Αφροδίτης,
“Γένναται θυγάτηρ, λυσιμελής, Ποδάγρα.”

And in the same sense runs the Latin verse:

“Ut Venus enervat vires, sic copia vini,
“Et tentat gressus, debilitatque pedes.”

ham describes this cause, as giving "the seeds of the gout;" and this position has some little truth; but I conceive that the agency which brings the disease forth, consists in the associated habits of living. Voluptuous passions, and the excesses of the table go hand in hand; and many of the gouty are too correctly characterised in the Greek and Latin epigrams just quoted.

Climate, Season of the Year, and State of the Atmosphere.—Although no climate can exempt the gouty constitution from the disease, if too indulgent habits of living are pursued; yet, a hot climate, by the copious and regular perspiration which it induces, furnishes, I believe, but comparatively few examples of the first occurrence of gout. I have met with a few rare instances in which I learn that the disease has first taken place, wholly acquired, without hereditary influence, both in the East and the West Indies; and, more frequently, I have been consulted by patients who inform me that they had several of their attacks while residing in one or other of these climates. All such individuals have been inconsiderate in the use of wine and rich living. Others, and certainly the more careful, have profited very much by the influence of a hot climate.

Long continued heat of weather is favourable to a few constitutions, and particularly to those in which rheumatism occurs in conjunction with gout; but if I judge from the summer of 1818, it does with the majority rather predispose to gout; and much more especially it has had this unfavourable effect, with those persons who have been confined by their pursuits in the metropolis, and scarcely enjoyed the beneficial change of country air. It has appeared to me that the body has acquired a feverish condition, together with some debility, from the excitement

of a constantly hot atmosphere, to which, it may be said, our English dress, and houses, and customs, are not suited. Exercise is taken with less inclination and less advantage than in more temperate weather.

In the intense heat of June 1817, I witnessed both the predisposing and exciting operation of this state of atmosphere on gouty persons.

Variable climates certainly operate the most in predisposing the body to gout; and locality of situation has a powerful influence; so that the gouty patient sometimes finds himself compelled wholly to change his place of residence, for one which is dry and protected.

Changeable weather, and especially in spring and autumn, when cold winds prevail, with frequent wet, acts very strongly on the gouty diathesis; and causes almost habitual achings, and increased stiffness in the joints which have been affected; also predisposing the system to a paroxysm. The regular functions of the skin, so important to health, suffer material interruption; and hence, the chief explanation of the ill consequences which ensue.

Some naval gouty patients inform me that they have not been more free from gout, when at sea, than on shore. They state that they have been much more inconvenienced in regard to a confined action of the bowels; which circumstance ranks so pre-eminently in the list of predisposing causes.

A MORBID STATE OF THE DIGESTIVE ORGANS, CONSIDERED BOTH IN ITS PARTICULAR RELATION TO THE GOUT, AND IN A GENERAL VIEW.

WHEN we consider that gout is truly a disease of repletion, it follows as a necessary consequence, that the

over-excitement of the stomach and chylopoetic functions must lead to relative derangement and debility. Sydenham observes, "that upon a thorough attention to the various symptoms of the disease, he judges it to proceed from a weakened concoction both of the solids and fluids." This is synonymous with the more modern language of unhealthy assimilation, which in quantity, and probably in quality, may be viewed as the parent source of the disease. It is in this, that the acquired predisposition, or that *not hereditary*, consists; and in most instances, the hereditary predisposition is aggravated by the same cause, before any developement of gout takes place. In a first fit of gout, and sometimes also in subsequent attacks, there is no very sensible disturbance of the digestive organs; but it does not follow that they are really in a state of healthy action. The patient judges only from partial indications. An active appetite is often joined with an unhealthy state of stomach, and particularly with an unequal performance of functions below this organ, after the food has undergone its first change. In the case of the *epicure*, there is much of artificial appetite produced by the habitual and excessive gratification of the palate. The patient is often deceived into a belief, that being possessed of appetite, and apparent powers of stomach for the enjoyment and benefit of a hearty meal, his digestive organs cannot be in fault. But we must keep in view, that in the stomach itself, the important process of digestion only begins. Its next stage, and a very material one it is, takes place in the duodenum, and comprises the proper secreting action of the liver, and the healthy influence of the bile. The due excrementitious function of the bowels, constitutes also an essential part of the process towards healthy chylication; and it may be fairly concluded, that upon the correct per-

formance of these preparatory steps, the subsequent healthy and complete assimilation of the chyle into blood, will greatly depend. A morbid interruption to any part of the train of digestive functions, may become a predisposing or an exciting cause of gout, in the same manner as it is a cause, which commonly exerts an influence in all other diseases.

The particular symptoms which arise, will, on accurate examination, generally indicate, with considerable fidelity, the particular seat of morbid action. On this head, I shall endeavour to offer some remarks, taking a general view of the subject.

When the stomach is the most affected part, the marks of indigestion are felt in the most sensible manner, by some or all of the following symptoms:—Heartburn; eructations which are sour, attended with a sense of heat, and often conveying the odour of yesterday's meal; a craving appetite, which does not become comfortably satisfied; oppression after a meal, with a painful sense of distension, and soreness of the whole epigastric region. This sense of soreness is sometimes experienced in so extreme a degree, that only slight pressure from dress can be allowed; and even a gentle touch at the ensiform cartilage, or just below that part, is dreaded by the patient as an act of violence. To this account may be added, a furred tongue and clammy state of mouth; with viscid saliva, which is experienced especially in the first of the morning. Its taste is often remarkably saltish. In dyspeptic persons, in whom the nervous temperament predominates, the tongue is coated either with a white, or yellowish-white fur; but in those of the sanguineous temperament, the colour of the fur is a deep brown; or brown mixed with white. In the former class of patients, the complexion of the tongue is very commonly

pale; in the latter, quite red. But of all the appearances of the tongue, which indicate a debilitated state of the stomach in its greatest degree, is the cherry-red colour of the whole surface, with more or less of cracks in its substance, and a prominence of the papillæ. By these last characters, the pale tongue also, after a long duration of dyspepsia, is sometimes distinguished, and is to be considered as giving a very unfavourable evidence of the weak powers of the stomach. An extremely clean tongue, with streaks of white and red, is another distinct appearance, as the index of debility. Accordingly, as nervous irritation prevails in the general state of the constitution, a frothy appearance of the tongue and fauces, may, in addition to what I have above described, be noticed. Nausea, occasional sickness, flushings after eating a stimulating meal, a giddiness on sudden change of posture, and an uneasy or painful state of head, also occur. With this dyspeptic condition of stomach, the bowels are irregular, but for the most part torpid. Many patients relate that the only predisposing circumstance as introductory to the fit, to which they can make reference, has been a costive state of bowels. The secretion of the urine is variable, both in its quantity and in its properties. Sometimes it is deficient in quantity, of a deep colour, and of high specific gravity; at others, abundant, pale, and much diluted; in which case it is passed with much nervousness and irritation, but without difficulty.

When the chief seat of complaint is below the stomach, the tongue often bears an almost healthy appearance, and the appetite is strong, and for the most part regular. The symptoms are occasionally such, as are in common language called bilious; with which, a general sallowness, or partial stains of yellow in the skin, and a dark

colour around the lower eye-lid, are more or less associated. While indigestion thus takes place in the intestinal canal, the stomach itself appears to be in too strong action, and prepares more food than can be afterwards converted into good chyle, and assimilated into healthy nourishment. I have met with instances among persons subject to gout, in which the functions of the stomach alone have been much improved by medical treatment, so that the digestion which was formerly weak, has become sensibly stronger; yet, from the subsequent process not being healthy, the good appetite has led to a false nourishment of the body, and frequent fits of gout have ensued. All the secretions become vitiated. The bowels suffer much irritation, being often affected with painful excitement, fruitless of relief, and occasionally amounting to tenesmus. The peristaltic action is irregular; the *fæces* have an unnatural consistence and colour, and are peculiarly offensive. Sometimes they are passed as *pellets*; and when more formed by the bowel, it occurs, not unfrequently, that they are so contracted in size as to convey the apprehension of some stricture of the canal; but the effect appears to be really owing to the unhealthy state of secretions, and to muscular contractions of the bowel irregularly occurring from irritation. It sometimes happens that air is formed so abundantly in the intestines, either from undigested aliment, or possibly from actual secretion, or both, that its effects in producing painful distention, alone serve to cause a most distressing state of complaint. The discharges are commonly not formed, but are remarkably tenacious. They assume the appearance of pitchy blackness, or are muddy, or resemble dirty clay; and when the complaint has been of long continuance, an excessive quantity of mucus is secreted in the intestines, which incorporates itself with the *fæces*,

and sometimes exhibits almost the appearance of purulent discharge*. This mucus is to be distinguished from the gelatin-like appearance, which is seen occasionally in the evacuations, when the bowel suffers excessive irritation from an acrid purgative, or when under dysenteric affection; in which case, it appears detached from the fæculent matter, apparently almost organised from its firm consistence, and is indeed altogether different from the slimy accumulation before mentioned. This habitual mucous secretion has always appeared to me an indication, that the morbid condition of the alimentary canal has been of long standing. In the same manner we see the urine loaded with mucus, when the bladder is diseased, or under permanent irritation from a morbid condition of its contents. In investigating a disordered state of the digestive functions, frequent observation of the excretions is an essential source of information to the practitioner. Mr. Abernethy, in addition to his original and highly instructive views of the present subject, has laboured very usefully in exciting among the profession an increased attention to these points. He observes, that “the colour of the alvine excretions in a disordered state of the viscera is various. Sometimes they appear to consist of the residue of the food, untinged by bile. Sometimes they are of a light yellow colour, which denotes a very deficient quantity of healthy biliary secretion. Any kind of brown which dilution will not convert into yellow, I should consider as unhealthy, since the colour of healthy bile is a bright yellow, which by concentration appears brown†.”

* If water be poured on these fæces, the mucus separates into small flakes.

† On the Constitutional Origin and Treatment of Local Diseases, p. 25.

It may be stated, that a bright yellow appearance indicates a fresh and hurried secretion of bile, and shews irritation rather than regular healthy action. A clay-like appearance is a sure indication of an obstruction of the bile, so that it does not enter the duodenum; and where the opposite colour is seen, sometimes to the degree of pitchy blackness, we may conclude, that not only the bile in particular is in a most unhealthy state, but also, that the secretions of the alimentary canal itself are much disordered. Another evidence of a wrong action of the liver, is very frequently found in the greenish appearance of the discharges from the bowels; proving unequivocally a highly vitiated state of the bile. With this, there is often further attendant proof of derangement in the general secretions of the canal. It is objected by some, that conclusions drawn from the observation of the excretions are fallacious, because many of the appearances which have been mentioned, will be produced occasionally in healthy persons, from irregularities in diet. In answer to this, it may be affirmed, that such persons will have their feelings of comfortable health and tranquillity disturbed, in however transient a manner, when the alimentary secretions thus become vitiated. The occasional hypochondriasis which is experienced by many who enjoy general good health, will, I believe, on examination, most commonly be found connected with the faulty state of the digestive organs, and corresponding indications in the appearance of the excretions.

I must repeat, that the care to be informed, by actual and repeated observation, of the state of the several secretions, is most assuredly, in many cases, quite indispensable towards forming a correct medical judgment.

More or less of jaundice usually accompanies the most urgent of the disordered functions which I have just de-

scribed; and in proportion to the degree of this evidence, the nature of the complaint is apparent to common observation.

I have known in several instances, that a dyspeptic and bilious state of complaint has occurred as a preliminary, many weeks before a first attack of gout; and in such instances, the patient has found himself gradually becoming larger in the abdomen; oppressed at the stomach; and affected with many nervous sensations. Very lately, I saw a gentleman labouring under a first fit of gout, who, for two or three months before, had suffered from an obstructed state of liver, with many attendant symptoms of derangement in the stomach and alimentary canal. So strongly was his nervous system affected, that, together with flutterings near the heart, and occasional palpitation, he was frequently seized with almost a loss of sight when walking in the street. Sometimes objects appeared double; and much dimness of sight occurred on the slightest occasion. The whole of this sympathetic defect of vision yielded to a course of medicine; of which, mercurial purgatives formed a material part. Returning to some freedom of living, before the complete correction of the internal functions was produced, the gout arose.

In another case, attended with very similar symptoms, almost every attempt to stoop, as in the act of adjusting the shoes, was instantly followed by that sudden spasm of the diaphragm, which the patient describes as the cramp of the stomach. This symptom occurs chiefly with very corpulent persons; in those especially, in whom the nervous temperament predominates; but it also occasionally affects others, whose temperament is sanguineous. Thus it sometimes happens to the gouty patient, that when he is most complimented on his florid looks, and good appearance, the disease is making its insidious approach.

In such instances, I have invariably found that unhealthy corpulence about the abdomen has been gradually forming, and, in other words therefore, a state of congestion in the vena portarum system.

In those who have had frequent attacks of gout, the occurrence of urgent premonitory symptoms may be said to take place, chiefly in proportion to the degree of the abdominal visceral derangement previously existing.

From the connexion by structure which exists between the diaphragm and abdominal viscera, and the liver especially, the respiration is very commonly imperfect, and, in some instances, even difficult and painful, on every occasion of increased exertion, when a morbid condition of these parts is existing in any considerable degree. This disturbance may be produced in a transient manner, from the influence of disordered functions simply; or, may be the more permanent result of altered structure of contiguous parts.

I have seen this sympathetic affection of the diaphragm, in one case, amount to so great a degree of spasmodic asthma; attacking the patient periodically, at first twice, and afterwards once in the twenty-four hours; that each paroxysm was dreaded with horror, from the fear of suffocation. It was of three or four hours' continuance; and until active treatment was adopted, the attack followed with certainty, and almost instantly, the least incaution either in the quantity or quality of the dinner meal.

One gentleman relates, that for a considerable time previously to his last severe fit of gout, he not only complained of oppressive abdominal corpulency, which had been slowly coming on, but also of the sensation of a strong cord across the epigastric region, just below the diaphragm; causing very great stricture of his breath in stooping.

In the dyspepsia of gouty persons, both in consequence of their temperament, and from the established influence of the gout on the nervous system, the severest state of hypochondriasis sometimes prevails. Under such circumstances, the connexion of the functions of the brain with those of the digestive organs, is particularly exemplified; and a striking picture is seen of the reflected and mutual influence, which prevails between the mind and the body in disease. When that state of the system is taking place which is introductory to the return of the gout, the bowels fail in their regular action; the secretion of the urine is variable in quantity and in appearance; the patient becomes sensible of enlargement in the abdomen, while the limbs sometimes are shrunk, and the muscles are soft; he complains of various nervous feelings; and is occasionally troubled with night-mare, and with cramps.

Dr. Cullen, in describing the atonic state of gout, remarks of the affections of the alimentary canal, "that they are often attended with all the symptoms of hypochondriasis; as dejection of mind; a constant and anxious attention to the slightest feelings, an imaginary aggravation of these, and an apprehension of danger from them*."

The active sympathy which exists between the stomach and the limbs, in certain states of chronic gout, I shall hereafter have to consider.

Under a disordered state of the digestive functions

* I have had occasion to observe that the dyspeptic invalid finds the morning his most healthy and comfortable time; but that, on the contrary, the truly nervous patient is wretched through the day, and re-animated only in the evening, by the influence of dinner, stimulants, and cheerful society.

(whether the patient is gouty or not), and in proportion to the duration of complaint, and its dependance on change of visceral structure, the symptoms which arise from sympathy of distant parts, or contiguous organs, are strongly and variously marked. In addition to palpitations, an occasional disposition to fainting, and other symptoms already described, I may mention as another effect of the morbid sensibility of nerve which is prevailing, a remarkable tendency in the nerves of the arms and legs to be affected with numbness both by day and night, on the application of very slight pressure from accidental posture; and of disagreeable tinglings, which usually most affect the fingers.

The bowels are frequently distended with undigested food, and with the air which is generated in consequence; so that many uneasy sensations are felt through the whole tract of the canal, which sometimes assume the character of spasm.

Flatus in the stomach and bowels, or in the bowels alone, is now and then the sole symptom of indigestion, of which the patient confesses himself sensible; and this is, perhaps, detected only on attentive inquiry: but it may be received as a sufficient proof of an unhealthy condition of the digestive powers. In this case the stomach and bowels are very deficient in muscular tone, and a full meal with certainty occasions a distressing feeling of weight, and general distension, which latter symptom is sometimes very painfully oppressive.

The hæmorrhoidal veins are occasionally painful; and a discharge of blood, which in some persons is quite irregular, and in others is almost periodical, very commonly accompanies the gouty diathesis. The discharge in question sometimes occurs very profusely, and is of a dark venous colour, or of a blackish hue. When the hæmor-

rhage is slight, it is more usually arterial, and of a florid red.

The local abdominal sensations vary according to the parts most affected. It is difficult, however, to fix on the precise seat of diseased action, concerning which, we are indeed often liable to be deceived. With parts of associated functions, there will be many sympathies spreading from contiguity. Mr. Abernethy observes, "It is probable that no material disorder can ordinarily take place in one of the digestive organs, without disturbing the functions of the others. When digestion is imperfectly performed, the functions of the intestinal canal will soon participate in the disorders of the stomach. Under these circumstances, the secretion of bile will also become irregular. Should disease commence in the large intestines, as about the rectum, it disturbs the functions of the stomach and secretions of the liver, and becomes augmented in its turn by its sympathy with these parts. Should the liver be disordered in the first instance, the stomach and bowels may not immediately sympathise, although they will probably soon become affected*."

Sometimes the patient describes his feelings of uneasiness to be exactly in the line of the duodenum, passing through also to the back in the opposite direction. The sensation is that of aching dull pain, with occasional shooting, and of heat, which often amounts to burning. In other cases, the right or the left hypochondrium is the seat of heavy dull pain, which is increased on pressure†.

* On the Constitutional Origin, &c. p. 44.

† In examining the abdominal regions by pressure, in suspected disease of the contained viscera, we should not from our *tactus eruditus* hastily draw conclusions of the soundness, or

This is very dependant in its degree on the state of the bowels, and is mixed also with pain in the back and shoulder blade. It frequently happens that the only part which is tender or painful on pressure, is just at the ensiform cartilage. In association with a morbid state of the alimentary functions, this tenderness, especially if it be found when the stomach is not distended by food, is certainly indicative of a wrong condition of the liver. An irritable state of the abdominal muscles is a common attendant on a disease of the liver of long standing, and it is found very remarkably in some constitutions. In these cases, the muscles contract suddenly, even on gentle examination; and the abdomen, in consequence, conveys the feeling of hardness and rigidity. There are examples, in which, with uneasiness and even occasional strong pain in the parts which I have described, strong pressure is usually borne without complaint. In these cases, we may infer, that the viscera are most affected in their internal structure, or at their posterior surface.

An inability to lie on one or the other side is a frequent diagnostic of a tender state of the liver, and of the particular part which is affected. It usually happens that the patient chooses his posture on the diseased side; for, in this way, the muscles of the opposite ribs alone are called into action, for the purpose of assisting respiration.

morbid alteration of any of the organs. The mode in which the examination is made, is of importance. When the patient is in bed, the legs should be drawn up, so that the abdominal muscles may be relaxed. If not in bed, and a sofa be not at hand, the patient's most favourable posture for examination, is that of resting the hand on a table, and leaning with the body bent; or, more conveniently, leaning forward at ease in a chair.

The sympathetic pains are sometimes more remote and anomalous, affecting the shoulder blades, shoulders, or even the arms, as with rheumatism ; for which, they are often mistaken.

The pulsation in the epigastric region, which is sometimes felt in an alarming degree by dyspeptic persons*, is seldom met with in persons who are subject to acute gout ; and appertains rather to those, who, together with an unhealthy state of stomach and liver, have an obstructed and vitiated state of mesenteric glands ; are badly nourished ; and possess a morbid sensibility of nerve, in a high degree. It is occasionally found in those who have been weakened by frequent attacks of gout ; have lost their flesh ; and now have the disease only in its chronic form.

A lady, forty-eight years of age, who suffers from enlargement of the liver, and always more or less from a faulty action of this important organ, strongly possessing the nervous temperament, is affected with the pulsation in question, so as almost to imitate the aneurisinal beating of the aorta. She experiences this troublesome symptom, most, when her stomach is disordered ; more when in bed than up ; and it is invariably aggravated when lying on the left side. Her pulse is always regular.

I must add to this description, that in many instances, and probably in most, the pulsation in the epigastric region is found in patients, in whom, no visceral enlargement can be discovered. I have met with five cases of this description. In two of these, the pulse at the wrist was

* Dr. Baillie has offered many valuable observations on this subject, in a paper, intitled, " Upon a strong Pulsation of the Aorta in the Epigastric Region."—*Med. Trans. of the College*, vol. iv.

on some days intermittent, on others, very regular. I have been quite convinced of the justness of Dr. Baillie's observation, that this very troublesome symptom is not to be considered really alarming. It seems to be, and as Dr. Baillie appears to view it, a sympathetic action of the artery dependant on a disordered state of the digestive organs, in an irritable constitution. I conceive it analogous in its nature to a palpitation of the heart, the latter symptom usually being more associated with general perturbation of the nervous system, and more under the influence of the mind, than the pulsation in the epigastric region. But I must add, that this symptom is also very commonly dependant on some error in the structure or functions of the abdominal viscera.

One gentleman, on the accession of a severe paroxysm of gout, was first affected with a strong palpitation of the head, and afterwards in the back, in the situation of the right kidney. The symptom was most troublesome at night.

Another gentleman, under the united influence of gout, and of bilious obstruction, suffered from a palpitation of the heart, and the pulsation of the aorta in the epigastric region, alternately.

Again, in a case of gout attended with a marked degree of bilious obstruction, and unhealthy state of the bowels, the symptoms complained of, were, heart-burn and violent eructations, palpitation of the heart and flushings in the chest, together with now and then a curious feeling of explosion in the head, preceded by a sense of tightness of its integuments; and a starting of the limbs at night, which latter symptom I have before described as a frequent precursor of an attack. It also happens as a consequence of a long paroxysm having enfeebled the nervous system.

To conclude with cases of this description, I shall make reference to the symptoms of a gouty patient aged forty, who, when labouring under the chronic symptoms of the disease, experienced flatus in the stomach and bowels, torpid action of the bowels, occasional palpitation of the heart, and pulsation in the epigastric region, general languor, great depression of spirits, broken rest by night, and in the day, a remarkable sense of weariness in the arms.

In the dyspepsia which attends an obstructed state of liver, and the various symptoms of indigestion of the lower bowels, the pulse at the wrist seldom conveys any corresponding indication of such derangement; but I must add, however, that the circulation is by no means equally performed in such cases, under particular excitement of diet; and in proportion as the temperament is sanguineous, marks of morbid determination of blood to the head, attended with pain, throbbing, heat, weight, and vertigo, are readily produced. At night, together with much restlessness, the palms of the hands are affected with dryness and burning heat, while the feet are remarkably cold. When the powers of the stomach, in a person of nervous temperament, have been weakened from errors in living, or other unfavourable causes, the circulation is universally languid, the extremities are easily chilled*, and the general appearance is unhealthy and delicate.

The threatenings of apoplexy which afflict some gouty persons, and others, I can only advert to in this place, as

* It happens with a few to suffer habitual coldness of feet to such a degree, that, even in summer, it is found necessary to wear very warm stockings.

connected occasionally with permanent obstruction and congestion of the liver. The aortal circulation not being carried on with due freedom in the abdomen, an unhealthy distribution of blood to the brain becomes the consequence. Dulness of the faculties, occasional head-ache and vertigo, flushings, and the feeling of morbid heat about the head and neck, a sensation of the head as if covered by a tight hat, and of the neck as if too much girt by the neckcloth, are the most ordinary symptoms.

Sometimes the symptoms are more remote, and afflict one or both arms. The patient complains of a sense of bigness in the limb, with numbness, and upon holding it up, it seems heavy, and soon aches in a distressing degree.

In the case of a lady fifty-four years of age, of the nervous temperament, for a long time subject to obstruction of the liver, and occasionally of the common duct, having a dark and bilious complexion, the bowels themselves in an unhealthy state, and the fæces remarkably loaded with mucus, I have found in two or three of her attacks a very slow pulse, the most extraordinary sensation of sinking at the pit of the stomach, as if there were an entire vacuity in the epigastric region, and such general faintness, as to alarm the mind with the apprehension of immediate death; a coldness of limbs; but no pain in any part of the body. I shall hereafter discuss the treatment of the case, and only now mention that the complaint yielded to a perseverance in regimen, and the use of purgative, corrective, and tonic medicines, in methodical succession.

How much dyspeptic persons are sensible to the influence of variable atmosphere, is familiarly known; and with gouty persons in particular, any considerable check which is given to the ordinary action of the skin, is with certainty succeeded by some uneasy internal feelings,

together with sympathetic pains in those limbs which have been affected with gout. A harsh and dry state of skin is a common attendant on bilious dyspepsia; and a distressing itching, particularly affecting the back and arms, is very much complained of by those who are liable to gout. It not unusually occurs as one of the premonitory symptoms. An erythematous rash; and urticaria, or nettle rash, also occur; and in one remarkable instance, I saw the latter irritation exist in a violent degree, during two days before the paroxysm; to which, and the treatment adopted, it immediately yielded. Other forms of cutaneous complaint also occur to the gouty patient, in common with all those who have an unhealthy condition of the digestive organs.

Itching of the skin, with or without eruption, is a very common consequence of an indiscretion being committed in the taking of certain acids, as pickles and unripe fruit; or, if this do not happen, the stomach and bowels are the more irritated, and diarrhœa follows.

Having now taken a cursory view of several points relating to the derangements of the stomach, the liver, the intestinal canal, and the skin; and noticed the sympathetic states of the circulation and of the nervous system; it remains that I should deliver some account of the function of the kidneys, as constituting a very important link in the pathology of the digestive organs.

With respect to the gout, an attentive observation of the changes and condition of the urine is an abundant source of information. It is very commonly found upon inquiry, that the patient has noticed a very deficient secretion of urine a short time before the occurrence of the paroxysm, and consequently it has been of a deeper colour than is natural. It also now and then certainly occurs, that for a day or two before the fit, the urine is passed copiously, and of a pale colour; but this happens

only in persons quite of the nervous temperament, and whose constitutions have been much weakened by gout.

The quantity of urine which is secreted in the paroxysm is usually scanty; and even when it is in natural proportion, is much concentrated. A portion of that which is passed the first in the morning, should be the particular example for medical observation and examination. The colour of the morbid urine is various, and is affected by different causes; but whether more or less deep, it is found under these circumstances much increased in specific gravity, beyond the standard of health. This, from my experiments, which have been very numerous, I should state as ranging from 1·010 to 1·015*. In the morbid urine to

* With a view to form some opinion of the degree in which the specific gravity of the urine, in health, might be influenced by modes of diet, I examined the morning urine of two healthy persons daily, for ten days; during which period, their habits, as to fulness of living and moderation, and exercise and rest, varied exceedingly; and my results were, that under the greatest extremes of their regimen, the specific gravity of their urine was remarkably uniform; differing certainly, but not as might probably be expected. In several trials, I found that the urine of health possessed its highest specific gravity about two hours after dinner. It does sometimes happen that the early evening urine deposits the brick-dust or pink sediment, when that which is passed the first in the morning is almost or wholly free from it. In such examples, the evening urine has a higher specific gravity than any that is secreted in the twenty-four hours. Yet, notwithstanding, the general rule which I have just now mentioned, of selecting the first morning's urine for particular examination, will be most satisfactory. In the peculiar cases to which I have alluded, I have considered that the assimilation of the food has taken place very quickly. The patients have been of the indulgent class in their diet, and of irritable constitution.

which I have referred, I have found the density sometimes increased to 1·035, and even to 1·040; and very usually to 1·025, and 1·030. It possesses the natural acid character, invariably, when in the recent state; but such highly animalised urine soon becomes putrid and alkaline. It proves an unnatural stimulus to the bladder and urethra, exciting much irritation; and sometimes in a degree to cause the apprehension, that a calculus obstructs the passage, or that a stricture exists.

A deposition of pink or brick-dust sediment on the cooling of the urine, is of such ordinary occurrence when any active symptoms of gout are present, that its connexion becomes forcibly impressed on the mind of the patient, and he gives it the name of gouty urine. A copious precipitation of mucus is invariably an attendant on either of the sediments in question, partly intermixing with it, and in part forming a distinct stratum upon it*. This mucus also appears in the form of a pellicle on the surface of the urine, and stains the sides of the vessel with a milk-like fur. In some of its modifications, it has very much the appearance of floating oil. However much the urine may be loaded with these coloured and mucous sediments, it is transparent when first passed from the bladder, and does not, according to my experiments, become turbid until it has cooled down to 63 or 62 degrees of Fahrenheit. By the application of heat it becomes completely re-dissolved, when the temperature is raised to 98°. The whole sediment in question, uric, mucous, &c. adds but little to the specific gravity of the

* The mucous stratum appears above or beneath the coloured one, according to the quantity and density of the two depositions.

urine. I selected an example in which the pink sediment was unusually abundant, and found, that before filtration, its specific gravity at 60° was 1·0342; after, 1·0332. In the case to which this experiment refers, no tendency to gout belonged to the constitution; and in this part of my subject I shall only thus briefly advert to it, as one among numerous proofs which I could produce, that the appearance of these sediments is entirely dependant on a faulty state of the digestive organs; and upon unhealthy assimilation. It sometimes happens that the sediment changes from a red colour to white; and a considerable portion of mucus also accompanies this kind of deposition. When this happens in a slight degree, the urine is transparent on being first passed, and precipitates its sediment only on cooling. Whether gout be present, or whether a patient, free from the tendency to that disease, be affected with the disposition to secrete urine of this character, we may consider not only that the digestive functions are irregular, but that the different secreting organs are much influenced by a nervous state of the constitution.

When the sediment assumes a crystallized appearance, the state of complaint comes under the well-known denomination of *gravel*. I may observe, that the information communicated by patients on this point, is given with much incorrectness; some declaring that they are affected with gravel, if any deposition of the pink or brick-dust sediment appears; while others do not connect such an opinion with its appearance, although abundant, unless they suffer painful irritation in passing the urine; but when the crystals are seen, the representation that the gravel is affecting the kidneys or bladder, is invariably stated. Of these urinous precipitates, I have now to offer a few observations.

Any elaborate view of this important subject would be incompatible with the limits of this volume; but as the crystallized sediments of the urine now to be considered, fall equally with the preceding, under the head of a morbid state of the digestive organs, a brief description of their external character may not be altogether uninteresting.

The establishment of a correct pathology respecting them, is of all possible importance; as, in addition to the painful symptoms of irritation which they occasionally produce in the kidney and bladder, they not unfrequently augment, from neglect or ill treatment, so as to lead to that most painful of all diseases, *the stone*. These depositions may indeed be said to constitute the stone *in miniature*.

It happens, without any exception, that gouty persons are, at some period or other, either affected with gravel, or, with the deposition of the pink or brick-dust sediment. I have met with many instances in which the patient has related that he was painfully affected with gravel, before he became gouty, to the degree indeed of apprehending the sufferings of the stone; but since, has only noticed the thick and coloured sediment. Some experience the gravel in the interval of the paroxysm, but very few are so affected when gout is present; and an explanation of this phenomenon, I shall hereafter endeavour to suggest.

The appearance of that species of gravel, which, from its physical resemblance to small gravel or sand, has given rise to this familiar term, varies in several particulars. It is either angular and shining, and of an orange yellow colour; or very minute, orbicular, shining, and of a bright red; or of irregular form, dull, and of a yellow or reddish colour. With the two first varieties, the urine

does not contain much suspended mucus; but with the last, more in quantity. This last, indeed, is the intermediate kind between the angular bright crystal, and the pink or lateritious sediment. The urine which possesses either of these deposits, invariably reddens litmus.

The next species of gravel is not only of a white and shining appearance, but is altogether free from coloured deposit. These crystals vary like the former in size; are sometimes angular, and in fragments of no inconsiderable size; but much more usually, they are in the state of very fine lamellated scales, and shine with silvery lustre. When they are very minute, and in small quantity, they do not appear visible until the urine has been some time at rest, when they become aggregated. When these minute shining crystals alone appear, the urine is transparent, and occasionally, as I shall mention at a future page, it is alkaline, and communicates a brownish red to turmeric paper. It usually happens that a mucous pellicle appears on the surface, entangling many of the minute crystals, which so refract the light, as to afford prismatic colours. The urine, in this case, is transparent, when passed from the bladder. When a very copious white sediment appears, partly crystallized, and partly in the state of rough powder, involved with much mucus, the urine itself is turbid, when first passed; but, according to my observation, it reddens litmus. The mucus in this sediment, as in the coloured uric, subsides both intermixed and in a distinct stratum; and sometimes appears to the eye, firm and gelatinous. On the surface of the urine, the floating mucus a little resembles oil. It now and then occurs that the two species are mixed; each partly crystallized, and in part in the state of powder; and a further variety is sometimes presented in the appearance of the crystals, from their being blended with

dark-coloured animal matter, which is either greenish or blackish, in addition to the mucus, which in this case is very abundant. In short, so far as my experience has gone, the sediments in question present as much diversity, as the formed calculi of the bladder, and require the same pathological reasoning, and medical treatment. These points, together with some discussion of the chemical composition of the respective deposits, are to be considered in a subsequent division of the subject.

I return now from this long digression, and resume my more immediate treatise.

OF THE EXCITING CAUSES,

Excessive Intemperance.—THE sudden repletion, and inflammatory action of the general circulation, which follow extreme indulgence at the table, sometimes introduce a fit, in the course of a few hours, where the predisposition strongly exists. It rarely fails to occur, if the excess be much repeated; and now and then, a debauch of the stomach appears to be the sole occasion of introducing the first fit.

One patient, neither descended from gouty parents, nor having reason to expect such a disease, after three or four days of successive conviviality, in which he drank *champaign* very freely, was seized severely with the gout.

In another instance also of acquired gout, the patient, by his external characters, not appearing predisposed to the disease, suffered a first attack, which had a very

evident connexion with the frequent and free use of champaign.

One, of a temperament bordering on the pure sanguineous, with a circular chest and large veins, not born of gouty parents, after committing the extraordinary excess of drinking four bottles of port-wine at a sitting, was seized on the same night, and for the first time, with the gout.

In those who have long been gouty, any great excess in wine is a pretty certain fore-runner of a fit. A gentleman, for a few years subject to the disease, in whom the predisposition was hereditary, relates to me that on one occasion, after strong horse exercise, and supposing himself, from his comfortable feelings, quite secure, he drank a bottle of claret, in addition to some general indulgence at the table; and on the following morning, the gout took severe possession of both ankles. Although wines of any kind which are of bad quality, give rise to certain injury in the gouty patient, yet it is a familiar fact, that the excitement of a paroxysm will much more probably happen from a wine abounding in acid, as *champaign*. The stomach being over-excited by the stimulus of the liquor, derived from its alcohol, is weakened, and cannot resist the acetous fermentation, which its acid nature excites; and hence a double source of irritation is produced.

The heating, exciting, and certainly injurious qualities of champaign, with regard to gout, are much more remarkable than are found from any other liquor. If it has power to excite a first fit, we need not wonder that it is a fruitful occasional cause, in producing the returns of the disease.

I have met with several instances, in which, the patient

thinking, from his feelings of general health, that he might indulge a little with impunity, has taken a few glasses of champaign; but before twenty-four hours have elapsed, the gout has followed.

A gentleman, much disposed to gout, in the summer season, persuaded at the time of his security, drank six or seven glasses of champaign, at a convivial dinner; and in twelve hours he paid the penalty of his indiscretion, in the arrival of a decided fit: which, however, was not of long continuance. He had never before been attacked in summer. Some are so susceptible to the influence of champaign, that even a single glass very soon produces some sensation of gout in the limbs. One gentleman assures me, that either by the slight use of this liquor, or even of perry, he could, at any time, bring on more or less of gout in his feet.

If the error of drinking champaign be committed on one occasion only, and not in great excess, I have usually seen that it has given rise only to a fit of short duration. The effect, however, must depend on the state of the digestive organs, and the accidental degree of the gouty diathesis at the time*.

The more habitual use of liquors of this class, and in proportion as it can be carried on long without exciting the fit, produces effects which are more lasting and severe. I saw lately a gentleman, who states, that, in his last visit to France, he indulged every day for some time in the

* I have now and then, yet very rarely, met with instances, in which the gouty patient has not considered himself to be injured by occasional indulgence in champaign, even to the extent of two or three glasses. Such exceptions should not, however, give any encouragement to the use of this luxury.

light wines, and without seeming disagreement; but on his return home, he was attacked with gout, which, with its immediate consequences, lasted *forty weeks*.

When a fulness of the general habit prevails, the patient feels heated, as he expressively describes it; and when the gouty diathesis is present, we see the active influence of stimulating liquor as an exciting cause of this disease, very strongly exemplified.

Briefly to illustrate this statement, I shall mention the following instances :

A gentleman, deriving the gout only remotely from his family, on two occasions took a pint of strong ale, and a little port wine after dinner. Each time, in a few hours after, he felt tinglings, pricking and shooting, in the great toe, and two of the other toes.

One gentleman expresses his certainty of having gout, if he takes both malt liquor and wine, although only in moderation.

Another, that if he drinks port wine for a week, although but sparingly, he finds the great toe hot, throbbing, stiff, and very painful.

A gentleman, who has had the slightest visitation only of gout, finds that merely two glasses of port wine, if it be of a strong quality, immediately produce pricking pains in the toe.

The gouty in inferior life, who cannot afford to excite a fit by wine, produce the effect with strong malt liquors and spirits: for in this way, also, indirect debility of the stomach is produced, acid matter is abundantly generated, and the inflammatory diathesis of the system takes place.

In this metropolis, examples continually come before me, of persons, whose indulgence consisting in the very

free use of porter as a daily habit, and in the drinking of gin also, not to the extent of great intemperance, produce the gout, entirely as an acquired disease, without any hereditary influence.

A man, not apparently predisposed to gout, kept a public house and drank usually two gallons of ale every day, some brandy and water, and a little wine. He scarcely took any exercise, and acquired the gout severely.

An eminent Physician informs me, that he met with an extraordinary prevalence of gout in a country town, in which it was the custom at the best tables, to drink freely a very potent ale, and scarcely any wine.

I may add to this general account, that in most of the instances, in which, the sudden seizure of gout takes place in the day time, the circulation has been actively excited by stimulating liquor, and the attendant provocatives of the table. In three distinct cases, I learnt that the patient sat down to a convivial dinner, with scarcely the sensation of gout; but on rising to depart, the feet were completely disabled by the inflammation and swelling, which had made such rapid progress.

Acidity, when much accumulated in the primæ viæ, will always powerfully concur with other causes to excite a fit, and will sometimes prove alone sufficient. During the paroxysm, if occurring in a considerable degree, it is always a cause which produces much aggravation of the symptoms. When rejected from the stomach, it is very acrid to the throat; and this may be considered as a proof of its irritating power in the stomach. As I have already observed, it is often of a grass-green colour, sometimes yellowish green, and now and then, but more

rarely, colourless*. In proportion to its acrimony, mucus is mixed with it more or less abundantly. In the first volume of the Medical Observations and Inquiries (p. 41), an interesting case is related, in which, this kind of green acid vomiting proved critical to the paroxysm of gout. I have met with several instances, in which the discharge of acrid fluid from the stomach has produced immediate and very sensible relief; but I do not remember any example of such a complete removal of the symptoms following, as described in this case.

* This fluid appears to be composed of the peculiar acid matter generated in a morbid condition of the stomach, together with bile and mucus. I believe that the green colour often arises from a vitiated state of the bile; but may also in part be referred to the change produced in it by the acid matter. What the nature of this acid is, seems questionable. An artificial admixture of acetic acid and bile in any proportion, does not produce the green colour; but it is readily affected with diluted muriatic acid and bile. I added recent bile to some colourless acid fluid, recently discharged from the stomach, in the proportion of one part to sixty, and a light green colour immediately appeared, which, on standing, became grass green. If the bile be added in any considerable proportion, the yellow colour predominates; and in the same proportion that this occurs in the stomach, the patient complains of a bitter taste. The green fluid in question rejected from the stomach when in a weak and irritable state, is very weakly acid, as is proved by the slight degree of neutralising power which it exerts when added to an alkali.

In the specimen now mentioned, two drams were rendered neutral by three drops of the liquor potassæ of the Pharmacopœia. The patient's sensation of intense acidity, in this case, therefore, must be connected with a morbid sensibility of the surface of the œsophagus, continuous from that of the stomach.

Obstruction, and Excess of Bile.—A jaundiced skin, with general fever, pain in either hypochondrium, costive bowels, with bilious vomiting; or, acrid bilious diarrhœa; and scanty, deep-coloured urine, sometimes occur as the short precursors of a paroxysm.

A gentleman, whose gout was not due to hereditary influence, had been very industrious in gardening on a hot day in autumn. In the evening he was seized with bilious vomiting and diarrhœa; and on the following day, the gout appeared in one foot.

Cold, with or without wet, applied to the body generally, or to the lower extremities only, especially when in concurrence with fatigue, proves, in an equal degree, exciting to the gout in an individual who is predisposed to the disease, as to the phlegmasiæ in general; and it is by far the most frequent of the exciting causes. The act of walking in thin shoes on damp ground is always hazardous. One gouty patient informs me, that on several occasions, when in the occupation of his business (surveying buildings), he has been standing for some time on cold, damp pavement, his bowels have been quickly attacked with pain, with subsequent constipation; and the gout has followed in two or three days. This same patient was formerly in the habit of getting wet in the lower limbs with salt water (in an employment at sea), and continuing so for several hours, with impunity. I should add, however, that at that time he had never suffered the gout. I know several gouty patients, whose sensibility to the immediate influence of the east wind, or to the act of standing on damp pavement, or very damp ground, is almost incredible! We see that cold, whether applied locally or generally, acts most powerfully when conjoined with wet; but certainly the

east wind*, by itself, is a severe and active agent. On visiting a patient just recovering from a paroxysm, upon an afternoon when the wind had suddenly changed from the east to the south-west, I was amused by the very emphatic description which he gave of the happy alteration in his feelings. It appears to me, that, of all the forms in which cold acts upon the body, a wet fog, with a penetrating east or north-east wind, appears to be most certainly injurious. It may be useful to mention the following facts as affording cautionary hints.

A gentleman, subject to a severe attack once every year, having had his accustomed fit, some time after rode in a damp and cold state of the atmosphere, returned home rather chilly, and unguardedly changed his clothes without putting on his flannel gown, as usual, while dressing. This continued operation of cold excited gout in both feet on the following day. On a subsequent occasion he suffered the same consequence from riding on the outside of a coach, insufficiently protected by clothing, when the wind was in the east.

Another traced an attack to the simple act of putting on stockings which were rather damp.

Another, who was just convalescent from a first and severe attack of gout, imprudently walked on the wet pavement in showery weather. On the following day, he suffered a severe relapse; but the symptoms entirely lost their former regular character, and assumed as much the appearance of rheumatism, as of gout. It would be almost endless to enumerate the cases in which a continued exposure to wet and cold has excited the paroxysm,

* Dr. Gregory, in his lectures, observes, that the genial south and west have alone been invoked with praise by the poet.

either as the immediate consequence, or as a sequel to some other of the phlegmasiæ.

In either case of the application of cold, whether in conjunction with wet or not, its first effects appear to be very general; but injury soon becomes manifested in the weakest part of the body. The capillary circulation is checked near the surface, the blood flows to the internal parts in preternatural quantity, and congestion in the weakest organs, or most morbid part of structure, is consequently produced. The secretions become disturbed and vitiated; and the nervous system partaking in the irritation, many sympathies, and various sensations of disease, are the fruitful consequences.

External Injuries.—Strains, concussions, contusions, or any mechanical violence offered immediately to the part which becomes affected with inflammation, or even to some other part of the body, very commonly prove exciting causes of a paroxysm. Even a first fit is sometimes excited in this manner, and the patient imagines that he labours under a common strain, only.

One gentleman suffered his first attack, which was in the foot, after unusual exertions in dancing, being at the same time under the influence of the effects of a convivial entertainment. In another case, an accident, by which one leg was broken, gave rise to severe gout in the other leg, on the following day.

A female, whose gout was *acquired*, had her first fit excited, in the great toe, by a severe blow on the foot. I am acquainted with seven other cases, in which the first attack was imputed to some local contusion or strain; and was treated at the beginning, by the usual means employed for such accidents. Under the head of my present subject, I may add the following facts:

A gentleman, much subject to gout, when considering himself unusually well, underwent the slight operation of having the nail of the great toe cut, on account of its improper growth. The toe was much pressed, and gouty inflammation was the immediate consequence.

In another case, the patient, never before having suffered the gout, tore off a broken portion of the thumb nail, so as to make the part tender. Very soon, the thumb and part of the hand put on a swollen and shining appearance, and was exquisitely painful. A poultice was applied. Suddenly, on the third evening, the pain quitted the thumb and seized the toe, next the ankle, then the knee, and lastly the great toe of the other foot. Throughout, he procured ease and sleep as the first light of the morning appeared; and hence facetiously observed, that the gout, in this respect, assumed all the behaviour of a *ghost*.

A gentleman had the misfortune to break his thigh by a fall from his horse. In a week after, he was attacked with gout in the great toe of the foot of the other limb. He described the pain of the gout to be infinitely more severe than that which he had suffered from his accident.

I have seen one case, and been made acquainted with the particulars of another, in which a paroxysm followed in a few days after the operation being performed for cataract. The patient was previously well, and, as may be supposed, not having the least expectation of gout; which was evidently called forth by the influence of the constitutional disturbance from the operation. It is with much pleasure, however, I add, that no untoward symptoms occurred to the eye, and those of the gout were very tractable.

A very gouty patient had his great toe trodden upon by a heavy person, and a paroxysm began in the part on the following day.

It seems necessary however to the effect of local injury, that the *constitutional* disposition to an attack should exist. In a gentleman for many years severely subject to gout, the knee received a severe contusion in a fall, and the muscles were violently strained, but only common inflammation followed. Another gentleman who had suffered repeated fits of gout during eight years, strained his foot violently; but the accident did not give rise to any gouty inflammation.

The following interesting illustration of this principle of pathology lately came under my observation. A gentleman, gouty for many years, received a severe blow on the knee, which produced the tenderness usual from such an injury for two days. It then recovered completely. A week after, having two days previously indulged in champaign, he was seized with gout in one foot; and before forty-eight hours had elapsed, the knee, so recently bruised, became affected. On all former occasions the disease had only occupied the feet.

When the gouty diathesis is present, any considerable degree of local injury seems with certainty to be followed by a fit. A gentleman who had been subject to gout fourteen years, at a time when he considered himself in more than usual health (yet living very freely), strained his ankle, in slipping from a stool. On the same day, that foot was attacked with gout, which continued a fortnight. The other foot next became affected, and was under the disease three weeks. This was the most severe fit which he had ever experienced. Even a fall, as from a horse, giving only a general shock to the frame, sometimes proves an exciting cause. I have known three instances of this, in which the gout took place on the following day, although no injury was received on the parts which became affected.

The pressure of too tight a shoe, and more especially if the patient walk much under such irritation, will occasionally induce gouty inflammation. That degree of straining of the disorganised and weakened ligaments and tendons, which appertains to an excess in walking always proves injurious, and sometimes excites a fit.

A corpulent and elderly lady, seldom using exercise, from a pressing necessity walked a distance of four miles; accidentally, also, exposed to rain. She considered herself in unusual good health. On the following day, however, the gout seized both feet; and, afterwards afflicting the knees, continued four months. In this case, there was certainly the two-fold cause of fatigue and cold; but she described, that she felt such immediate sensible injury from the over-exertion, that she judged the consequences to be principally ascribable to it.

One patient assures me, that he is convinced, from the occasional injury which he has experienced, that he could at any time produce an attack, by over-walking in any extraordinary degree.

I may add, that any of the modes of local injury, which I have enumerated, will powerfully operate to produce a relapse, at a time when the convalescence has begun apparently in the most favourable manner.

A gentleman, much subject to gout, informs me, that when quite convalescent from the paroxysm, he received a kick upon that leg of which the foot had been affected; and a consequent return of gout immediately followed.

It happens very commonly, that premature exertions in walking, just as the paroxysm is taking its leave, produce a relapse.

I should before have mentioned, how remarkably local injury predisposes a particular part to be affected with gout, at a distant subsequent period.

I might relate many cases in illustration of this position, but I shall limit my narration. A gentleman, when a youth, accidentally received some small shots from a gun in one knee. They were removed, but the knee was rendered permanently weak in a slight degree; and to this part, the gout in after-life first attached itself, and always afterwards with more severity than in any other situation.

Another gentleman, many years before he had the gout, injured his ankle by a fall from a coach. This ankle was first visited by the disease, and was in all the fits much more affected by pain than any other part.

I have indeed often witnessed that a part, which has been the seat of former accident, has caused the severest suffering in the fit.

Fatigue and Anxiety.—That fatigue, which is the joint offspring of mental anxiety and bodily exertion, is a very common cause of exciting a paroxysm. I have lately attended two female patients, who had suffered the gout in its severest form; one of them for the first time; brought on from the nervous exhaustion which they had sustained, each in nursing a relation dangerously ill, several nights in succession. Late hours, with intense application to business, and in proportion as the feelings are anxiously concerned, lead with certainty to an attack, when the predisposition is only moderately strong. I have seen many proofs of this fact. A fatiguing exercise of the attention is sometimes injurious, even when the mind is agreeably engaged. Thus, one gentleman, much subject to gout, assures me that he has more than once excited a fit by a game at chess.

The Passions of the Mind.—Although in most of the

instances related by authors, the strong passions have rather had the credit of *curing*, than the reproach of *causing* the gout; yet we may believe, that on some occasions, they will serve to excite a paroxysm*. One patient informs me, that some of his worst fits have immediately succeeded a violent irritation of temper; and in other examples, some mental shock has been quoted as the cause of attack. In such cases we may consider, that the system is at the time in a state of great readiness to take on the disease; the nerves are morbidly susceptible in a high degree; and the temper also is more irritable. In the fit itself, the irritability of disposition is almost proverbial with every author.

Sydenham observes, indeed, “that every paroxysm may be as justly denominated a fit of anger, as a fit of the gout.”

Van Swieten relates, *I had long an intimate friendship with a person of very great learning, and other-ways of a sweet, peaceable, and mild disposition, who, taught by his experience, could at last foretell when he was to have a fit, by his being peevish a little before, and out of humour with every trifle.* I must, however, add to this account, which will not appear very flattering to the gouty patient, that I have met with many happy exceptions, and have seen a serenity of mind preserved throughout all the sufferings of a severe paroxysm.

* The active influence produced both on the appetite and digestion, by violent emotions of the mind, is constantly exemplified. The secreting action of the liver is also affected very suddenly by the same cause. Horace, in his description of jealousy, has, with very little of poetical fiction, noticed this circumstance :

“Fervens difficili bile tumet jecur.”—Ode xiii.

THE PROXIMATE CAUSE.

THIS subject of investigation has, with regard to diseases in general, proved an *ignis fatuus* to medical theorists; and the remark may be strongly applied to the case of gout. Aretæus considered, that the proximate cause was known only to the gods. By less pious inquirers, ingenious speculation has been exercised, and profusely substituted in place of solid principles; and various opinions built upon hypothesis (that is, upon conjecture without proof, or even reasonable evidence), have been at different times advanced, with all the pride and confidence of truth.

The uric-concretions (or *chalk-stones*, as they are improperly called) which belong to the gout in a few individuals, appear to have been the chief source of the doctrine of a morbid matter, from Hippocrates to the present time.

A *materies morbi* has been differently conceived by every new author, and amongst the various denominations assigned to the supposed substance, I may mention the following:—*phlegm; acidity in the semen; bilious humour; mucilage; tartareous, or urinous salt; an earth; a volatile alkali; an æther; a superfluous part of the chyle;* and the list might be greatly extended.

Some writers* have contended that an excess of uric acid, or its elements, in the system, is the *proximate* cause of gout. This inference seems to have been de-

* Forbes, in a "Treatise on Gravel and Gout, London, 1787;" and Parkinson, "On the Nature and Cure of Gout, London, 1805."

duced, from the occasional instances of the external uric formations, already noticed; from the presence either of pink or lateritious sediment; or the reddish crystals (commonly called *gravel*), so generally found to exist in the urine in connexion with the gouty paroxysm; and further, from the remark of Berthollet, "That paper coloured blue by litmus, becomes red on being exposed to the perspiration of a part affected with the inflammation of gout*." Dr. Wollastan mentions, that "In gouty persons there is always a redundancy of uric acid†." In reference to a similar theory, and upon chemical principles of practice, Sir Everard Home and Mr. Brande advised the employment of magnesia in gravel, and also in gout‡.

In respect to the uric concretions, as a consequence of gout, it must be observed, that their formation is, comparatively, so rare an occurrence, and so much confined to certain peculiar habits, that the phenomenon cannot be assumed as a basis for the construction of a general theory of the disease; and much less, can this singular morbid process be considered as its *proximate cause*. In two hundred and six cases of gout, I have not discovered any trace of this kind of concretion, in more than twenty-one persons.

While on the one hand, I have also to observe, this process is sometimes described as a common effect of highly inflammatory gout, being thus considered as a general circumstance; it is on the other asserted, that it occurs occasionally, where no previous evidence of gout

* Parkinson, p. 22.

† Philosophical Transactions, 1810, Part II.

‡ Ibid. Part I.

in the constitution has yet existed. The error of the first conclusion I have already pointed out; and I have sought, without success, for an example, to prove the correctness of the last opinion. In all the cases which have come within my knowledge, gout has more or less preceded.

The sediments in the urine, to which I have adverted, are not the peculiar attendants of gout; but occur under various kinds of disease, both constitutional and local, in connexion with a morbid state of the digestive organs.

Upon the question also of an excess of uric acid secreted by the kidneys, as indicated by the deposition of the crystals, or the pink or lateritious sediment, and its supposed relation to the proximate cause of gout, I may state, of the crystals more particularly, that they are not to be considered as a proof of an excess of uric acid; but rather as a separation of this principle from the urine, and a new combination with some other of its elements. If nitric or muriatic acid be added to urine of moderate specific gravity which has deposited these crystals, scarcely any further uric precipitate is produced; and having in different experiments carefully estimated the quantity so obtained, with that procured from healthy urine of the same specific gravity, by the same methods, I have found the position in question to be fully manifested; equal portions of such healthy urine, furnishing an equal quantity of uric precipitate, with the whole of that spontaneously and artificially precipitated from the morbid urine, with which it was compared.

Such urine as deposits the pink or lateritious sediment in an abundant quantity, is, according to my experience, always of high specific gravity, and from such specimens I have invariably procured a much larger *total* quantity of uric acid, than from any urine of moderate specific gravity; but I must add, that if this sediment be

in small quantity, and the specific gravity of the urine be not considerable, almost the whole of the uric acid which such urine possesses, is found by experiment to have been precipitated in the sediment; further confirming the argument, that the relative quantity of uric acid in the urine has a great correspondence with its specific gravity, and cannot be so much inferred from appearances, as we have been hitherto taught to believe.

As such sediment, and in abundant quantity, is a very common occurrence in the urine secreted during a paroxysm of gout, a basis appears to be presented for the assumption of the theory under discussion, in relation to the proximate cause; but it must first be taken into consideration, that such sediments are neither necessarily nor regularly attendant on a paroxysm of gout, and that they are found, as I have before stated, under other circumstances of disease, in connexion with unhealthy chylopoetic functions. In proportion, therefore, as gout is associated with such disordered functions, and not further, are these urinary evidences connected with that disease.

An important fact is also to be added, of which I have obtained very abundant proof by experiment, that not only the uric acid is in excess, in urine of high specific gravity, in the examples of the gouty specimens to which I have now alluded (and also in urine of similar density in other diseases); but, other principles composing this secretion, as the phosphoric, sulphuric, and muriatic acids, urea, &c. are also secreted in preternatural quantity.

An opinion suggested by Berthollet with regard to the peculiar nature of gouty urine, is of such importance, that it has engaged my particular attention. The following is a translation of the original, as quoted by Barthez. *That from numerous observations, he was con-*

vinced, that the phosphoric acid (which is always in the urine combined in excess with calcareous earth) is naturally in much less quantity in the urine of persons subject to gout and rheumatism, than in healthy persons; but that at the approach of a paroxysm, and during its continuance, the urine contains as much of phosphoric acid, as that of persons of strong constitution, and much more than belongs to the gouty in their ordinary state†.*

I have submitted the truth of these doctrines to the test of careful experiment; and the results of this investigation, I now offer in the annexed statement.

Richerand, in his *Elements of Physiology*, advances an opposite theory, and observes‡, “On the approach of fits of the gout, the phosphoric ingredients of the urine diminish, and seem to be carried to the joints, to produce, in their vicinity, arthritic concretions.”

* Berzelius has pointed out the error of the supposed existence of free phosphoric acid in the urine, in the following words: “As by the laws of chemical affinity, these acids will unite with any alkali that may be present, and saturate themselves with it in the order of the force of their respective affinities, it must follow, that where the quantity of alkali is insufficient to saturate all the acids present, the weakest acids must be those that will remain uncombined, and will give the urine its acid properties. These, therefore, must be the lactic and the uric.”—*Medical and Chirurgical Transactions*, vol. iii. p. 257.

† *Traité des Maladies Goutteuses*, tome i. p. 50.

‡ Translation by De Lys, p. 141.

FROM FOUR OUNCES OF URINE.

EXP.

	Sp. gravity.	Precipitate by Nitr. of lead. Grains.	Ph. of lead. Grains.	Ph. acid. Grains.
I. J. W. under a paroxysm of gout; urine of orange colour, with minute uric crystals, and some mucous cloud	1.016	22	13.7	2.88
II. Ditto, in health, and two months after recovery; urine natural.	1.0199	18.4	4.6	.96
III. A. L. under a paroxysm of gout; urine of deep orange colour, with copious pink sediment and mucus	1.028	66.2	43.7	9.2
IV. Ditto, in health, and two months after recovery; urine natural.	1.0168	26.3	14.1	2.97
V. J. C. under a paroxysm of gout; urine of amber colour, and with mucous cloud only, but precipitates albumen with nitric acid	1.014	28.8	4.8	1.02
VI. Ditto, in health, and three months after recovery; urine natural	1.0137	16.2	4.3	.91
VII. C. M. under a paroxysm of gout; urine of deep orange colour, with copious pink sediment and mucus	1.020	50.5	20.2	4.3
VIII. Ditto, in health, ten weeks after recovery; urine natural	1.0107	23.4	7	1.47
IX. E. P. under a paroxysm of gout; urine of deep orange colour, with very copious pink sediment, and mucus	1.029	71.5	30.4	6.41
X. Ditto, in health, ten weeks after recovery; urine of orange colour, with uric crystals, and mucous cloud.	1.0191	40	19.2	4.05

FROM FOUR OUNCES OF URINE.

EXP.		Sp. gravity.	Precipitate by		Ph. of lead.	Ph. acid.
			Nitr.	Grains.	Grains.	Grains.
XI.	W. W. (see Case I.) one day before an attack of gout, feeling in health ; urine of amber colour, with some uric crystals and mucous cloud....	1.0094	13.8.	3.8	.8	
XII.	Ditto, second morning of the paroxysm ; urine of deep orange colour, with copious lateritious sediment and mucus.....	1.0201	55.2	25.4	5.36	
XIII.	Ditto, seventh morning ; urine of amber colour, with mucus, but a trace only of the pink sediment.....	1.0099	30	15	3.17	
XIV.	Ditto, eighth morning ; urine of deep orange colour, with dirty brick coloured sediment, and much mucus.....	1.0242	46	18.4	3.88	
XV.	Ditto, ninth morning ; urine of amber colour, with the mucons cloud of health, but no other sediment.....	1.012	24	4.8	1.02	
XVI.	Ditto, tenth morning ; urine as yesterday, with the addition of some minute uric crystals.....	1.0105	28	7	1.47	
XVII.	Ditto, eleventh morning ; urine of light straw colour, with minute uric crystals, and more of mucus than yesterday.....	1.0106	32.8	16.4	3.47	
	Ditto, twelfth morning ; urine natural.....	1.0085				
XVIII.	Ditto, in health, about two months after recovery ; urine natural.....	1.0172	26.8	13.3	2.8	

FROM FOUR OUNCES OF URINE.

EXP.		Sp. gravity.	Precipitate by		Ph. of lead.	Ph. acid.
			Nitr. of lead.	Grains.	Grains.	Grains.
XIX.	J. G. (see Case) hepatitis, with gouty diathesis; urine of deep orange colour, with copious pink sediment, and mucus.....	1.0207	57.6	19.5	4.1	
XX.	Ditto, eleven months after the former period; digestive organs not in sound health; urine of orange colour; with abundant uric crystals, and mucous cloud.....	1.014	20.4	10	2.1	
XXI.	J. W. aged 45, under a paroxysm of gout; urine of deep orange colour; with pink sediment and uric crystals, and much mucus.....	omitted	42.5	16.1	3.4	
	Ditto; no opportunity of comparison.					
XXII.	J. M. aged 38, under a paroxysm of gout; urine of amber colour, with some pink sediment and mucus.....	omitted	35.6	10.6	2.24	
	Ditto; no opportunity of comparison.					
XXIII.	T. A. aged 37, in health; of free habits; not subject to gout or rheumatism; urine natural.....	1.0113	25	13	2.75	
XXIV.	L. S. aged 37, ditto; of free habits; ditto.....	1.0183	40.8	20.5	4.33	
XXV.	A. J. aged 24, in health; of free habits; subject to violent acute rheumatism; urine natural.....	1.0173	17.4	5.6	1.18	

FROM FOUR OUNCES OF URINE.

	Sp. gravity.	Precipitate by		Ph. of lead. Grains.	Ph. acid. Grains.
		Nitr. of lead. Grains.	Ph. of lead. Grains.		
EXP.					
XXVI. E. J. aged 19, in health; not subject to gout or rheumatism, but plethoric, and liable to tonsillary inflammation; urine natural.....	1.0217	49.2	20.4	4.31	
XXVII. H. T. aged 48, not subject to gout; suffering under chronic hepatitis and morbid alimentary secretions; urine of deep orange colour, with copious pink sediment and mucus.....	1.0334	49.2	22.8	4.82	
XXVIII. Ditto, in two or three months after, when in improved health from a course of medicine; urine of amber colour, and free from sediment.....	1.018	37.2	13	2.75	
XXIX. J. T. aged 34, a strong man, under severe continued fever; urine of deep orange colour, with much pink sediment and mucus.....	1.025	32	18.5	3.9	
XXX. Ditto, in six months after, in perfect health, and of moderate habits, but very plethoric, and of sanguineous temperament; urine natural.....		48	24	5.07	
XXXI. S. P. aged 39, a robust man, of free habits, plethoric and sanguineous; under acute hepatitis; urine of deep orange colour, with much pink sediment and mucus.....	1.023	38.4	24	5.07	

FROM FOUR OUNCES OF URINE.

Sp. gravity. Precipitate by
Nitr. of lead. Ph. of lead. Ph. acid.
Grains. Grains. Grains.

EXP.

xxxii. Ditto; feeling in general health, two months after; but certainly the functions of the liver not yet correct; living freely; urine of light orange colour, with some pink sediment and mucus..... 1.0245 40 28 5.92

xxxiii. J. B. aged 13, robust, and not of unhealthy appearance; bowels constipated, with much spasmodic pain and nephritic irritation; urine of amber colour, depositing the phosphates, and also holding them abundantly in solution, although it vividly reddens litmus..... 1.018 36.6 17.7 3.75

xxxiv. L. S. (a female) aged 23; robust; under acute rheumatism: urine of orange colour, with slight pink and mucous sediment... omitted 29.2 5.1 1.07

No subsequent opportunity of comparison.

xxxv. L. M. a boy, aged 4, diseased with tabes mesenterica; a large belly, with distension and hardness; narrow chest; cough; irregular fever and debility; urine of amber colour, with much laceritious sediment and mucus..... 1.0263 51.2 23.5 4.96

FROM FOUR OUNCES OF URINE.

EXP.

xxxvi. H. C. aged 42, robust, and engaged in laborious employment; ill with continued fever; urine of orange colour, with pink sediment, and mucus.....

1.023
44.8 omitted
24.6 ditto

xxxvii. Ditto in health; two months after recovery; urine natural.....

xxxviii. A. B. a female, aged 3 years; a well grown child, now very ill with cough and fever, and disordered bowels; the fæces blackish; the urine of deep orange colour, with copious deposition of the phosphates, although reddening litmus in the usual manner: so much loaded with urea, that without concentration it immediately crystallized with nitric acid.....

1.0214
39.2
27.4
5.8

xxxix*. R. T. a female, aged 2 years and a half; of rather a healthy countenance, but suffering from mollities ossium, in a very great degree, both in the upper and lower extremities.....

1.0299
35.6
20
4.22

xl. M. D. female, aged 5 years; robust and of a full habit, and in a state of inflammatory diathesis.....

1.024
43.2
28
5.92

• It has been conceived that in mollities ossium, the kidneys secrete a preternatural quantity of phosphate of lime, depriving the bones of this principle. So far as one example may be quoted, I may refer to this Experiment in contradiction of the hypothesis.

In these experiments, nitrate of lead was the precipitant employed; and the urine was diluted with distilled water. A portion of the urine, first and separately passed in the morning, was, in each case, chosen for examination. The precipitate was carefully dried and scraped from the filter. Of this, a certain quantity was boiled in distilled water, that the muriate of lead might be removed; and with it also the uric acid was separated. It was then burnt in a crucible for about half an hour, that the several animal matters should be destroyed as much as possible. It was next boiled in diluted nitric acid; and being allowed to rest, the clear liquor was decanted from the sulphate of lead and some insoluble animal matter. To this, ammonia was added in excess; the precipitate was collected on the filter, dried, and weighed. This was the phosphate of lead, from which the relative proportion of phosphoric acid was estimated, by means of Dr. Wollaston's logometric scale. Equal quantities being always employed, and all circumstances of experiment being alike, it may, I hope, be presumed, that this investigation, by comparative experiments, is entitled to confidence.

The method which was thus adopted may appear improperly complicated; but I was led to employ it, from the comprehensive opportunity which it afforded me, of judging with tolerable accuracy of the relative quantities of the different principles of the urine in every example. I derived these results from estimating how much the given quantity of precipitate under examination had lost by each process. By the agency of nitrate of lead as the precipitant, the several principles of the urine are almost wholly thrown down. Some of the muriatic solution, with a portion, also, of animal matter, in union with lead, remain in the filtered

liquor. It will probably be suggested as an objection against my conclusions from these experiments, that the urine is a secretion of such variable density, in the same individual in the course of each twenty-four hours, that fallacious inferences, deduced from the particular portion examined, can scarcely be avoided.

In answer to this objection, it can with truth be stated, that if the first urine of the morning, *separately procured*, be made the subject of examination, great accuracy of results will be obtained; and by observation and investigation, carefully conducted, I may assert, that a very faithful judgment may be formed, of the state of the important function which the kidneys are performing.

The general accordance of the specific gravity* of the urine with the principles which were sought for in these experiments, is further satisfactory, as proving the utility of the information, which may be obtained from examination of this point alone, with a view to forming general conclusions, when it is desirable to obtain some judgment, with convenient promptness, of the condition of the secreting action of the kidneys.

With respect to the position of Berthollet, it is manifest that in the urine of persons under a paroxysm of gout, the proportion of the phosphoric acid is greater than in their healthy state; but the same truth is also apparent in Experiment xxvii. in a person whose liver was diseased, and who never had gout. In Experiment xxxii, the same contrast to the former state is not presented, as the patient was under the stimulating excite-

* In all my experiments, I employed the temperature of 60° of Fahrenheit. I found the following variations of specific gravity in the same urine, at different temperatures: at 60° 1·027, 70° 1·0256, 80° 1·0251, 90° 1·024, 100° 1·0201.

ment of free living, and was not quite free from hepatic complaint. In Experiment xxx, the actions of the system were vigorous, under a full circulation, and a sanguineous temperament. In Experiment xxxvi. the case of continued fever, the precipitate by nitrate of lead is as much more in quantity than that obtained in the health of the patient, as in most of the examples of gout. It seems, therefore, that the phenomenon in question cannot be considered as a specific occurrence in gout alone; and we are consequently forbidden from saying, that an excess of phosphoric acid in the system is the *proximate cause* of gout; and the more especially is such an hypothesis opposed, when we take into consideration the general fact already stated, that in urine of high specific gravity in gout, the other principles of this secretion also are in excess. It is incumbent upon us to avoid partial opinions founded on partial deductions.

In regard to the statement of Berthollet, "that the phosphoric acid is naturally in much less quantity in the urine of persons subject to gout and rheumatism, than in healthy persons," it may be remarked, that so far as these experiments apply to the question, this inference must not be hastily drawn. Each of the gouty persons was pursuing more or less a regulated regimen, and may be supposed also to have retained some corrective influence over the digestive functions, and the general secreting actions, from the previous course of medicine; while in the healthy examples which were chosen, and particularly in Experiments xxiv and xxvi, the habits of living were indulgent, and the circulation was plethoric.

In experiments iii and ix, the proportion of phosphoric acid was in great excess, over the examples to which I have last referred; and in the Exp. xi and xii, the comparison is particularly interesting, together with the

subsequent series. Exp. xi points out that the kidneys were deficient in their function of removing matter from the blood requiring to be excreted; and in two days after, a disease was set up in the constitution, for the relief of which, Nature immediately employed her own powers; and the kidneys, in part of the salutary effort, became excited to increased secretory action. In how great a degree, a previous retention in the system of the elements of phosphoric acid, or the other principles to be discharged by the kidney, may be considered in the relation of cause and effect to the gout, it would be hypothetical to contend, if the question of *proximate cause* be considered; but the general fact of an excess of excretion of all the principles of the urine, in the paroxysm, is very apparent.

The theory of Berthollet, of an excess of phosphoric acid in immediate connexion with the paroxysm, as cause and effect, has not only been adopted by others; but has, I know, been carried to its utmost limits, in a comparison drawn between the phenomena of gouty inflammation, and of phosphoric combustion. The Experiments xxxv, xxxviii, xxxix, and xl, do not apply to the present question, as they are not comparative with respect to the individual children; but they serve to shew the large proportion of phosphoric acid, which belongs to the urine of children. It is a curious fact, that the child, whose urine was chosen for Experiment xl, and which possessed such a large proportion of phosphoric acid, was attacked in the evening of that day, when she was considered to be in her usual state of robust health, with a violent inflammation and swelling of half the face. The hypothesis of Richerand is altogether founded upon false principles. According to my experience, and so far as I have ever learnt, the phosphoric acid does not enter into the composition of the arthritic concretions.

The proximate cause of a disease must not only be something which is invariably antecedent, but also that which is distinctly peculiar to such disease; and the whole of this position I am not able to demonstrate; although some general points are, I trust, satisfactorily proved.

The results which the preceding experimental investigation has afforded, appear to me interesting, both in a physiological and pathological point of view. They exemplify the activity of the kidney as a salutary organ of excretion, and as constituting a very material agent, which Nature often employs to relieve the overloaded system. In this view of the subject, at the same time that I consider the excretory function of the kidneys as really a part of the curative process which Nature is attempting to accomplish, I do not mean to contend, that the *specific cause* of the disease is in this way removed; for such a conclusion would be at variance with the results of my experiments. The investigation appears to me of additional importance with reference to the doctrines of the humoral pathology, which have probably been too violently discarded in modern medicine. Notwithstanding, however, that I advance this sentiment, I would not be understood to lean to the extravagant notions of the mathematical sect of physicians, Langrish, Bryan Robinson, &c. who offered very refined calculations of disease, deduced from the state of the blood. Also, in admitting or contending, that the different fluids of the body vary in their chemical composition, in health and disease, we should abstain from all particular conclusions, which are not founded on careful experiment; and we should studiously shun the wild hypotheses of the humoral pathologists in general.

In pursuit of my present immediate inquiry, I have

now to notice some further theories which have been advanced, as attempting the perfect explanation of gout. The acid nature of the transpiration, as indicated by the reddening of litmus paper when applied to the skin under gouty inflammation, has been improperly quoted to suit the theory of an acid matter being a proximate cause; for the same fact may be constantly demonstrated in the healthy action of the skin. In numerous trials which I have made to this effect (in cases even of long standing jaundice), I have not found an exception*.

Barthez, a reputed French author on the subject of gout, appears to me obscure and perplexed both in his theory and practice; but his work contains much useful and interesting reference, and claims the attention of the medical reader.

He considers that the production of gout depends on two causes; the one, *a particular disposition in the constitution to produce a specific gouty state both of the solids and fluids; the other, a weakness (natural or acquired) which the parts appearing as the seat of the disease, suffer relatively to the other organs.* He does not attempt to explain a proximate cause. His hypothesis of *la situation fixe, in the muscular fibre and other textures*, does not convey to me any comprehensible idea. He entertains, after the example of Van Helmont and Van Swieten, the absurd notion, that the gout is a *contagious* disease. Amongst his varieties of gout, he includes the hot and the cold, according to the authority of Aretæus and Cælius Aurelianus, and from

* Berzelius remarks, "The matter of transpiration is always acid, and reddens litmus paper very distinctly."—View of the Progress and present State of Animal Chemistry, p. 95.

the example of Liger, a preceding French author. In this doctrine, he refers the *hot* kind, to the disease when seated in the neighbouring parts most external to the joint; and the *cold*, accordingly as the disease is more internal and deeply fixed; grounding this distinction also, on the particular effects produced by hot or cold applications to the affected part.

M. Guilbert, the latest French author* upon gout, appears to me to have exceeded every other writer, in the *universality* of character which he has given to this disease; and in following his opinions, we should very readily range every nervous and spasmodic complaint, of doubtful nature, under the head of anomalous gout. Also, some of the complaints of young children; various disorders of the adult; uterine irregularity; and the diseased condition of every texture of the body in a person gouty, or supposed to be so; would come within the convenient denomination of arthritic. In explaining the nature of gout, M. Guilbert states, that under the influence of causes which have impaired the functions of digestion and perspiration, and given rise to a condition of plethora, a matter, intended to be excreted, has remained; the lymphatic system is left obstructed (engorgé) with this matter, which becomes the matter of gout†. He seems quite contented with this easy assumption of *cause*, and next describes the various affections, which the matter in question produces. Of the practice recommended, I shall occasionally take notice. The work is certainly deserving of an attentive reading; although, according to

* De la Goutte, et des Maladies Goutteuses. Paris, Jan. 1817.

† Ibid. p. 195.

the impressions made on my mind, it abounds in romantic opinions, and unsound pathology*.

Dr. Sutton, in a concise, but instructive Essay on Gout †, appears to believe that a peculiar secretion in the alimentary canal is the proximate cause; and that the disease, in its origin, is very local in its foundation. He remarks, "The symptoms attending gout give every reason to suppose, that its principal and exciting cause resides in the alimentary canal." And again, "The action of purgatives however, and their quick and decisive effects in subduing a fit of the gout, might lead to the inference, if other proofs were wanting, that this cause" (the cause of gout) "resided in the alimentary canal."

This pathology, although useful in the practice to which it directly leads, appears to me not only unsupported in its theory, but much too restricted in its views.

The author gives the history of a paroxysm, which he considers to have yielded entirely to the influence of active doses of opium, by which the bowels were confined, at the very time that the symptoms gave way. Is the preceding theory quite consistent with this conclusion? I am disposed to say, in comment on this case, that the gout is a constitutional disease, bringing into disorder the whole nervous system, and thereby inducing irritation, and many secondary symptoms, which require

* The substance of this Treatise has been translated with much spirit, from the French of M. M. Guilbert and Hallé, as drawn up for the *Dictionnaire des Sciences Medicales* in 1817, by James Johnson, M.D.; and very lately published, by the same ingenious author, in a separate form, with his own observations, under the title of "Practical Researches on the Nature, Cure, and Prevention of Gout," &c. &c.

† Tracts on Gout, &c. by Thomas Sutton, M.D. &c.

collateral treatment. In the paroxysm now alluded to, the anodyne treatment was followed by the use of an active purgative. It seems to me probable, that otherwise the symptoms would have returned. In every consideration which I am able to give the present question, no very limited theory, and certainly no one peculiar hypothesis, can be found applicable to explain the whole nature of gout.

Dr. Parry, in his elaborate work on *Pathology and Therapeutics*, considers that the gout is a disease depending on certain conditions of the circulating system; and offers it as one of the "exemplifications of salutary processes*." He speaks "of the final cause of the malady being merely the correction of the irregularly directed circulation†;" and further, as being a mode of "the evacuation of the habit, and the consequent reduction of a plethora, which is relatively excessive; and that another end, is the restoration of the due balance of circulation, previously determined in excess towards other and more vital parts‡."

In this view of the subject, some general circumstances of the disease appear to be correctly comprehended; but the problem still remains, What is the peculiar condition of the system, upon which the *specific* inflammation of gout depends? A preternatural afflux of blood to the affected part, is a condition of a general nature, belonging equally to any other inflammation depending on the constitution. This, indeed, might be assumed as the *proximate cause* of gout, if we considered the local characters as the whole disease, because it is the nearest antecedent to their occurrence. But such an hypothesis does not carry with it any explanation. The question

* p. 428.

† Ibid.

‡ Ibid.

under discussion is not, I conceive, more difficult than analogous questions as to many other kinds of inflammation; in relation to which we content ourselves with the general expression, that the tendency to a particular disease in an individual, arises from the peculiar predisposition of his constitution.

Although any definite and essential condition either of the solids or fluids, standing in the immediate and certain relation of cause and effect, to gout, cannot be demonstrated with the precision which may be desired, yet our information appears sufficiently complete, to conduct us to a clear arrangement in theory, and a solid foundation in practice.

I shall pursue the present investigation under the next head of my subject.

RATIO SYMPTOMATUM, OR THE THEORY OF THE SYMPTOMS;

INCLUDING THE

Chemical History of the Urinous Sediments.

OF great affinity to the last question, and nearly allied also in difficulty, is the theoretical consideration of the various symptoms of gout. It may, however, be entered upon with less danger from hypothesis; and as it involves many interesting phenomena, its discussion becomes important.

It will be useful first to bring under review, some general circumstances which are associated with the invasion of the disease*.

* Those who possess the hereditary predisposition to gout in a strong degree, require only few of the adventitious aids to-

The gouty, for the most part, have a circular form of chest; and, as we may infer, a corresponding capacity of lungs. Such is the structure most fitted for the production of the plethoric habit. In the early part of life, they usually exhibit the stamp of health. Their indulgent habits of living concur with the structure which I have described, in forming an excess of blood. In proportion, chiefly, as the habits of bodily exertion are inadequately active with relation to the plethoric state of vessels, does a state of corpulency take place. As already noticed under the head of *temperament*, they are not the truly sanguineous, but possess a temperament of a mixed character; so that the consequences of a redundant circulation rather affect the venous than the arterial system, and do not give rise to the strong actions of common inflammation. I must here again advert to the enlarged and distended veins, which are so commonly seen in gouty persons; and in the lower extremities often to a degree highly varicose. The hæmorrhoids are also a complaint in them, of frequent occurrence; and when they are attended with much occasional hæmorrhage, may be considered as one of the indications of fulness and obstruction in the circulation of the vena portarum system.

I have met with several instances of patients, who state, that previously to their being in any degree affected with gout, they were occasionally subject to epistaxis, or to hæmorrhage from the hæmorrhoidal vessels, in a much more remarkable manner than since; by means of which natural evacuation, we may consider

wards the production of the disease; and in such persons, the exceptions which occur to the general statement which I offer, will probably be almost exclusively found.

that a fulness of habit was very much relieved, and further disease prevented.

While, therefore, the *vis medicatrix naturæ* can, in these or other modes, actively counteract the bad effects of plethora; or, as long as the powers of the general circulation are adequate to the establishment of a healthy balance; bearing a just relation to the quantity of blood requiring to be circulated, and adapted to all its ultimate purposes, the gout does not make its first invasion.

I know a gentleman, now about thirty-two years of age, whose case appears to me to afford an illustration of this latter position. His father was a martyr to the gout. He has the structure and temperament which I have just described, and is a free liver. At present his exercise is active, his strength is vigorous, and gout has not befallen him; but when the period arrives that his vessels lose their tone, his indulgent habits continuing, I expect that he will become a gouty subject*.

This is a very general statement only of the question, and a closer view of it must be attempted. In a first fit of gout, the plethoric state of vessels, either absolute or relative (of which Dr. Parry has taken ample notice), appears the predominant, and often the only circumstance which can be detected in the errors of the constitution. In the returns of the disease, more or less of irregularity in the functions of the abdominal viscera becomes conspicuous; and it gradually assumes a more complicated character. In a general statement of the

* In my first edition, I made this report. The gentleman in question has recently had occasional slight pains, some sense of heat, and just a degree of throbbing in the great toe, so as a little to rouse his attention; but no visible symptoms of gout have yet taken place.

fact, it may be said, that the plethora which exists is of a partial kind. That determination of blood to the extremities, which, in its peculiar actions, exhibits the phenomena of gout, becomes more and more obviously connected with congestion in the whole system of the vena portarum; with a vitiated secretion of bile; costive bowels; and unequal function in the kidneys.

The stomach is truly the medium through which the gout is created. Excess of ingesta, beyond the powers of healthy assimilation, and the supply of blood demanded for the useful purposes of the body, is the material foundation of the disease. In those instances of sudden and unexpected attack, when the patient considers himself in the most vigorous state of his health, he is pursuing free habits of living, and carelessly producing a state of repletion, which insidiously grows into a fit of gout. The increased specific gravity of the urine depending upon an increase of its solid principles, which constantly takes place in a paroxysm, appears to me one certain evidence that the blood-vessels are surcharged with blood, unhealthy in quantity, and probably also in quality. In addition to the excess of the saline ingredients of urine, so constantly found in the paroxysm, with relation to the time of health, the fact, of which I have obtained abundant proof, that *urea* is also excreted in preternatural quantity, deserves particular attention. In several comparative examinations which I have made with reference to this point, I have invariably found that the urine secreted in the paroxysm, has furnished urea more abundantly than the healthy urine of the same individuals; and in some instances, its proportion has very much exceeded the measure of the general healthy standard. The excess of urea has, also, very remarkably corresponded with an excess of the phosphates; and

hence additional evidence is afforded that the animalising principle (if I may be allowed the expression) too much abounds in the system.

A copious appearance of the pink or lateritious sediment, which is to be taken in connexion with an increased excretion of other animal principles, is an indication that the kidneys are secreting from the blood much unassimilated matter; and according to the degree and duration of this symptom, we are enabled to form a strong conclusion as to the magnitude and importance of abdominal visceral complaint. I consider that we are to view this preternatural secretion of the kidneys at once as the *sign* of disease, and as a salutary process, which Nature is performing, to relieve an overloaded and faulty state of the circulation of the liver, and the organs associated in its functions. I view this deposition therefore as a favourable circumstance, in a case, in which I have reason to be assured that there is an unhealthy condition of the abdominal viscera; but as the general and particular evidences of improved health appear, it is also then satisfactory that the urine should lose its sediment. I have repeatedly seen, that just at the commencement, and more particularly a few days before the invasion, of active symptoms of derangement of the abdominal viscera, the urine has been light in colour, and of low specific gravity; and when, by the salutary efforts of the constitution, and the influence of proper medical treatment, the kidneys have secreted with more freedom, the urine becomes of a deeper colour, and deposits sediment. This is now and then of a whitish appearance, consisting of the phosphates with mucus; but much more commonly it is pink or lateritious, containing chiefly uric acid, in variable proportions.

In a morbid condition of the digestive organs, as-

suming the chronic form, whether in a gouty patient, or in one not liable to the disease, we find this sediment deposited only in the urine of digestion; as in that of the morning, or in that which is passed in some given time after the dinner meal. In the most aggravated cases, the urine which is secreted in the middle of the day, and more especially after a draught of any diluting fluid, will scarcely afford any of this evidence; but in the acute paroxysm of gout, and when the liver is much in fault, every portion of urine throughout the twenty-four hours, although the patient is constantly and copiously drinking diluting fluids, deposits the sediment in abundance. In every inflammatory disease, the urine which is secreted bears a specific gravity much beyond the natural standard, although the ingesta consist only of thin fluids, which scarcely possess a nutritious quality; but this circumstance, together with the excess of matter which it can only hold in solution at a high temperature, is by no means so remarkable in any state of inflammatory action of the heart and arteries, as when the liver and organs of assimilation are immediately concerned in the disease; and of which, we have the purest specimen, in a severe paroxysm of gout occurring in a patient, whose liver is obstructed and under some degree of chronic inflammation; whose state of vessels is plethoric; and whose temperament approaches to the sanguineous.

Mr. Cruickshank, in speaking of the pathological sign of the pink-coloured sediment, remarks, "this peculiar sediment we consider as in some measure characteristic of diseased or rather schirrous liver." I must here observe, that, although this sediment is a certain indication of some error of the digestive functions, it does not necessarily denote so serious a state of complaint, as is here mentioned. In many instances I have seen it ap-

pear for a short time only, as an attendant upon transient derangement. In proportion, however, as its duration is continued, we must be led to believe in a faulty state of the functions of the liver; and if the sediment continue to take place for months, the inference, that some alteration of structure in this organ may have followed such continued derangement of action as is indicated, would present itself.

Amongst the indications of a redundant circulation, existing with a relative debility of vessels, I may state the occasional occurrence of apoplexy in gouty persons, as life advances; provided that the habits of living are improper; and that the regular action of the bowels is not carefully maintained.

How much the œconomy of the circulating system is concerned in the immediate production of a fit of the gout, is sufficiently exemplified in the following phenomena. Previously to the attack, sometimes for several days, the lower extremities are constantly chilled; and the symptoms of inflammation do not, under these circumstances, take place, until the patient has been some time in bed, when the circulation becomes more equalised. This is at least one explanation why the paroxysm usually makes its invasion about midnight. Also, I apprehend that from twelve to three in the morning the last stage of digestion takes place most actively; the chyle is then freely entering into the circulation; and the natural plethora of the vessels, if I may so describe it, is produced.

When the first symptoms of the paroxysm do take place in the day, it happens as a consequence of some sudden excitement of the heart and arteries, from excess in stimulating liquor; as, after a convivial entertainment.

We are now brought to the general conclusion, that

gout is a disease depending upon a redundancy of blood with relation to the powers of the circulation, particularly affecting the system of the vena portarum, and the consequent functions of the liver; together with the production of a morbid change in the secreted products of the alimentary canal in general, and of the kidneys in particular.

In proportion as the constitutional susceptibility to gout is stronger or weaker, whether founded in hereditary predisposition, or derived alone from habits of excess in living; or increased by the influence of the disease long established in the constitution; so it is obvious, that a fit will be accordingly excited at a more early period of life, and will be reproduced by a greater variety of remote causes, and a slighter degree of their application.

Enough has already been said to explain the common existence of dyspepsia in gouty persons, the stomach being the organ first over-excited in relation to its healthy powers, in the original institution, and in the subsequent recurrences of the disease. The liver not only suffers certain injury to its functions from the same cause; but it is also a fact, that a considerable proportion of gouty persons have the evidences of a bilious diathesis of constitution, both before and subsequently to the gout, as is proved in their being very subject to jaundice, and other chronic irregular actions of the liver; and occasionally, also, to cholera morbus. The pure or mixed nervous temperament of the gouty, induces frequent and severe hypochondriasis, in connexion with the influence of an impaired state of the digestive functions.

When the gout has been frequent in its attacks, the sensibility of the nervous system is much increased, as is strongly exemplified by the various premonitory symptoms which mark the approach of a fit, and the

numerous sympathies which attend its continuance. The prevalence of cramp in gouty persons, which occurs before and during the paroxysm in the most exquisite degree, is a strong indication of high nervous irritation. The agitation of the mind is much more remarkable than in any other inflammatory disease. The disposition to return at certain intervals (a character by no means so strongly marked in any of the other phlegmasiæ), although mostly to be referred to a periodical state of plethora, either general or partial, also shews how much the disease is connected with the nervous system, in which, the laws of habit are centered.

The influence of local causes in exciting a paroxysm is worthy of observation. A severe strain or contusion in a gouty person, sometimes produces only common inflammation, as I have mentioned some examples to shew; but at other periods, is immediately followed by a fit. Mr. Hunter remarks, "It is probable that the gout is not always an act of the constitution; but that parts may be so susceptible, or rather disposed for this action, that they may immediately run into it when deranged." The truth of this opinion appears to me in some degree questionable. The gouty inflammation is an external evidence of a morbid condition of the system. When from local injury in a gouty person, *common* inflammation alone is produced, I should infer the absence of the constitutional gouty diathesis, and vice versa; so that I would rather affirm that gout is always, in a greater or less degree, an act of the constitution. This also appears to me, both the safer, as well as the more just, conclusion to be adopted.

It seems beyond our scrutiny to point out the essential condition of the constitution, which respectively produces the several kinds of specific inflammation. Peculiar

liar texture may be viewed as the entire source of the distinctive phenomena in the effects which arise from common inflammation, as we see in the example of a mucous, and a serous membrane. But this consideration does not illustrate the difference of the two diseases, *gout* and *rheumatism*; as each disease is known to attack the same textures, with very different symptoms and effects. The argument, indeed, might be much extended in application to other kinds of specific inflammation; in which, we see that each, although affecting the same textures in common, has its own peculiar laws of action.

It is unquestionably in the constitutional circumstances, that the grounds of our different pathology of these two diseases are to be found.

This part of my subject will again come under consideration, in the sequel.

The uric acid constituting the chief ingredient in the composition of gouty calculi, has led to that universal pathology of the disease with many theorists, which I have already noticed. From the same source, also, the hypothesis has probably been derived, that the local action of gout is dependant upon the mechanical obstruction of the minute vessels; which was the favourite hypothesis of Boerhaave.

With regard to the curious phenomenon of the uric concretions, which occurs in occasional examples of gout, it is necessary in this place to advert to the opinion which some entertain, that this local secretion of the capillary vessels is a constant effect of the gouty inflammation in all cases of the disease; and it is assumed that the *concretion* does not follow, necessarily, because the secreted matter may be removed by the excretion of perspiration, and by the action of the absorbent vessels. This appears to me an extreme argument, and forced far beyond the

bounds of fair reasoning. It could not happen in this manner, as the fact really is, that so large a proportion of gouty persons, as I have mentioned, should pass through life under the martyrdom of the disease, remaining wholly free from all evidence of these concretions.

It is also to be observed, that, in some of the few individuals in whom this peculiarity occurs, it is found that repeated years of gout have passed away, before any trace of the concretions was found to exist. This morbid secretion does not necessarily require for its production, an active inflammatory action of vessels, or, in other words, an acute state of gout. It sometimes takes place under the chronic form of the disease.

We continually meet with proofs of the capricious action of the secreting vessels; and of such anomaly, the gout, in the manner now stated, occasionally affords a striking example. We have no proof of the existence of uric acid in the system, independant of secretion; or even if it were present, there is no apparent cause why it should not always be secreted by the kidneys, the glands obviously designed to separate saline matter; or, why it should be thrown on the secreting vessels of the joints, or other distant parts.

I am disposed to conceive, that, in these cases, the capillary vessels of the part affected with gout may act *vicariously*, in a greater or less degree, to the secreting vessels of the kidney. In six examples of the existence of *chalk stones* both in the hands and feet, I have found, by repeated experiments, a deficiency, and sometimes almost total absence of uric acid in the urine; in reference to the usual mode of obtaining it, by means of nitric or muriatic acid. I must add, however, that even in patients of this class, during the paroxysm, this principle has existed in the urine in considerable quantity; and in

some cases, to the extent of a copious deposit of pink sediment.

In taking notice of the *local* phenomena of gout, in its ordinary forms of occurrence, I have first to advert to the almost constant termination of the inflammation *without suppuration**. Only four cases have occurred under my immediate care. In each instance, the result was curiously modified by an attendant secretion of urate of soda†. Additional cases of this description have been represented to me; but certainly they are of very rare occurrence.

Ligament is probably the texture which is the most frequent seat of gout; but the bursæ mucosæ, the sheaths of tendons, and the muscular aponeurosis, together with the respective vessels and nerves of these parts, may also be enumerated as textures *primarily* affected. *Secondarily*, the cellular membrane and skin share in the effects of the inflammation. The textures just now mentioned, belonging to the functions of the joints, do not appear susceptible of the suppurative inflammation. In the cases which I shall relate of gouty abscess, the purulent secretion was formed wholly in the common integuments.

* The abscesses which are occasionally produced by the influence of the concretions, acting as extraneous bodies, must not be confounded with this description.

† I may here take occasion to notice, that, in every example of gout concretion which I have examined, uric acid and soda have proved to be the constituent ingredients. The nature of these concretions was first pointed out by Dr. Wollaston, who shewed, with his usual clearness of demonstration, that they are composed of uric acid and soda (see Phil. Trans. 1797, p. 387). Berzelius since speaks of them, as *super-urates* of soda.

Nor do we find that coagulable lymph is produced from the inflammation of gout. The thickened state of parts which is caused by repeated attacks, arises from a change of structure in the ligaments, the bursæ, and the tendinous sheaths; and also from the morbid secretion of the two last textures.

Mr. Hunter observes, "The inflammation of gout is very different from the adhesive and suppurative in its sensation;" and further, "local inflammations depending on the constitution seldom or ever suppurate."

This last observation is, I conceive, rather exceptionable; and the explanation in question may apparently, with more correctness, be referred to the particular textures which are affected.

In gout, the inflammatory action of the vessels receives some abatement from the effusion, which in certain parts takes place abundantly, into the cellular membrane; and, at the same time, the circulation is in a considerable degree restrained by the stricture of the vessels, which the great distension of cellular membrane and skin produces. Also some changes of sensation occur, in proportion as the swelling appears, and as the inflammation becomes more superficial. The severe sense of tightness, screwing, and boring, abate; the burning and throbbing, prickings and shootings, continue; and the cumbrous sense of weight and numbness, with entire disability, increase. It may be stated, that the most painful situation of the disease is in the deep-seated affection of the ligaments, and before the swelling and redness nearer to the surface take place.

In the course of the acute symptoms, the nerves of the affected part acquire the highest state of tenderness and irritation. In no inflammation is the sense of throbbing so remarkable, as in gout. The pulsatory action

of the minute arteries, which appears to be in part owing to the distension of the contiguous parts operating as a ligature, is increased to the feelings of the patient, by the preternatural sensibility of the nerves.

In many examinations by the thermometer, I have found that gouty inflammation, when attended with severe pain, produces a much stronger sensation of heat in the affected part, in relation to the real quantity evolved from the inflamed surface, than is caused by common or rheumatic inflammation.

To exemplify this fact, I shall offer the following statements.

EXPERIMENT I.

W. W. *acute gout* in the left foot. The temperature at the external ankle, which he feels as the hottest part, and describes to be "in a strong burning heat," is 97° ; two inches above, where he feels comfortable warmth, $94\cdot5$; at the external ankle of the other foot, which is beginning to be affected with the gout, but is not, he says, "half so hot as the other," 96° . Temperature in the axilla, 99° .

EXPERIMENT II.

J. P. *acute gout* in one great toe, with a sense of heat as if in boiling water; the temperature at the hottest part is 84° ; at the same part in the other toe, which is free from complaint, and feeling comfortably warm, 83° .

EXPERIMENT III.

W. C. *acute gout* (see CASE II.); April 15th. Temperature at the middle joint of the fore-finger near the abscess, which gives to the patient a sense of heat, as if a hot iron were in contact with the part, accompanied with throbbing, pricking, and cutting, 94° ; same situation in the fore-finger of the other hand, which is free from complaint, $70\cdot5$; palm of gouty hand, $93\cdot5$; palm of sound hand, $81\cdot7$; the last joint of the affected fore-finger, 94° ; axilla in each arm $96\cdot5$; under the tongue, 99° . He remarks that each part of the fore-finger at 94° feels very much hotter than the palm of the same hand at $93\cdot5$.

Ditto, April 17th. Temperature of the last joint of the affected fore-finger now feeling warm, but not very hot, and almost free from the painful sensations before mentioned, $87\cdot5$; the same part of the other hand, which he describes as comfortably warm, $75\cdot5$; of the middle finger of the gouty hand, which is beginning to be affected, is not yet red, but is very painful, and gives him as much sense of heat as water nearly boiling would occasion, $75\cdot5$; of the same situation in the middle finger of the other hand, which is healthy, and to him feeling pleasantly warm, $69\cdot5$; palm of gouty hand, which has a much stronger sense of heat than the other palm, but is not expressed by him as painfully hot, $87\cdot5$; the other palm, $75\cdot5$.

EXPERIMENT IV.

W. H. *chronic* gouty inflammation of the back of the hand; much swollen, and accompanied with a sensation

as if a lump of lead were upon it, and affected with occasional numbness and prickings. The temperature in the middle of the inflamed part, $96\cdot5$; of the same part in the sound hand, 84° . To himself, the sound hand seems the warmest; but he does not complain of disagreeable temperature from either.

EXPERIMENT V.

J. B. has had chronic gout in one hand for two or three weeks past, with much increase of pain and swelling within these two days. The back of the hand the part affected. It pits much on pressure. The colour of the skin is dark blue. He complains of considerable heat and throbbing, and suffers severely from the sense of weight and tightness. He describes the fingers as constantly cold, and usually almost without feeling.

Temperature between the fingers of the affected hand, 63° ; of the back of the hand, where he feels most heat, 86° ; over the radial artery of this hand, where he has no unusual sense of heat, 91° ; at the back of the other hand, which is perfectly free from inflammation, 74° ; and between the fingers, $68\cdot1$.

EXPERIMENT VI.

J. D. *chronic gout*. Slight inflammation of the right wrist, with some preternatural sense of heat*, much the-

* It is to be considered, that in all these examples of gout, the examinations were made in the day time; a period of the

cal distension, so that the tendons seem matted together; the parts feeling tight, but scarcely painful, except on motion. Temperature at a spot near the radial artery, which he describes as the warmest part, 95° ; at the same part in the other hand, which is free from complaint, 93° ; and over the radial artery of this hand, 94° . His left foot is much swollen, and pits on pressure; towards the little toe the skin is vividly red, not very painful, and not feeling more than very warm. The veins of this foot much fuller than in the other limb, and particularly so near the inflamed part. Temperature at the warmest spot, 96° ; at the same part in the other foot, which is free from complaint, 93° ; just beneath the ensiform cartilage, 95.5 .

In every instance in which I have examined by the thermometer the state of parts under gouty action, however slightly existing, I have found the temperature more or less increased. This is illustrated by the three following experiments.

EXPERIMENT VII.

T. W. the last stage of *acute gout* in the hand and foot. A little remaining redness of a dark hue; much pitting on pressure; but this is borne almost without uneasiness. Temperature between the thumb and finger, where most appearance of the complaint remains, but no particular sense of heat is felt, 93.5 . At the back of the hand, 92.5 . Between the thumb and finger of the other

twenty-four hours, at which, the several painful sensations abate more remarkably in this disease, than in the other kinds of inflammation.

hand, which is free from complaint, 92·5. Between the great and second toe, where very slight gout remains without sensible heat, 89·5. At the ham of the same limb, over the popliteal artery, 88°.

EXPERIMENT VIII.

L. P. convalescent from *acute gout* in the foot. Expresses a slight degree of preternatural warmth at the ball of the great toe, where the temperature is 84°. Same part in the other foot, which is not affected, 83°.

EXPERIMENT IX.

J. S. the last stage of *acute gout* in the side of one foot, slight aching only remaining; the most uneasy part scarcely warmer to his sensations than the sound parts, 86·5. Same situation in the other foot, 83·5.

EXPERIMENT X.

W. C. *rheumatic inflammation* in one hand, chiefly at the wrist. He expresses that he feels "fever heat, with some throbbing, gnawing, and pulling," and appears in very severe pain. Temperature at the ulnar side of the wrist, where he has most sense of heat, 92°; at the same part in the other hand, which is free from complaint, and comfortably warm, 85°.

EXPERIMENT XI.

W. C. *rheumatic* inflammation of one hand, and the whole of the wrist. She complains of pricking and shooting, and intense pain, and expresses the heat as that of hot water. Temperature at the most inflamed part, 97° ; at the same part of the sound hand, the skin of which she feels as temperate, 93° .

EXPERIMENT XII.

M. H. has a whitloe at the middle finger; much surrounding inflammation; vivid redness of skin; the sense of heat as if in water just warm. Complains of severe pain, of much tightness, and some throbbing. Temperature at the most inflamed part, 83° ; at the same part in the other hand, which has just been exposed to cold, uncovered, 66° . Palm of the affected hand, 88° ; palm of the other hand, 69° .

EXPERIMENT XIII.

H. A. ulcer on the back of the hand, produced by a recent burn. The integuments are swollen, and pit on pressure; the skin vividly red; she complains of throbbing, pricking, and shooting, and describes the sense of heat as from scalding water. Temperature at the hottest part, 94° ; of the same part in the other hand, which she feels pleasantly warm, 93° .

It appears from these experiments, that the sensation of heat (or of pain like that produced by a great degree

of heat) which is experienced by the patient, even in gouty inflammation, corresponds rather to the degree of pain which is present, than to the thermometrical* temperature of the affected parts; but it seems to me clearly proved, that, other circumstances being equal or nearly so, the gouty inflammation is productive of more intensity of sensation, both with respect to the heat and other sufferings, than in the contrasted examples of inflammation which I have offered. Haller considered the chief seat of the diseased local action in gout, as being in the nerves themselves. Without contending for the truth of this extreme position, which I consider too much an assumption, it certainly appears that the nerves are affected in a very remarkable degree.

In the case of inflammation from a burn, the subject of Experiment xiii, it is shewn, that, as in some of the examples of gout, the sensation of heat was much more excessive than might be expected to have been present from the indications of the thermometer. In this kind of injury, we know that the nerves, in common with other parts, suffer violent lesion, and high consequent irritation.

The intense degree in which the gouty inflammation affects the nerves, is further exemplified by the following curious fact.

J. S. having the gout in both feet very severely, when much swelling had not yet taken place, and suffering such pain and sense of heat at the time, that he could imagine his feet in a furnace, or that some one was

* In these experiments, I employed a delicate animal-heat thermometer, covering its bulb with a thick cork hollowed out for the purpose, with a notch to receive the stem. As in neither experiment, the temperature of the apartment was more than 64°, I have not noticed the exact degree.

actually forcing into them red hot spikes, accidentally dropped on the inflamed part, a piece of tripe immediately from the boiling kettle. The skin became instantly vesicated, but he was not sensible of any increase of heat in the part.

R. D. having gout at its height in one foot, with much swelling, accidentally received a considerable quantity of *boiling* water on the part. The previous pain and sense of heat were only slightly increased; and, in this case, the cellular membrane being already extremely distended with serous effusion, no vesication ensued.

To the account which I have offered of the peculiar force of sensation which prevails in gout, I may add, in recapitulation, the subsultus tendinum, tremor of muscles, severe cramps, and irritable state of the mind, which usually occur in a violent paroxysm; demonstrating altogether the acute sensibility of the brain and whole nervous system, which is connected with this disease*.

Another circumstance which shews a prevalence of nervous irritation, is a fact with which some of my patients have made me acquainted, that in regard to the predisposing cause last alluded to at page 76, they have often foretold an attack of gout, by finding a more than usual state of excitement, without a correspondent share of strength.

I must not pass over entirely unnoticed, the circumstance of the desquamation of the cuticle from a part affected with gout, as the inflammation entirely ceases. Dr. Cullen notices it as among the characters of regular gout. In ninety-eight patients, with whom I have ex-

* Pliny, in Epist. xii, mentions the commission of suicide by his friend Corellius Rufus, on account of his severe sufferings from the gout; for which, his physician had not been able to afford him relief!

amined this point, I find that 25 have never experienced this symptom; not more than six of the seventy-three invariably; and many of them, in some fits, and not in others: and in no instance, I believe, except from the hands and feet. This exfoliation of the cuticle appears to be principally owing to the acrimony of the cutaneous secretion; but may probably, in part, be referred to the distension which the integuments suffer, obstructing the free circulation of the extreme vessels. Those who have almost always noticed the desquamation to result after a long fit, state, that if the symptoms have been quickly removed by medical treatment, this circumstance has not occurred.

Of the Chemical Composition of the Urinary Sediments.—A knowledge of the composition of the different sediments of the urine is so materially connected with the theory and practice of medicine, that I shall present, in the most compendious manner that I am able, some details on the subject; first taking notice of a refinement of opinion which has been advanced, respecting the peculiar acid existing in the pink sediment.

Proust*, many years ago, endeavoured to prove that the pink or brick sediment, to which he refers, *as the substance which, during fevers, separates itself from the urine at the moment of its cooling*, consists in part of a distinct acid, different from the uric, and which, from its colour, he termed the *rosacic*. In order to determine the propriety of this conclusion, I have, in repeated experiments, examined favourable specimens, both of pink sediment, and of the red crystals (gravel); and of this comparative investigation, as relating to the present question, I shall briefly offer the results.

* Annales de Chimie, vol. xxxvi. p. 265.

The crystals appear to be very little acted upon by boiling water, except sustaining some loss of colour, from the separation of animal matter ; but if previously triturated to a fine powder, they acquire a considerable degree of solubility.

The pink sediment, collected on the filter and dried, is found to be an impalpable powder. It dissolves, with the exception of one tenth of its proportion, in boiling water ; but partly separates into the solid state on cooling. Nitro-muriate of gold added to either of these solutions (of the crystals, or of the pink sediment) while yet warm, instantly produces a purple colour, and a precipitate subsides ; but the effect takes place slowly, if applied to the liquor when cold.

Each substance is readily soluble in pure potash ; and, during the solution, apparently yields to the muriatic acid stopper, the white fumes characteristic of ammonia.

The potash solutions, by the addition of muriatic acid, precipitate small grains of a greyish white colour, which, when washed, and treated with nitric acid in the usual manner by evaporation, and carefully heated, afford the rose hue* distinctive of uric acid.

* The rose hue which is obtained either from the crystals or the pink sediment, is heightened to a beautiful carmine by the addition of ammonia ; and is rendered permanent in the evaporating dish for several weeks, if carefully dried and kept free from moisture. Without ammonia it more readily deliquesces, and assumes a yellowish green colour ; but the former hue may be revived by heat ; and, in a dry apartment, will continue a few days, more or less perfect. With or without ammonia, if transferred to paper, and preserved from light and moisture, the first tint is permanent. Lastly, the tint with ammonia, moisture being entirely excluded, remained unimpaired after daily exposure to light for about a month, when it began to fade.

The crystals, separately treated with nitric acid (more favourably if diluted), afford the rose hue.

Precisely the same result is obtained from the pink sediment.

Muriatic acid added to the watery solution of the pink sediment, previously filtered, produces a precipitation of minute whitish grains; which, washed and treated with nitric acid, afford the rose hue.

Nitric acid added to the watery solution, evaporated and heated, presents the rose hue.

From these experiments it appears evident, that the uric acid is a constituent of each substance, and that the theory of a distinct* acid is not demonstrated.

The crystalline form of the one sediment, and the divided impalpable nature of the other, in addition to its easy solubility in hot water, are circumstances of difference which are necessary to be considered.

Mr. Cruickshank observes of the lateritious sediment, "We have examined several portions of this sediment, and have generally found it to be composed of lithic acid, phosphate of lime, and some peculiar animal matter but little soluble in water. It has by some been sup-

* Subsequently to these experiments, I have read the paper on the rosacic acid, in the *Annales de Chimie*, No. 287, by M. Vogel, who concludes with noticing the similarity of the rosacic acid to the uric, and the easy transition of the one into the other, by means of the action of the acids. This fact alone, however, of its being changed indiscriminately by any acid, appears to me to invalidate his opinion, that the uric acid, and the rosacic so called, are distinct substances. All the facts which are stated, both here and by M. Vogel, seem to demonstrate, that the phenomena do not depend upon mere conversion of the pink sediment into uric acid; but upon a separation of the uric acid, from some principle with which it was combined.

posed to consist entirely of lithic acid, but this substance for the most part constitutes by far the smallest part*.”

I have derived the following results from my examination of different specimens of the pink and brick-coloured sediment. Neither of the varieties effervesces with muriatic acid, nor affects vegetable colours. Heated before the blow-pipe or in the crucible, each kindles, blackens, emits pungent fumes, among which, the odour of prussic acid is perceptible, and consumes to a greyish ash. That afforded by the pink sediment forms about a tenth of the original matter. The ash of each sediment changes turmeric paper to brown, renders violet paper green, and dissolves with effervescence in muriatic acid, it having been converted into a carbonate, from the decomposition of the uric acid by heat. Ammonia added to this solution, causes a white granular light precipitate. Oxalate of ammonia produces a white heavy precipitate. The ash, on being exposed to the atmosphere, does not deliquesce†. This ash is also considerably soluble in boiling water; and, on the addition of ammonia, affords by its precipitates, the evidence both of phosphate of magnesia and of lime. Hence it seems to follow, that

* Rollo on Diabetes, 2nd Edition, p. 449.

† In formerly believing that the ash was deliquescent, I conceive that, from some accidental circumstance, I must have fallen into an error. In the several specimens of pink and lateritious sediment which I have lately examined, I have not found the ash left from long burning, to be in the least degree deliquescent; and it has invariably reddened turmeric, proving therefore that the alkali was *soda*. My notice was directed to the particular examination of this last fact, by learning from my friend Dr. Prout, that Dr. Bostock had found urate of soda in the pink, and in the lateritious sediment.

the pink sediment is composed chiefly of uric acid in combination, and animal matter, on which the colour may be considered to depend ; together with a small proportion of the usual phosphates of the urine. It appears evident that the uric acid exists in combination with soda, for which it has a strong affinity. I have constantly found, that the sediment which is of the deepest pink colour, affords the least residue on being burnt, and contains the least proportion of the phosphates ; and that in proportion as the shade of colour is lighter, the ash which remains is more abundant, and consequently the phosphates also. We may, therefore, form a good presumptive conclusion of the nature of the sediment in the urine of a patient, from the colour alone. I have not found any difference of composition in sediments of similar external characters, whether procured from the urine of a person under gout, or any other disease connected with a faulty state of the digestive functions.

On examining the *crystals*, I obtained the same results as the above, except that the ashes which they left after combustion, were less than one-twentieth of their original weight, and consequently so minute, that when dissolved in an acid, they scarcely yielded a sensible precipitate on the addition of ammonia in excess. The two substances therefore appear to be very similar in their composition ; and the probable reason why the uric acid does not crystallize in the pink sediment, is, that the animal matter or mucus discharged with it, so mixes with it, as to prevent it from assuming a regular form ; for I have uniformly observed, that such urine as deposits the pink or lateritious sediment, is much loaded with a mucus-like substance ; and that, on the contrary, such urine as deposits the crystals, is almost free from this mucus.

I lately had the opportunity of examining a small calculus, which was passed by the urethra from a gouty person. For a considerable time before its expulsion, it had occasioned irritation of the bladder, and, for two days previously, excessive pain in the whole course of the cord on one side, and difficulty in the passing of urine. Since its discharge, he has been free from such inconvenience; but the urine has most commonly deposited the lateritious sediment. The calculus was of the size and form of a full lemon kernel. Its colour brownish red. On being divided, the disposition of its component parts appeared in concentric layers; the outer one yellowish, the inner brown; and within, a blackish substance seemed like a nucleus.

The following were the results of my analysis. Almost wholly soluble in potash ley. Treated with nitric acid, it gave the rose hue. One grain boiled in distilled water left $\cdot 4$ of a flaky yellowish residue; and this, on being burnt, afforded a white ash of inappreciable weight, which reddened turmeric; did not effervesce with an acid; and proved to be pure lime. The filtered water on cooling deposited a profusion of minute pellucid needle-like crystals; which, with nitric acid, afforded the rose hue. A portion of the water, concentrated with its solutions, on the addition of muriatic acid, precipitated minute grains, which also, with nitric acid, gave the rose hue. Pure ammonia added to some of the watery liquor produced a flaky precipitate, which, on trial, proved to be magnesia; and, with the filtered fluid, oxalate of ammonia instantly yielded the evidence of lime. Before the blow-pipe, the brownish red portion of the calculus blackened and gave pungent fumes. The blackish nucleus-looking portion kindled with a faint flame. The ash affected turmeric, and effervesced strongly with

mmuriatic acid. In this case, the uric acid, which, in the boiling process, was dissolved in the water, had been decomposed. The ash also afforded evidence of phosphate of magnesia, and phosphate of lime. Hence, with the exception of the presence of oxalate of lime, this calculus furnished results very similar to those of the lateritious sediment.

I now come to speak of the nature of the whitish or yellowish white sediment, and white shining crystals, which, as to their external character, I have before described.

When urine, as it is first passed from the bladder, appears turbid, it may immediately be pronounced to contain the phosphate of lime, and the ammoniaco-magnesia phosphate, both suspended and in solution; and on the cooling of the urine, a part of the crystallized precipitate forms with some mucus into an iridescent pellicle on the surface, and also subsides to the bottom, being the mixed sediment before mentioned. It is invariably accompanied by an abundance of mucus, which may be considered to be partly derived from the inner coats of the kidneys, but principally from the surface of the bladder; the arteries of these parts being excited to increased secretion, for the purpose of defence from such preternatural stimulus.

This urine, like that which precipitates the pink or lateritious sediment, is always of high specific gravity. It has been stated by authors, and it is the prevalent theory, that such urine is certainly alkaline; and the conclusion has led to a rule of practice, which I shall have occasion to notice. I have applied the test of litmus paper, in at least thirty instances, to the most favourable specimens of urine of this description, and have invariably found it reddened in the usual manner. It is, however,

highly animalised; very soon runs into the putrefactive process; and often contains such a remarkable excess of urea, that, even without concentration, from the application of nitric acid in the usual proportion, crystals of urea very soon appear.

In order to the examination of the urine depositing the mixed sediment in question, the following steps are to be pursued.

Filter the urine, and ascertain its specific gravity and other general characters. Collect the sediment from the filter, and adding diluted nitric acid to a small portion of it in the platina dish, evaporate it quickly over the lamp. Should the least portion of uric acid be present, the rose hue will appear, just at the moment when the evaporation is about to cease. From these trials, I have derived the knowledge that in every instance in which the sediment is entirely unaccompanied by red, reddish white, or pink, or pinkish white colouring matter, the uric acid is absent. The phosphate of lime thus treated, exhibits a greenish white appearance, which quickly passes into yellowish white.

We are now to presume, that the sediment consists of the phosphates; and this may be proved by digesting it in dilute nitric or muriatic acid. If the whole be dissolved without effervescence, and be again precipitated by pure ammonia, it evidently consists of the phosphates. The triple phosphate of magnesia appears in minute grains, and being of light specific gravity, chiefly floats on the surface, and adheres to the sides of the vessel; while the phosphate of lime in small flakes immediately subsides to the bottom. I have invariably found the surface pellicle, which forms in urine depositing the sediment now under consideration, to be composed of the phosphates; and almost wholly of the triple phosphate.

Whether the white or yellowish-white sediment of the urine consist of rough powder alone, or of shining crystals, or of both in mixture, and involved with mucus, the following process, as more demonstrative, may be employed. I am aware that it is liable to some objections, where great accuracy is required; but from repeated trials which I have had occasion to make, I can recommend it as affording a proof of the existence, and an useful presumptive evidence of the proportions of the two salts.

Subject the mass to a red heat until the animal matter is destroyed, in a platina or earthen crucible. Supposing that it consist of the ammoniaco-magnesian phosphate, and the phosphate of lime, the ammonia having been expelled, the ash will be composed of phosphate of magnesia and phosphate of lime. This ash is found not to effervesce with muriatic acid; nor to affect vegetable colours. It is now to be analysed. It may be put into diluted acetic acid, which quickly dissolves the phosphate of magnesia, only slightly affecting the phosphate of lime unless heat be applied. What remains, therefore, undissolved, after a few minutes digestion, may be considered as phosphate of lime.—To the phosphate of magnesia now in solution, add pure ammonia in excess; which, neutralizing the acid, combines with the phosphate, and unites to form the insoluble ammoniaco-magnesian phosphate; and this salt, first appearing on the surface, slowly subsides in small shining crystals. The remaining part of the ash is to be treated for phosphate of lime, with muriatic acid and pure ammonia, as just now described. The urinary sediment in question kindles and blackens before the blow-pipe without flame; emits an ammoniacal odour; and partly fuses, and forms into very thin enamel plates.

In the analysis of a compound calculus, which not long since I had occasion to make, I found the following formula successful for the separation of the phosphates of lime and magnesia; but certainly the process is more troublesome than the occasion would, in general, make desirable. Clear presumptive conclusions are, in general, sufficient for our purpose, as to relative proportions.

^a Boil the mixed salt with three or four times its weight of carbonate of potash in solution; filter the fluid, and wash the insoluble matter by repeated affusions of distilled water.

^b Dissolve the residue which was left in the last process (carbonates of lime and magnesia) in diluted nitric, or muriatic acid, and decompose the solution by adding neutral carbonate of ammonia in excess, whence is produced a precipitation of the carbonate of lime.

^c To the fluid from which the carbonate of lime has been separated, add a concentrated solution of phosphate of soda. The precipitate which now falls down, and which may be collected upon a filter, is the triple ammoniaco-magnesian phosphate; from the quantity of which, as well as of the carbonate of lime, the relative proportions of the two salts may be deduced.

^d To determine the proportion of phosphoric acid (if desired), let the fluid which was left in the first process (^a), be neutralized by the careful addition of nitric acid, so as only just to saturate the excess of alkali; and to the neutral solution add a solution of nitrate of lead, until a further addition ceases to produce a precipitate. The phosphate of lead thus obtained, whenedulcorated upon a filter, dried and weighed, will serve to indicate the quantity of phosphoric acid contained in the salts.

The dark sediment, possessing a blackish green colour, and appearing chiefly crystallized, alone remains

for description. It kindles before the blow-pipe, blackens, and apparently, from the peculiar animal matter which it contains, burns with a faint flame. To the ash obtained from sufficient burning in the crucible, add diluted acetic acid, as in the former case. The kindling before the blow-pipe has already led to the conclusion that the sediment contained oxalate of lime; and this will be confirmed on finding that the ash (if extreme heat have not been used) affords a strong effervescence on the addition of muriatic acid. The crystallized appearance of this sediment, together with the peculiar colour of its accompanying animal matter, distinguish it to the eye from the pink and lateritious sediment, which also furnishes, on burning, a carbonated salt. For the analysis of this ash, acetic acid is to be first employed, which proves a ready solvent for lime or its carbonate, as well as the magnesian phosphate. The acetic solution thus obtained, may be decomposed by oxalate of potash, which will separate the lime; and the subsequent addition of carbonate of ammonia will throw down the magnesian phosphate. The remaining steps of the operation for the phosphate of lime, may be such as have been already described.

The ashes of the different sediments which I have examined, have invariably left some particles of a matter insoluble in muriatic acid, even with the aid of heat; and which appeared to be silex.

DIAGNOSIS OF GOUT,
OR
THE DISCRIMINATION OF ITS PRESENCE
BY ITS SIGNS.

From Rheumatism, the gout is to be distinguished not so much by any one mark, as by the concurrence of several circumstances. In the first fit of gout, it seldom happens that more than one part is affected; and still more rarely, that more than one part is attacked at the same moment. This solitary disposition of gout, as to its situation in the first attack, is a striking point of discrimination; to which may be particularly added, the adult age. In ordinary examples, the diagnosis is not difficult. The absence of distant premonitory symptoms, though occasional in gout, is almost constant in rheumatism. The remissions from pain and fever during the day, are much more distinct in acute gout than in acute rheumatism; and among the local characters which are striking in gout, the following may be selected:—A serous effusion into the cellular membrane, which takes place almost immediately after the invasion of the fit in the parts already enumerated (p. 30), and which yields a pit to the finger as the inflammation ceases, and in a slight degree even before this, when the skin is not extremely tense;—a turgescient state of the neighbouring veins, more marked and general than in rheumatism, sometimes appearing in the whole limb, and occasionally preceding the inflammation, a day or two, or even longer; the severe intensity of pain accompanied with extreme sense of heat and violent throbbing; the acute sensibility of the parts to touch, and particularly to any concussion; the excessive sense of weight, with numbness

and total disability. A gentleman relates to me, that, on one occasion, just as he was convalescent from gout, after partial exposure to damp, he was seized with rheumatism in both arms; the pain was agonizing, peculiarly gnawing, and seemed to him to be in the very substance of the bone. He described it as quite different from his usual pains of gout. I have witnessed, in the last year, several instances exactly similar, as to the contrasted sensations produced by gout and rheumatism; the latter existing in one part, and the former in another, at the same time; or, the rheumatism occurring as a sequence. The symptoms which I have recited, if not entirely distinctive, are strongly diagnostic. Sometimes the inflamed surface has a shining appearance as if varnished; and, occasionally, desquamation takes place, a circumstance which I do not remember to have seen in rheumatism, unless as the consequence of particular applications. Our diagnosis is further assisted by collateral considerations, such as those derived from the habits of the patient, the constitution of the parents, and the nature of those remote causes which may have led to the disease. For example, a young man, aged thirty, above the middle height, and slight; of the nervous temperament; had suffered two attacks, at distant intervals, of rheumatic inflammation of the ankle joints, with flying pains in different parts of the body. The local appearances were, a slight vivid blush of redness, and a distension of the bursal texture, but no œdema of the cellular membrane. The sensations, severe gnawing pain as if in the bone, with occasional pricking and shooting; but scarcely any heat. About two months after a recovery from one of these attacks, and during which period he had been living indulgently in all respects (his circumstances in life being improved), and had become much more corpulent, the gout,

for the first time, took place, affecting the great toe and instep of one foot. I was much interested by the clear comparison which my patient himself drew between the rheumatism and the gout. "Now, the part was swollen to a great size; the skin shone as if varnished, and was uniformly of a deep red at first, and afterwards became rather purple; finally, the skin peeled off from the whole surface." In the rheumatism, he had not found any particular intermission of pain in the twenty-four hours; but in the gout, his extreme sufferings were from twelve to three in the morning, and about five he procured sleep. His sensations under gout were, "of heat, as if scalding water were poured on the part; throbbing, as if receiving blows; pricking and shooting much more intense than from his former rheumatism; and above all, a deep-seated sensation in the joint, as if the parts were undergoing dissection."

His father was subject to gout. His only sister has suffered severely from chronic rheumatism.

From Erysipelas.—The species, *phlegmonides*, can alone baffle the discrimination of the practitioner; and this only happens when it affects parts also liable to the gout. The inflammation in this complaint is more superficial than in gout, affecting primarily different textures, and is usually more disposed to spread. It is ushered in by stronger constitutional symptoms, as drowsiness, headache, and severe rigors, if the attack be urgent; if it be slight, the distinction may be deduced from the specific local characters. In erysipelas, the heat of the affected part resembles rather the stinging of nettles than the strong burnings of the gout; nor is there in erysipelas, the same disabled condition of the moving powers, as in gout.

From phlegmon, or common inflammation, the diagnosis can never be rendered difficult, except in a first case of gout; and even then, an attentive consideration of all the relative circumstances, constitutional and local, will remove every doubt.

PROGNOSIS.

IN a first fit of gout especially, the favourable progress and termination of the symptoms are indicated almost solely by local appearances; and the event of first attacks is often so favourable, that the patient congratulates himself on the benefit which he has received, or at least expects to receive. In the severe and complicated returns of the disease, it affords a very different prospect. Mr. Cruickshank remarks*, “In gout, the termination of the paroxysms is most perfectly indicated by a copious lateritious sediment; and when this suddenly disappears, and the urine at the same time affords a precipitate with the muriate of mercury, a fresh attack or relapse may be expected.”

If this statement were correct, the practitioner would be furnished with a very simple guide in forming a judgment of the case; but I have invariably found, that, except with very dilute urine, the muriate of mercury always produces a precipitate, even in health. With respect to the lateritious or pink sediment, it must be again observed, that its appearance and quantity are connected with the state of the alimentary canal, and of the circulation, and with the specific gravity of the urine itself. It begins and goes on with the fit, in a disordered state

* Rollo on Diabetes, p. 449.

of the digestive organs, connected with some degree of inflammatory excitement in the system; and does not, as is sometimes stated, denote the close of the paroxysm, but the contrary. If, however, having disappeared, it again return, it certainly indicates, that the internal functions are not restored to health, and that, consequently, a relapse may be expected.

On the subject of prognosis, I am induced to mention, among the favourable signs, the visceral organs being sound in structure, and not materially disturbed in their functions; an entire cessation of the sympathetic fever; the tongue becoming moist and clean; a return of natural appetite; the *fæces* recovering and maintaining a healthy character; the urine *ceasing* to deposit sediment, and at the same time losing its high specific gravity; the nervous system becoming tranquil, and the spirits improved; the local sensations readily yielding in their severity to remedies, and the inflammation soon abating*, and not shewing a disposition to quick transference from one part to another; or, if it be fugitive, not fixing severely in

* Dr. Hamilton (Letters on Gout) who assumed an opinion that the disease "depends on a fluid matter *sui generis*," speaks of a sweat issuing from the pores of the skin at the inflamed part, as the first sign of the resolution of the inflammation. He adds, "This discharge from the pores is accompanied with a very peculiar *fœtor*, which is the first and strongest proof, that is obvious to the senses, of the existence of the genuine gouty fluid." He pursues this idea of gouty effluvia at great length, and affirms, that it is "like no other existing in nature." Except in two or three instances, I have not been able to find that there is any truth in this asserted odour of a gouty part; and I can only consider the occurrence as the effect of a cutaneous secretion peculiar to the idiosyncrasy of some persons.

new situations.—The skin which has been affected, becoming pale and permanently cool, affords a favourable omen; and with regard to a desquamation of the cuticle, when it does occur, it may be considered as a strong indication that the inflammation is wholly removed from that particular part.

In proportion as the opposite circumstances are observed to take place, we have, according to my experience, the assurance of a tedious and difficult form of disease; and among the unfavourable signs, I consider the strongest to be, a quick transference of severe inflammation from one part to another, joined with painful sympathy of the stomach or the head; with irregular fever; and with exquisite sensibility of the whole nervous system. I must also add, that, in every case, in which, the internal secretions fluctuate between the condition of health and disease, we should give a guarded prognosis as to the probable period of true recovery. Until I had acquired considerable experience (that powerful corrector of the judgment) in the management of a severe gouty paroxysm, my expectations of an uninterrupted convalescence were sometimes sadly disappointed by a troublesome relapse. The chief grounds on which the prognosis is to be formed, are, an attentive observation and accurate knowledge of the state of the respective secretions. When they are become *permanently* healthy; when all local complaint, except some weakness, has disappeared; and, when in conjunction with such change, the tranquillity of the nervous system is restored; we may look with confidence to the stability of the cure.

TREATMENT OF GOUT GENERALLY CONSIDERED.

THE prohibition of Sydenham against the least interference in the paroxysm, long exerted a powerful influence on medical practice; an influence which continues to this day to have considerable force. It is obvious that he derived all his opinions from the doctrines of the humoral pathology, which at that time pervaded the whole theory of medicine. His practice, as he himself declares, was still more rigidly cautious in the gout, "than in most other humoral diseases." In this disease, he says, "Nature seems to have the prerogative to expel the peccant matter according to its own method, and throw it off from the joints, there to be carried off by insensible perspiration." In his *Treatise on Gout*, he equally deprecated "bleeding, purging, and sweating, the only three ways proposed of expelling the morbid matter;" but in a subsequent treatise, "*Of Bloody Urine, from a Stone in the Kidnies*," he expresses the following qualification: "With respect to purging, therefore, in the case of bloody urine, provided only manna be used according to the method above delivered, I must retract an assertion I formerly published in my *Treatise on Gout*; namely, that it is absolutely improper to purge gouty persons, either at the beginning, declension, or in the intervals of the fits: for I did not recollect then, that the fit, which I feared might be occasioned by the purgative, might be prevented by giving an opiate at night." He still adds, however, to this passage, "If the gout only be attended to, all manner of evacuations are very pernicious therein, and therefore not to be used, unless the above-mentioned symptoms require them."

Much, then, as the moral sentiments, and a portion, also, of the medical reasoning of this amiable Physician command our admiration, it would be unworthy of the great improvements with which time has since enriched our art, if we should continue fettered by the peculiar doctrines which I have quoted, and remain content to yield the cure of gout to *the nurse* alone. I freely admit what I have already stated, as a general position, that Nature is seeking a remedy for herself in a fit of the gout; and, also, that if her purpose were always accomplished with as much success as in a *slight* first fit, the physician might with propriety stand by, as a looker on merely, admiring the *vis medicatrix Naturæ*. But, in truth, the disease not only returns frequently, but returns with constant increase of strength; and the remedial efforts of Nature, which were at first so satisfactory, become tedious, irregular, and uncertain. The gouty sufferer, gradually reduced to lameness, constant pain, and many secondary evils, suffers at length the fate described by Lucian, thus expressed in the Latin translation:

“ Multæ formæ infortunatorum,
 “ Meditatio pœnæ, et consuetudo,
 “ Podagros miseros consolentur.”

Sydenham himself, in the feeling description which he drew of “the racking pains, unfitness of motion, and other disorders which afflicted him during the greatest part of his life,” was a striking example of the imperfection of Nature considered as the physician, and of the erroneousness of his own doctrines of practice. Several instances have come before me of patients, who, being prejudiced against every kind of interference with the gout, have left the paroxysm to its own course. A few,

of strong constitution, and being accustomed to have the disease mildly, do not appear to have suffered any serious inconvenience from such proceeding; one has now and then appeared, who has praised the gout; but the majority have found ample cause to regret their neglect of regular treatment, either in the injury sustained in the general powers of the constitution; or in the textures which have been affected with inflammation; or in both ways.

Lately I saw a gentleman, who, in his second fit being attacked in both feet, and one hand, hailed his visitor with satisfaction, and cherished his pains with flannel; resolved to be the patient of *Nature* alone. Six months had already elapsed when he consulted me; and even then, his tongue was furred; the secretions from the alimentary canal and kidneys were in a very unhealthy condition: he was lame in the feet, from the weakness of the ligaments, and the distended state of the bursæ, and of the sheaths of the tendons; and his hand was almost useless from weakness, and the contracted state of his fingers. All this state of evil gradually yielded to a suitable plan of general and local treatment; and this gentleman fully admitted the propriety and benefit of medical assistance in the gout, as much as in any other disease.

Under the artificial condition of man in social refinement, Nature is not always able (and seldom indeed permitted) to employ the wisest, shortest, or safest methods of cure. I trust that we are paying her sufficient respect, in attentively seeking to discover the kind of evil which she has to remedy; in taking all her indications for our guide, and acting truly as her servants; but not with *fettered hands*.

I would assume it therefore as a principle, that we

should attempt the prevention of a fit of gout, if warned of its approach; and interrupt its progress when formed, *unless* such a state of the constitution exist, that the gout has taken the place of another more serious disease; or may be expected to prevent one which is threatening, and more to be dreaded than itself; but even in this case, it is incumbent upon us to moderate the violence of symptoms; to study and fulfil particular indications; and carefully to estimate the balance of the present evil, with the expected good.

In the view which I have taken of this part of my subject, I shall first consider,

THE TREATMENT OF THE PREMONITORY SYMPTOMS.

THESE may sometimes be remedied with the effect of averting the threatened attack; and even when from the state of the system, in concurrence with general causes, this preventive object cannot be accomplished, the paroxysm will be rendered milder, and more tractable, by the previous removal of evident causes of irritation. Much may be effected by timely anticipation of that relief to the constitution, which Nature is about to attempt.

If the true inflammatory diathesis be present in any considerable degree, general bleeding should be employed; or, if congestion be indicated in the vessels of the head, or liver, or other internal organ, without the action of the pulse being increased scarcely beyond the usual standard, local cupping will be more advantageous. Any tendency to hæmorrhoidal discharge which is manifested, should be promoted. With this view, aloetic and saline purgatives may, with great advantage, be administered in con-

junction. The costiveness, which is so usual a fore-runner of gout, should be removed by an active purgative; and, with this intention, calomel, antimonial powder, and the extract of colocynth, at bed time, followed by sulphate of magnesia, and infusion of senna with some aromatic tincture, in the morning, produce very good effects. Or, in place of these medicines, the more particular treatment, which I shall speak of under the head of Prophylaxis, may be adopted. If a furred state of tongue, with *heart-burn**, nausea alternating with a craving appetite, and acid eructations, be present, an emetic of ipecacuanha should be administered. Should the threatening symptoms continue troublesome, we are to consider that the constitution is labouring under the causes of the paroxysm, almost as much as if the full symptoms had actually arrived; and a great part of the same treatment which I shall presently have to describe for the paroxysm itself, must be adopted. In numerous cases, I have, in the most satisfactory manner, entirely averted an attack, which, according to all former experience, would certainly have taken place, severely. If, after the employment of suitable evacuations, the internal secretions continue to assume a vitiated appearance, as indicated by the state of the fæces and urine, already described, small unirritating doses of mercury, administered at a proper interval, according to the principles of Mr. Abernethy, and a gentle aperient once a day, with or without the addition of a stomachic bitter, according to circumstances,

* A medical gentleman, who has had severe fits of the gout, informs me, that he has often taken warning from these symptoms; and considers, that, by prompt appropriate remedies, and subsequent regimen, he has completely succeeded in averting a threatened fit, on several occasions.

will constitute an efficacious mode of treatment. Alkaline medicine has its advantages, but produces only palliative relief, unless the bowels are also excited. Particular indications are to be met by corresponding remedies; but to the use of medicine should be joined careful moderation in diet*, or, on some occasions, a course of rigid abstinence; horse and foot exercise in the country air; repose of mind from anxious employment; early rest, early rising, and early meals; in one word, all the *lædientia* should be changed for the *juvantia*.

I have now to offer the arrangement which I have attempted.

OF THE TREATMENT OF THE PAROXYSM.

IN our choice of remedies for the particular symptoms which appear in every individual case, we should reflect upon the kind and degree of the predisposing and exciting causes, by which the fit has been introduced; and our practice should be relative to such considerations, as well as to the age, constitution, and temperament of the patient.

I propose a detailed view of the several remedies which I have thought most deserving of attention, in the following order:

Of Bleeding.—The employment of the lancet in gout is not allowable with the same freedom as in the other phlegmasiæ, in most of which it is our chief dependance. It usually happens in the gout, that the increased excitement affects the nervous system much more than the

* Quæcunque parit repletio, inanitio curat.

heart and arteries; and, as I have before stated, the redundancy of blood appears to belong rather to the circulation of the vena portarum, than to that of the general system. It may also be assumed as a practical fact, that this kind of partial plethora is more favourably and effectually removed by purgatives and diuretic medicine, than by the detraction of blood itself. With respect to the loss of blood from the system, as a remedy in gout, it appears to me wholly unnecessary for the removal of the local inflammatory action, to which other treatment proves very adequate. Nor does general bleeding, however freely employed, afford that relief to the local inflammation, and the pain more particularly, which might be expected. Of this fact I have seen many strong proofs.

A gentleman, aged forty-two, of full habit, but of the nervous temperament, subject to severe and protracted fits of gout, contracted a violent cough with urgent sensations of soreness in the chest, from exposure to wet and cold. He was bled from the arm to the quantity of sixteen ounces; a blister was applied to the chest; and other usual means were adopted. In two days after, he was attacked with gout, which first affected the hands, and next, the knees and one foot, in succession. In this fit, and particularly on the day before the seizure of the knees, the thighs were very curiously affected with sudden disability, in such a degree, and in a manner so entirely unprecedented, that he was infinitely alarmed with the dread of paralysis. He continued to suffer much from this extreme weakness of the lower limbs. His general languor, and depression of spirits and of strength, were also excessive; to which were joined copious perspirations. In this case, the temperament of the patient, or probably in more correct expression, the peculiarity

of constitution, rendered general bleeding, to the extent mentioned, a very unfit remedy. A moderate abstraction of blood, from the chest, or neck, by cupping, would in all probability have proved sufficient to relieve the inflammatory irritation of the lungs, without so material an expence of the general strength.

In the present instance, I know that the inflammatory action of the affected parts was less acute than on former occasions; but it seemed questionable, whether upon motion, the pain was less in comparison. The nervous restlessness was greater; and debility and depression, as already stated, were truly distressing.

In one case, within my immediate knowledge, the patient was first attacked with inflammation of the liver, for which, from the urgency of the symptoms, seventy ounces of blood were taken from the arm in the space of thirty-six hours. In a few days, violent gout succeeded, and apparently not controuled in the severity of painful symptoms by the previous depletion. I could relate many more instances in illustration of my present subject; and certainly the point in question is one of great practical importance. There are good grounds, therefore, for the objection to the general principle of bleeding in gout; but it has been carried much too far, and a prejudice of very ancient date has been established against taking away blood *generally* in the gout, under any circumstances.

When the inflammatory diathesis is strongly marked by a full hard pulse, hot skin, scanty and high-coloured urine, with costive bowels, general bleeding will certainly be indicated, as it would be were no gout present. And when any internal organ is affected with inflammation, the observations which I have just now offered of the

unfitness of this remedy for the gout itself, alone considered, must not be allowed to have weight.—The greater indication must be obeyed. The quantity to be abstracted, and the repetition of the venesection, must be adapted to the urgency of the general inflammatory action; to the state of the affected organ; and to the effect produced. I may also add, that it should be in relation to the *powers* of the individual, rather than to his age, of which, if we judge medically by years alone, we take a fallacious guide.

In forming our judgment as to the propriety of this active instrument of benefit or injury, we should give a due consideration to any influence from exciting causes, of a temporary or transient nature, such as the effects of sudden excess at the table, or of the passions of the mind; and which effects may be removable by other means. But when, under a predisposition of the system to inflammation, wet and cold, or a continued course of excess in stimulating liquors, have excited the paroxysm, the inflammatory diathesis becomes more permanent. In the circumstances in which bleeding is a proper remedy, its *early* employment is a point of much importance. This remark has, indeed, a general application to the treatment of any of the phlegmasiæ. When bleeding is improperly delayed, it will be found that the depression of strength resulting from the excessive irritation of the nervous system, counteracts its advantages in a great degree; and although timely practice is important in all the remedies which are to be adopted, it is so especially with this of bleeding, in the few instances in which it is required.

The advantages of bleeding in the gout as a practice of familiar employment, have, I am persuaded, been

much too strongly insisted upon by Dr. Hamilton*, and Dr. Rush†.

Barthez, Guilbert, and Hallé, wholly condemn the practice of general bleeding in gout.

When there is tenderness to pressure in either hypochondriac region, not depending on a distended and faulty state of the upper bowels, but joined with strong indications of congestion in the hepatic circulation; or when marks of fulness in the vessels of the head appear predominant, the action of the pulse at the wrist not being increased, ample local cupping is to be preferred to the lancet.

Emetics.—From my own experience I am not induced to advise their employment, unless an evacuation of the stomach in a full degree is obviously required, from its being pointed out by indications of irritating contents. The influence which acid matter in the stomach may have in exciting or aggravating the symptoms, is such as to claim our particular attention.

I saw the good effects of an emetic exemplified in one very striking instance. A man, just seized with the gout in the hand, was in severe torture, so that he writhed and cried with anguish, and at the same time was affected with nausea, and had a furred tongue. He took a full dose of ipecacuanha; and, on discharging from his stomach much acrid acid, of a greenish appearance, together with heavy mucus, was so much relieved, that he fell immediately into a refreshing sleep; and

* Letters, &c.

† Rush's Medical Inquiry and Observations, vol. v.

although many other remedies became necessary for the removal of the paroxysm, this preliminary step was of great advantage.

In two other cases, I prescribed an emetic at the commencement of the paroxysm, with advantage in relieving the stomach from its immediate state of error; but the symptoms of gout were not apparently controuled.

Dr. Small* extolled, in his own state of gout, the use of an emetic, and gave the preference to tartarized antimony. This medicine he also gave with Peruvian bark, which, he says, in this combination acted very favourably as a mild laxative.

Cathartics and Diuretics.—On the choice and free employment of purgative and diuretic medicine, the successful treatment of the paroxysm chiefly depends. The least reflection upon the state of the system in gout, and how much it is connected with previous repletion, which exhibits its influence in various ways, and manifestly in the secretions, will necessarily lead us to this method of practice. When the eau medicinale, hellebore and laudanum, and various other specifics, have been most successful in the paroxysm, the action has been powerful on the bowels. Many authors have given their testimony to the propriety of purgatives in the gout; and Dr. Sutton, to whose Tract I have before referred, ably enforces the advantages of this treatment actively pursued†.

* Medical Observations and Inquiry, vol. vi. art. 20.

† Purgative treatment is strongly condemned by the French physicians, evidently upon the mistaken principles of the humoral

In considering the rationale of the evacuating treatment from the alimentary canal, in gout, we must refer to the pathological principles of the disease, upon which I have already expatiated; and in this view of the subject, the following intentions seem to be presented: to unload the bowels of solid accumulation, and excite vascular secretion and consequent discharge throughout the whole tract of the intestinal canal; to promote the excretion of vitiated bile, and remove also the acid matter and unhealthy mucus to which disordered digestion has given rise. In some degree we thus detract from the circulation at large, and more particularly from the vessels belonging to the system of the vena portarum. In imitation of Nature's efforts to remove redundant matter by the medium of the kidneys—an action, the existence of which I think myself entitled to infer from my experiments—we are to keep the corresponding treatment attentively in view; and I have invariably employed, with the greatest advantage, purgative and diuretic medicine conjointly; so that the exhalant vessels of the alimentary canal, and the secreting function of the kidneys, are stimulated to increased action at the same time. With this view, probably, some have selected elaterium; and, by Dr. Sutton, it has been advised in combination with opium. But when I reflect on the uncertain operation of elaterium, and its injurious violence occasionally, even in small doses, I can but consider, that unless in a sluggish state of bowels, which is also joined with ana-

pathology, and the consequent apprehension of producing a weakened action, so as to impede the course of the gouty matter to the extremities. They allow the use only of very mild aperients, and particularly praise “un électuaire lénitif sulfuré.”

sarcous swelling in the extremities, its active nature should be rather dreaded than chosen as a regular remedy in gout. Under certain indications, and within proper limitations, however, it is a medicine as useful as it is powerful.

In stating the kind of purgatives which I have found most favourable and efficacious, I shall offer, in as brief a manner as I can state, the general plan which my own experience has sanctioned, by its being attended with particular success. Occasional doses of calomel, in conjunction with antimonial powder*, compound extract of colocynth, and a little soap, fulfil, in the most useful manner, the first part of the intention which I have expressed; and they should be repeated each night, or each other night, according to the degree of vitiation which the bilious and other matters from the bowels appear to possess; and according to the advantage derived.

For the purpose of co-operating with the medicines just described, so as to affect the bowels and kidneys at the same time, without nausea to the stomach, and with the least uneasiness to the bowels, I have experienced the most remarkable success from a draught composed

* So far as my experience has gone, I am induced to give the preference to that preparation of antimonial powder called *James's Powder*. I have considered, that, in equal doses, it has nauseated less, and has yet acted with more certainty on the bowels, than the ordinary powder of the Pharmacopœia. As it comes from one individual source, it follows as a probability, that it should be a more uniform preparation, than what we meet with prepared in the usual way; and this conclusion will seem the more reasonable, when we reflect that antimony is a volatile substance, and being moreover the most active ingredient, the quality of the powder must be materially affected by the variable degree of heat, to which it is liable to be exposed.

of Magnesiae gr. xv. ad xx.*; Magnes. Sulphat. ʒj. ad ʒii.; Aceti Colchici ʒj. ad ʒii; with any distilled water the most agreeable, and sweetened with any pleasant syrup, or with 15 or 20 grains of Extract. Glycyrrhiz. When much feverish heat of skin has prevailed, I have been induced to add to this draught, gr. xxvii of Potass. Carbon†. accurately neutralized with fresh lemon juice; and, in this case, have preferred the carbonate of magnesia to the calcined; using a larger proportion‡. It should be repeated at intervals of four, six, or eight

* It has been suggested to me, that the combination of the alkaline earth with the acetic solvent of the colchicum appears *unchemical* and contradictory; and I think it necessary therefore to enter into some explanation of my views in regard to this formula. I find by experience, that magnesia and the sulphate, in conjunction, act very happily as a certain and easy purgative; and that the colchicum, brought to the state of a mere solution in water by having its acid menstruum neutralized, is in the most favourable state of preparation in which it can possibly be administered. A very small portion of magnesia proves sufficient to neutralize the acetic acid (becoming an acetate), and the remaining portion is left to act as an ant-acid in the stomach and bowels, and unite with the sulphate in the advantageous effects just described. When the patient is furnished with this medicine in quantity, for occasional use, I recommend that the magnesia be added to the dose at the time only of its being taken; as otherwise it adheres with much inconvenience to the bottom of the vial.

† The carbonate of potash is a medicine much to be preferred to the subcarbonate, which is a very impure salt; not unfrequently containing metallic impregnations injurious to health, derived from unfit vessels occasionally used in the process of preparation. I have detected both copper and lead.

‡ The carbonate loses exactly two-thirds of its weight, by the process of perfect calcination.

hours, according to the freedom of its operation and the urgency of the symptoms; and I must dwell upon the importance of adapting the activity of this part of the treatment entirely to the degree of gouty inflammation which is present. However reluctant we may feel to lower the powers of the patient by general bleeding, we must not be restrained by any unfounded fears of debility, which may be entertained as to the present mode of reduction. Such apprehensions must be satisfied by those sound arguments of good practice, which the practitioner alone can demonstrate. The inconvenience of the operation will be abundantly compensated by the important advantages produced; and if any treatment of the paroxysm is entitled to be considered *radical*, I am sure it is one which aims at the removal of those obstructions and vitiated actions of the visceral organs, which are the great supporters of the disease. In the advice even of particular remedies, it is obvious that I can only lay down general principles of treatment; but I may say, that the purgative and diuretic medicines in question should be actively administered, until the gouty inflammation subsides; and so long as the urine, which is first passed in the morning, retains a high specific gravity; or, as a rule of more easy application, so long as it deposits sediment. In proportion as improvement in these points is obtained, the frequency of repetition of the medicine should be lessened, to thrice, twice, and once in the twenty-four hours: but it should not be discontinued until all inflammation is removed; both the *fæces*, and urine* acquire healthy characters; and the

* It is not sufficient to authorize a discontinuance of the medicine that the urine becomes healthy; or that the state of the

tongue becomes clean and moist. The fluctuation of improvement and relapse which a severe paroxysm of gout so often produces, must be watched with the best attention; and the use of effective practice should not carelessly be conceded to the first flattering marks of convalescence.

I thought formerly, that, in two or three instances of particular idiosyncrasy, the colchicum, although thus mildly exhibited, was too acrid in its operation to be continued, from its appearing to cause some distressing heat and irritation in the bowels. I believe this to have been the fact in one instance; but having watched the progress of subsequent cases, in which the same suspicion had arisen in my mind, I have been led to the conclusion, that this symptom has arisen almost wholly from the morbid state of the secretions. Indeed this point has been rendered generally quite clear, because the irritation has ceased upon an improvement in the state of the bowels being acquired, although the same medicine has been regularly continued. The exception, therefore, which must occur to all general rules, scarcely ever applies, according to my experience, to the fitness of administering this medicine in the paroxysm, or when gouty action is in any degree present, agreeably to the rules which I shall offer, of combination with other general and particular treatment. I may affirm that this preparation of colchicum, in union with the other medicines, has never disappointed me in its effect, either to contribute to the production of watery

bowels is merely for the most part correct. Even when every point of convalescence seems well founded, this or other alterative medicine, should be given at certain more distant intervals than before, rather than at once be laid aside. Upon this principle of treatment, I shall have further occasion to expatiate.

evacuation from the bowels, or to increase the discharge of urine abundantly, or both. It also assists the excretion of bile; and its operation is remarkably easy and agreeable. I shall again have occasion to notice the grounds on which I prefer the acetie preparation of colchicum; but I may now further mention, that it does not tend to produce the least nausea or weakness of stomach; nor is the draught in question disagreeable to the palate.

Very rare occasions only, in the acute disease, will call for the preference of the more stomacheic* purgatives; which are probably often prescribed upon the theory, that an atony of stomach necessarily belongs to gout, as a primary character.

Mercurial Preparations.—The administration of mercury during the paroxysm, and more especially in the interval, appears to have been recommended by some authors from a theory of its *specific* operation; and by others upon more general principles. When employed occasionally as a mild alterative, at bed-time, in conjunction with the use of a gentle aperient in the day; or, joined in a full dose with purgative medicine, this active agent has full claims to our regard. With the former intention, calomel, or the quicksilver pill, in a small dose with James's powder, or the compound calomel pill, produce excellent effects; but if any of these preparations be given in frequent doses, so as to excite mercurial fever, more or less of serious injury follows as a certain consequence, without any corresponding advantages. I

* The London *gout cordial* is, I believe, composed of rhubarb, senna, liquorice extract, and aromatics, digested in proof spirit; and is very similar to the *gout cordial* of *Boerhaave*.

shall mention three strong instances which I have witnessed of this general fact; but slighter forms of the evil to which I refer, such as increased fur of the tongue, and nausea, attendant upon an irritated stomach, and in union with much general nervousness, have frequently come under my observation.

In one gentleman, who in three days took a quantity not exceeding six grains of calomel, in divided portions, at bed-time, in conjunction with small doses of opium, while the bowels were daily acted upon by purgative medicine, a very severe salivation took place, attended with high fever and irritation. The gout, which had almost quitted him, and had declined in the most favourable manner, was re-excited, and became more painful, intractable, and tedious, than almost any case which I remember to have seen.

In another example, mercurial inunction had been incautiously used by a gouty patient for a slight syphilitic complaint. The strong effects of mercury were produced, and a violent paroxysm of gout immediately ensued, which proved of unusual severity and duration. In a third case, the mercurial action, together with an attack of gout which seemed entirely *consequent*, produced such continued irritation and debility as proved fatal; the patient, however, I must add, having previously laboured under a diseased state of lungs. I have been induced to suspect, from these and additional facts, that gouty persons, in general, are more easily affected by mercury than others. Whether or not this opinion be correct, I am persuaded that they require more than ordinary caution in the employment of mercurial medicines.

Upon this conviction, it is my invariable practice in using mercury as an alterative with a gouty patient, to give it in a single defined dose at bed-time, each night,

or each other night, according to the circumstances of the case; never to employ a large dose, except as an immediate purgative; and very cautiously to avoid the excitement of mercurial fever and irritation.

Pretended Specifics. — Of the unbounded list of boasted remedies for the gout, with which every age, from the earliest history of medicine, has supplied credulity, I shall notice under the present head, a very few only, and those of modern date and having some existing repute. Wine of white hellebore with laudanum (which has been offered as *identical* with the *eau medicinale**) has gained considerable estimation with some practitioners, either as an active purgative, or as possessing *specific* properties. From some examples which I have seen of its operation, I am convinced that it is too dangerous a medicine to be familiarly employed, and also that it can seldom be given in free doses to the gouty patient, with any prudence. Even in careful and gradual doses, its accumulated action sometimes becomes alarming, and causes serious hyper-catharsis. In its severer effects, under a bolder administration, I have known it to produce, in quick succession, vomiting and purging, great depression of strength, spasms, and *burning* pain in the bowels, cold sweats, and, in some instances, tremors and faintings, and the very sensations of approaching death. I am informed, on good authority, of a case of gout, in which a saturated preparation of hellebore in alcohol proved fatal by its action on the stomach and bowels; and of another case in which the patient died apoplectic, after a free use of the tincture of

* See Mr. Moore's Letter to Dr. Jones, on the Composition of the Eau Medicinale.

hellebore and laudanum. This termination was attributed, and with great apparent probability, to the stimulating properties of the medicine.

A gentleman informs me that in his last fit of gout, which had attacked him severely in the feet, he took the tincture of hellebore to the extent of three drachms, in divided doses, in the space of twenty-four hours, joined with laudanum. It caused profuse perspiration; and the operation on the stomach and bowels was most alarming. The griping was so severe, and the depression of the animal powers so excessive, that, he declares, he wished for death.

Dr. Woodville mentions, that, "upon opening those who have died by the effects of this poison, the stomach discovered marks of inflammation, with corrosions of its interior coat; and the lungs have been much inflamed, and their vessels much distended with dark blood*."

I have been lately consulted by a gentleman, who, on a former occasion, having gout severely in the knee, took Mr. Moore's preparation of hellebore and laudanum, in correct doses, several nights in succession. With him, it does not appear that the medicine materially disagreed; although he describes himself to have had unpleasant cold perspirations, and some depression of strength, which I consider should be referred to the medicine; but the most material point respecting it, in this case, is, that it did not seem to have the least controul over the symptoms.

Even when the tincture of hellebore acts with comparative mildness, it has occasioned, under my observation, in two cases of gout, considerable irritation of the stomach, causing a distressing sense of heat, a white

* Medical Botany, p. 276.

tongue, with thirst, and, together with inflammatory excitement, much nervous depression. Finally, of this medicine, I am well persuaded that, in any form or combination, it should be entirely deprecated as a remedy for the gout.

The *gratiola*, or hedge hyssop, has been much extolled, and has, by some, been considered as the chief ingredient in the French medicine. I have employed a saturated tincture of the plant well prepared, without any satisfactory result; and indeed in the two cases in which I gave it free trial, it proved rather an inert medicine. An eminent practitioner informs me, that he has seen effects from this tincture very analagous to those of the *eau medicinale*, when that medicine acts mildly as a sedative.

The *colchicum autumnale*, or meadow saffron, has more particularly been asserted to be the essence of the *eau medicinale*. This is to be considered as the *hermodactyl* of the ancients, which formed an ingredient for many centuries past, in the most reputed formulæ for the gout*. It was almost discontinued in medicine, and particularly as a remedy in gout, when Mr. Want, I believe,

* In Bernard's edition, Leyden, 1743, of a very old Greek M.S. upon the Gout, written by Demetrius Pepagomenus at the desire of the Emperor Michael Palæologus, who reigned anno 1282, and which appears to have been first rendered into Latin by Marcus Musurus, at Rome, in 1517, I find the following prescription:

“ *Compositio simplicium Pilularum purgantium.*—Aloes pars una; *hermodactyli* dimidium; *glycasini* aut *cinnamomi* quod *hermodactylus* stomacho sit infestus, dimidium; *scammoniae pulcherrimæ* sextans. Ex iis fiant pilulæ; denturque pro viri-

first in any public report*, revived the attention of the Profession to its peculiar influence in this disease. From the many trials which I had made of the different preparations of colchicum in gout, I was convinced that this author had made too extravagant statements of the curative power of this medicine. In the first edition of this Treatise, I gave the following account.

I have repeatedly also made trial both of the powder and the tincture, but from neither have I been able to trace the smallest *specific* operation†. It was taken in free doses, but with effects altogether unsatisfactory, when used in either of these forms, and relied upon as the only medicine. I observed that the stomach was irritated; an increased fur of the tongue, with thirst, were produced; and no certain action of the bowels occurred.

I am again led, by my present subject, to notice the papers of Sir Everard Home on Colchicum, published

bus, materiæ copia, et natura ejus medicamenti quod datur, et tempore."

In several other prescriptions in this little Treatise, the hermodactyl was the principle ingredient.

Quincy, in his Dispensatory, 11th edition, 1789, observes, of the colchicum, "that it stands so much in the esteem of some writers, as to be dignified with the name of *Anima Articulorum*, the soul of the joints." "Because they believed it to be very efficacious in scouring the mucilaginous glands, and preserving them free from the lodgment of such gritty matter, as occasions the gout and arthritic complaints." Quincy makes reference to many officinal preparations containing colchicum, and thence styled *arthritic*.

* See Essays in the Med. and Phys. Journ. No. 185, &c.

† By this expression, I meant, so as to produce a certain and permanent relief.

in the Philosophical Transactions*, offering a strong recommendation of this medicine, and of the eau medicinale in common with it, as a *specific cure* for the gout. A recommendation so enthusiastic and general in its principle, and from such high authority, is, in my opinion, much to be lamented; it being calculated to prolong the popularity of that injurious medicine, the eau medicinale; and also to promote the exercise of empirical principles of treatment. Sir Everard observes: "For the cure of gout, the eau medicinale of Husson has been most fortunately discovered to be a specific remedy, and it is now ascertained, by experiments on different people, that a vinous infusion† of the colchicum autumnale or meadow saffron, is equally so, and therefore the two medicines must be considered the same." The most substantial answer which I can give to the first of these opinions, will be found in the details, which I shall presently offer, of the effects of the eau medicinale on those gouty persons who have come under my care, and from whom I have collected their own unbiassed reports. That the two medicines are to be considered "as the same," is surely a very unsupported opinion, when it is made to rest only on the general assertion of a similarity of power in removing the immediate symptoms of gout. A general claim of this kind (in the desire to represent the eau medicinale) has been maintained with regard to many other medicines; as the wine of hellebore, and wine of landanum; elaterium, guarded by opium; and, according to the

* Part II. 1816. Part II. 1817.

† "Made by macerating two pounds of the fresh roots in twenty-four ounces of Sherry wine, in a gentle heat for six days, the spirit being previously carried off by heat."

opinion of some, even the *gratiola* or *hedge hyssop* : and two quack medicines, called *Wilson's tincture*, and *Reynolds' specific*, may be added to the list. That all these medicines, except the *gratiola*, have usually, in different degrees, an immediate controul over the local symptoms of gout, I am well assured ; and sufficient observation has taught me, that the tincture of *colchicum*, and the vinous infusion of the same substance, have also a similar influence ; but, so far as I have seen, in a much less degree than the *eau medicinale* ; and less than *Wilson's tincture*.

Dr. Sutton, when speaking of the *eau medicinale*, states*, "When I became acquainted with the publication which announced its efficacy, I proposed to my medical friends a substitute for this secret medicine, and which should produce similar effects : this was *elaterium*, in doses of one or two grains, with forty or sixty drops of tincture of opium. In one case, I soon had an opportunity of finding this remedy to be completely effectual ; and Mr. Green, of Lewisham, administered it in smaller and repeated doses, in two instances, with the best effect ; in one of them with such rapid success, that the patient thought he had taken the French medicine." It is well known that Mr. Moore's preparation with *hellebore* † for a long time was considered to be the *eau medicinale* ; and, doubtless, this belief was, in a great measure, founded on a comparison of the effects of the two medicines. I have traced a considerable difference in the usual operation on the system, which all the various medicines I have mentioned, have respectively produced ;

* Tracts on Gout, &c. p. 201.

† Composed of three parts of the wine of white hellebore, and one of the wine of opium.

although they have had so much, in common, the effect of procuring a *palliative* cure. The construction which should be fairly given to the term *specific*, deserves some discussion. Sir Everard Home, in his zeal to commend the eau medicinale, and the vinous infusion of colchicum, describes its power over gout, as corresponding in point of efficacy with that which mercury possesses over the syphilitic virus, with no other difference than that of being more rapid in its operation. Dr. Parr, in his Medical Dictionary, thus expresses his article, "*specifica* (from *speciem faciens*, peculiarly adapting) specifics. By specifics, are meant such medicines, as infallibly and on all patients, produce given salutary effects; acting by some unknown power on the disease, without being directed by indications. Such was the bark supposed to be in intermittents, and mercury in lues: such are the boasted remedies of the quacks. In general, however, as sound science has increased, these vaunted specifics have disappeared, and we now find no remedy which does not, by some action on the system, point out the principles by which its salutary effects are produced." The very general success of mercury in destroying the syphilitic virus in the system, gives it a real claim to the title of specific. The Peruvian bark deserves almost the same general praise for its speedy and permanent controul over a regular intermittent fever. This, however, cannot be said of the eau medicinale; tincture of colchicum, and the vinous infusion; the preparation of hellebore and opium; elaterium and opium; Wilson's tincture; and Reynolds' specific. They do, in most instances, for a few trials, influence the local symptoms very speedily; but, so far from removing the cause of gout, they leave the disposition to the disease much stronger in the sys-

tem, with less powers, it is true, to produce violent inflammatory attacks; and lead to the still more calamitous, because more constant, sufferings of the chronic form of the disease. The patient emphatically describes, that his feelings make him in constant dread of something worse occurring than the gout; which his constitution no longer seems able fairly to produce. With the effects of elaterium and opium, I am the least acquainted; but I have had abundant opportunity to know that each of the other medicines, sooner or later, disappoints the patient of his expected cure, rendering merely a palliative assistance, and keeping the disease dormant for a time only, so that it is left to prey on the constitution with more lasting and serious ill effects.

After my first report of the properties of colchicum in its different modes of preparation, I again made trial of the tincture, in two cases of gout situated in the feet. It produced, in each instance, very similar effects; the fur of the tongue was increased, and there was much thirst. It acted on the bowels as an acrid purgative, causing remarkable heat and irritation, and also affected the kidneys as a diuretic, very favourably. The symptoms of gout yielded very soon, but returned after an interval of ten days; and the disappointed patients were desirous of other treatment. One very gouty person informs me, that in the summer season he took, every day for three months, six grains of genuine colchicum root dried. At first it affected the bowels and kidneys moderately, and he felt improvement in his limbs. Afterwards it had scarcely any sensible action. At the expiration of the three months, and first at the beginning of October, he was seized more violently with the gout, than upon any former occasion; and not adopting any medical treatment,

he was confined, with severe symptoms of gout, to his chamber for nineteen weeks, suffering for three weeks great torture, equally day and night.

I have also witnessed the effects of the infusion recommended by Sir Everard Home, and am persuaded, that it is a preparation much to be preferred to the spirituous tincture. It has afforded palliative relief to many persons; for a time, to their satisfaction; to a few, more permanently; but I have been consulted by some, who have found all the ill consequences to the stomach from this infusion, which I have ascribed to the tincture, without any satisfactory influence being produced over the severe symptoms of the paroxysm. Others have derived immediate relief, but the gout has recurred much more quickly than before. With some, the relapse has taken place almost immediately, and the repetition of the medicine has appeared of no satisfactory avail. Where this remedy has most succeeded, I have observed, that relapses have so much occurred, that the disease has gained in frequency what it has lost in force.

I have had abundant opportunity to be acquainted with the present and ultimate effects of Wilson's Tincture and of Reynolds' specific; but a slight notice of these nostrums will be sufficient.

One gentleman, who took it very steadily for many months, and was a warm advocate for its effects, relates to me, that the gout, which formerly observed intervals of at least three months, has, since the employment of this tincture, returned every ten or fourteen days. His previous favourite remedy was the eau medicinale, and from the similarity of operation in his case, he considers the medicines as the same.

Another very gouty subject, usually having one severe fit in the year, in the first use of this medicine,

thought himself possessed of a most valuable prize, and, in his joy, almost disregarded the gout. But see the quick sequel! His relapses were frequent, and the influence of the tincture over the symptoms became weaker. He was without appetite; he suffered sensations of weight, tightness and general oppression about the stomach and abdomen; and was highly nervous. At length, during even all the summer months, he was not free from actual gout for the space of a fortnight; and in vain did he seek relief from his former boasted antidote. I could relate almost an infinity of cases in which this tincture has, after a time, entirely disappointed the hopes of the patient, and given him more or less cause of regret, that he has depended on its palliative relief*.

According to the reports which I have collected, I find that Reynolds' specific; in small doses, does not produce certain influence over the symptoms; in large doses it often operates violently both on the stomach and bowels; but in no case within my knowledge, even when most successful, has it done more than effect a short cure; and the patient has laid it aside, either with dissatisfaction or complete disgust.

Upon the *eau medicinale*, I shall offer my observations at considerable length. Its credit appears truly to have declined so much in general estimation, that any laboured exposure of this baneful nostrum might appear unnecessary; but as it still finds some strong advocates, and as the panegyric bestowed on it by Sir Everard Home†, must tend to encourage an inclination for this

* For additional authority upon the injurious consequences of using Wilson's Tincture, &c. I refer the reader to some Observations lately published by Dr. Williams.

† I will not apologize for the freedom which I use in criticising the statements of this distinguished author; being well

remedy, in the minds of those persons who are impatient of the rules of regular treatment, I shall endeavour to discuss its real pretensions.

In the first trial of the medicine, it proves in most instances a powerful palliative or short cure; removing the paroxysm as by a charm, and not unfrequently without any very sensible operation upon the stomach, or upon any of the excreting organs. This curative power of the remedy gradually lessens on repetition, and with many persons becomes entirely lost. Such disappointment occurs very commonly, even where its first agency produced the most flattering expectations. In one example of peculiar idiosyncrasy, which came under my observation, the patient reported that he had in a severe paroxysm taken six bottles (procured from the dépôt) within a few weeks, and yet from this free use of the medicine no sensible influence, either as to operation or relief, was produced. On the other hand, the capriciousness of its action is now and then manifested by very alarming consequences, affecting the stomach and bowels with all the virulence of an active poison. Such opposite effects from the medicine may perhaps, in part, be attributed to accidental difference in its strength*, but chiefly, I presume, to the modifying influence of constitution, in different individuals.

assured, that, from his zeal in science, he will court the contest of opinion, where any important truth is to be established.

* In an old bottle of the medicine which I examined, the sediment, which adhered tenaciously to the bottle, was considerable.

To the objection derived from the uncertain immediate agency of this medicine, although certainly it is one of great importance, we might reconcile ourselves with less difficulty, than we can to that of its insidiously leading to a train of subsequent evils; the former being only an inconvenience of a similar kind to that which attends the use of every active substance derived from the vegetable kingdom. When the eau medicinale does not immediately debilitate by the violence of its effects, it often leaves behind an impaired condition of the nervous system; so that the head is affected with frequent giddiness; the stomach with weakened digestive power, and frequent sensations of sinking and vacuity; the limbs, and especially the parts affected in the paroxysm, suffer for many weeks with tremblings, numbness, and coldness, and very commonly with tedious œdema; these symptoms appearing variously in different individuals. The mind is affected with exceeding depression; and some persons declare their general feelings to have been rendered miserable for many months, even from having made a moderate use only of the medicine. It tends also to render the bowels inactive, to diminish the alimentary secretions, and materially to weaken the functions of the liver. In the general character of the medicine, it may with truth be stated, that sooner or later, in proportion as it is freely employed, it leads to a broken state of health. Of its deleterious powers in a more immediate manner, I may offer the following examples.

In one case, the contents of a single bottle taken as a dose, so paralysed the stomach, that for many days it was scarcely sensible to the strongest stimulants. The patient was recovered with much difficulty, and remained for a long time in a state of serious debility.

Mr. Ring* relates an instance of its causing the death of a gonty patient; by the direct violence of its operation.

Dr. Gregory, the distinguished Professor of Medicine in the University of Edinburgh, has received full conviction, from his own observation, of the injurious tendencies of the eau medicinale in gout. One of his patients almost fell a victim to the cholera which it produced. Another was soon affected with symptoms of hydrothorax. In the next attack of gout, this patient rashly repeated the medicine, and hydrothorax again followed. His health was as before fortunately restored by good treatment; and his lessons of future prudence were purchased at a sufficient price in the dangers he so narrowly escaped.

A few years ago, I was favoured with the following communication from Dr. Parry: "From the eau medicinale, I have seen in several cases great injury to the alimentary canal. In the majority, it has eased present sufferings at the commencement; has not diminished the actual quantities of inflammation; and, in process of time, has disappointed the patient as to its present effects."

I have now to offer some further, and more particular details.

CASE I.

J. S. aged 55, robust and plethoric; of nervo-sanguineous † temperament; gout first at 29, not hereditary.

* Treatise on Gout, p. 175.

† Sanguineous the chief character, but mixed with the nervous; and *vice versa*, when sanguineo-nervous is expressed.

In one paroxysm, just on its decline, took fifty drops; its effects delightful, and operating mildly on the bowels. Repeated it on several subsequent occasions, and believes that he has taken a dozen bottles. Gout returned every month or two since this use of the medicine, not so painful as before, but very harassing; and, contrary to former experience, the stomach became much weakened, with severe symptoms of dyspepsia; and the bowels were with difficulty regulated even by medicine. In every attack since, also has noticed abundance of the pink sediment in the urine.

CASE II.

L. M. aged 59; formerly robust; corpulent in the abdomen only, and constitution much worn; of sanguineo-nervous temperament; gout first at 25, not hereditary. Has taken the medicine many years past. It has always quickly removed the painful symptoms; but it has been only palliative, as the fits which previously returned twice, or at the utmost three times in the year, have since taken place every four or five weeks. Last week took three bottles; half of one each night; the stomach much nauseated every day, and on the morning following the last dose, he vomited so severely, that arterial blood came from the stomach; and this has happened before. Has recently complained of an unusual weakness in all his joints, and of an œdematous state of ankles towards night, which has much alarmed him. He looks very sallow and unhealthy, and is altogether highly nervous; much more so than formerly, but asserts that he does not consider his stomach weakened, or his digestion impaired.

CASE III.

D. T. aged 46, robust and plethoric; of nervo-sanguineous temperament; gout hereditary, and first fit at 30; took the medicine in one attack, in full doses, with much less effect on the immediate symptoms than commonly happens, and considers that it produced such uncomfortable and even alarming feelings of the head, which continued very troublesome, that he would not be induced on any account to repeat the medicine.

CASE IV.

C. B. aged 56, formerly muscular; corpulent in abdomen only, and now very infirm in the limbs; of nervo-sanguineous temperament; gout first at 26, not hereditary. When he began the medicine, he took a bottle at twice on succeeding nights; it did not operate sensibly, nor much relieve the paroxysm. For a year and a half after, no regular fit, but often affected in a distressing degree with flying pains in all the limbs, to which he had never before been subject. Doubtful whether this chronic state of complaint had been produced by the medicine, or from cold. In the next painful attack, took half a bottle, without the smallest relief.

CASE V.

C. L. aged 56; very corpulent and plethoric; of nervo-sanguineous temperament; gout hereditary, and

first fit at 42. Having severe gout in one hand and both feet, and quite disabled, at six in the evening took three parts of a bottle; it almost immediately produced delirium, which became violent, but yielded to a profuse perspiration. It acted moderately on the bowels. At six next evening, could walk across the room even without a stick. The swelling and soreness of the parts remained a fortnight, but without much pain. For a long time unusually nervous. Fit returned in four months. Took, as the first dose, half a bottle, with much relief to the symptoms; but it first produced delirium as on the former occasion. Four months after, in summer, again attacked, when he repeated the medicine. Has not resorted to it for three years past, and his final report is, that for some time after its use, he suffered much from flying pains in the head, with other uncomfortable feelings; from dyspepsia; cramps; œdematous swelling of the legs with much weakness; and indifferent sleep. He became extremely corpulent; as already stated, the recurrences of gout were frequent; and since the discontinuance of the medicine, he adds, the fits have observed an interval of twelve or fifteen months, and his health has latterly, from regular medical treatment, much improved.

CASE VI.

T. F. aged 44; corpulent in the abdomen; of nervous temperament; gout first at 37, not hereditary. Four years ago, both feet and hands being attacked, took half a bottle at bed time; no particular operation, except a gentle perspiration; was so much relieved, that he could

walk about the room next day. Much chronic gout afterwards, and in two months had a return of severe symptoms; then took four bottles in the course of the month. All acute pain again removed by the medicine; but says, that his limbs have been distressingly weak ever since, and much disposed to œdematous swelling.

CASE VII.

W. J. aged 52; robust and almost of the sanguineous temperament; gout first at 49, not hereditary. Took one bottle at twice; it nauseated exceedingly, and acted much on the bowels; the immediate symptoms quite relieved; but long after he suffered so much dyspepsia, and severe occasional spasms affecting the stomach, that no suffering of pain would induce him to take it again.

CASE VIII.

D. W. aged 40; plethoric and corpulent; gout hereditary and first at 28. In a severe fit took the third of a bottle; in about four hours, the pain subsided into numbness; perspiration followed, succeeded by a sound sleep. Next morning walked down stairs; the head uncomfortable, tight and heavy; no effect on the bowels, which were confined; thinks it was diuretic. Some purgative salt removed the unpleasant feelings of the head. Not aware that the next fit returned sooner than usual; on its occurrence, took half a bottle, its effect sudorific; on the following day was able to take a short journey in a carriage; next fit returned soon, and attacked both feet (the previous time one foot). Took a third of a

bottle; it gave immediate relief.—Gout returned very soon and very severely. Again took the medicine, when it acted both as a purgative and a sudorific, but only in a slight degree mitigated the pain. This fit, which lasted several weeks, left the head for a month after extremely uncomfortable; his feelings were wretched; had often *horrors*; was for a long time very dyspeptic, and suffered much from chronic weakness of limbs.

CASE IX.

H. B. aged 62; robust, plethoric, of sanguineo-nervous temperament; gout first at 47, not hereditary; has taken the eau medicinale freely, and has been a great advocate for its effects, finding it always prove a quick palliative; but confesses that the fits, which formerly observed distant intervals, have, since the use of this medicine, so much increased in frequency, that of late he is almost always more or less under the influence of gouty pains in the limbs.

CASE X.

O. J. aged 60; slight in limbs, but corpulent in the abdomen; of nervous temperament; gout first at 25, not hereditary. Took the medicine three years ago; half a bottle in the morning, the remainder in the evening; says that it proved almost fatal; produced constant nausea, violent pains in the abdomen, without any effect upon the bowels for five days. The paroxysm was very much relieved. Concluding that he had not taken the medicine properly, resumed it shortly after, both for the

purpose of removing the remaining symptoms, and to act as a prophylactic; but the gout returned soon, and with its usual severity.

CASE XI.

T. K. aged 57, corpulent and plethoric, of nervo-sanguineous temperament; gout hereditary, and first fit at 30. When attacked severely in the feet, took one-third of a bottle; it produced much delirium, and excessive perspiration. Opium affects his head in the same manner, in a slight degree. Next day took the remainder of the bottle, with a similar effect. It did not disturb the stomach and bowels. No relief to the symptoms. At intervals only of a few days, took a second and a third bottle in divided doses, being resolved to give full trial to the medicine. Again the same effects as before, but in a less degree. The gout not checked in its course, and it progressively attacked the feet, knees, hands, and elbows. Relates, that, after the third bottle, he seemed to gain some motion of the fingers, and felt that the medicine was searching every part; yet the fit was of six weeks' continuance, and, upon the whole, not apparently relieved by the medicine.

CASE XII.

C. L. aged 51, corpulent and plethoric; of nervo-sanguineous temperament; gout hereditary, and first fit at 42. In a severe attack, took at first half a bottle. It nauseated much, caused great perspiration, acted strongly

on the bowels, and removed the pain in three hours. Took the remainder of the bottle, at divided doses, within the next three days. In four days could just walk without assistance. Has taken altogether six bottles, dividing each into four doses. Does not think his stomach particularly weaker since, or that he is more nervous; but reproaches the medicine with having caused an extraordinary frequency in the returns of his gout.

CASE XIII.

P. A. aged 28, moderately robust; plethoric, of sanguineo-nervous temperament; gout first at 21, not hereditary. In a severe attack, took a bottle at two doses; it nauseated very much, but had not any other operation, and did not relieve the symptoms.

CASE XIV.

T. W. aged 41, formerly rather stout and corpulent, but now thin, and much worn in constitution; truly of the nervous temperament; gout first at 28, not hereditary. Has taken three bottles; the first at twice, with an interval of a few hours. It produced very severe cholera. The stomach was left remarkably weak for some time. The affected parts were relieved from pain almost immediately, yet remained a long time much debilitated. The second bottle acted as before, but less severely, and had not much influence on the symptoms. Never suffered in the loins, as he has done of late severely, before taking the medicine. The third bottle

had less operation than the second, and did not in the least degree influence the painful symptoms.

CASE XV.

C. W. aged 64, robust, corpulent, and plethoric; almost of the true sanguineous temperament. In a severe attack, took a bottle at two doses. It produced much perspiration, but no other sensible operation. The symptoms which were at first relieved, returned in a few days, and continued more tedious than on any former occasion; indeed he was hardly free from them before a fresh fit took place. For some time, "lost the natural feelings of his feet very remarkably. They were so cold and benumbed, that he feared he should never recover their use."

CASE XVI.

L. S. aged 46, corpulent and plethoric, of sanguineo-nervous temperament; of highly bilious diathesis; gout hereditary, and first at 35. Began the medicine three years ago. It acted as a narcotic, producing sleep; caused some perspiration; and affected the bowels moderately. One bottle removed the painful symptoms of the paroxysm, leaving only weakness; but the general debility was also considerable and of long continuance, attended with many nervous symptoms. On a second occasion took another bottle in divided doses, which left behind still more remarkable debility; and, for a long time, much trembling of the hands and knees.

CASE XVII.

S. B. a female, aged 50, corpulent, plethoric, and of nervo-sanguineous temperament; gout hereditary, and first at 47. In the first fit took one bottle in three doses, without any marked operation, and with slight relief only to the symptoms.

CASE XVIII.

C. T. aged 48, robust, plethoric, of strong bilious diathesis, and of sanguineo-nervous temperament. Gout hereditary, and first at 40. In an attack of both feet, took half a bottle with much relief to the symptoms. It acted as a purgative. Contrary to custom, the paroxysm returned in the same year. Repeated the medicine, taking a bottle at twice with only a few hours interval. It now acted as a strong poison, producing severe cholera. His stomach was left for a long time much weakened, his limbs very œdematous; and, in mind and body, he experienced a depression and languor, which he had never before known.

CASE XIX.

W. M. aged 42, slight, and of the nervous temperament. Gout hereditary, and first at 28. Has taken several bottles of the medicine. The first gave immediate relief; it acted slightly as a purgative and diuretic. In the subsequent use of it, the paroxysm became so

slightly relieved, and returned with such unusual frequency, that he gave up the remedy in despair. He adds, that it rendered him very bilious, highly nervous, and his limbs became exceedingly weak and much affected with cramp.

CASE XX.

S. J. aged 41, of nervo-sanguineous temperament; gout first at 30, not hereditary; very subject to violent attacks of the disease. In the beginning of a paroxysm, took a bottle in two doses; and not finding the smallest relief, did not persevere; but does not ascribe any ill consequence to it.

CASE XXI.

W. L. aged 42, corpulent, rather plethoric, of the nervous temperament, with a scrophulous diathesis. Gout hereditary, and first at 34. Has altogether taken twelve bottles of the medicine (procured from *Befort*) without any discoverable effect either of benefit or injury, although in one attack took two bottles, divided into four doses, with an interval of a few days only between each dose. In all this gentleman's attacks, the gout has assumed chiefly the indolent chronic character.

CASE XXII.

G. K. aged 55, thin and delicate, of the nervous temperament. Gout hereditary, and first at 28. A butcher.

Two years ago, having an attack in both feet, took a bottle at twice. The first dose stupified him. Awoke in a strong perspiration after seven hours' sleep. It had not any sensible effect on the stomach and bowels. Afforded great ease. He took the remainder of the bottle on the succeeding night. Appeared to the by-standers to be stupid and convulsed. So much mended on the following morning, that he stood in his shop, and, it being warm and settled weather in autumn, rode out in a cart. Suffered a severe relapse in the ankles, instep, and toes of each foot, and both knees. Repeated another bottle with similar immediate effects, but was left very weak and low. Gout returned, in the hand, within two months. Again took the medicine with similar inconvenient action on the nervous system, but with less controul over the symptoms. Reproaches the medicine with having produced such weakness in the back and lower limbs, that, ever since, he has been under the necessity of using a pair of crutches.

CASE XXIII.

B. R. aged 41, corpulent, plethoric, of nervo-sanguineous temperament, and of very bilious diathesis. Gout hereditary, and first at 28. In a severe attack, took half a bottle at bed-time, without any immediate effect; the remainder in the morning. Violent cholera ensued, with much perspiration. The symptoms quickly removed. No remote ill consequence on that occasion. In the next fit took two or three bottles, in divided doses, within a few days, but without much relief. On the following attack, still believing in the powers of the medicine to give ease, took four bottles in the course of eight days.

It did not at all mitigate the symptoms, and the following distressing results were produced. The bowels were left very inactive. He was afflicted with hypochondriasis. The limbs were extremely weak and œdematous, and even ascites took place in a considerable degree. The muscles of the whole arm were remarkably relaxed, and of the thumb in an extraordinary degree. Indeed, in this gentleman's case, the indulgence in this medicine brought with it such dangerous symptoms, that he considers himself as having narrowly escaped from death*.

Sir Everard Home himself in describing the power of the Eau Medicinale, mentions that he has ascertained it more than six times by experiment on himself, in the paroxysm of gout; and this very fact, taken by itself, appears to me to point out that the medicine does not possess more than palliative power, even when it acts most favourably; and cannot have the least claim to be called "a specific remedy for the cure of gout." How much evil it tends to produce in the constitution, I think, is sufficiently exemplified in the foregoing cases.

In apology for the diffuseness with which I am discussing this part of my subject, I wish it to be remembered that the practice in gout at the present day, in this country, lies between the empirical and the regular treatment; and therefore it is of the greatest importance that the present question should be exposed in all its bearings, and the truth be brought to light. Is the practitioner to advise the use of *vinum colchici*, Wilson's

* I could add very largely to this list of examples; but as the material point in question appears to be sufficiently illustrated, I am unwilling to allot more space for this purpose.

tincture, or eau medicinale, upon the faith of their *specific* agency? or is he to treat his patient upon solid general principles, varied and adapted to each particular constitution; and suited to those accidental circumstances which alter every individual case, in some more or less material point? The investigation into the nature of the various gout nostrums, still remains an interesting and unresolved question.

With respect to the chemical part of the inquiry, we are confined within narrow limits; for, as is well known, the substances of the vegetable kingdom do not admit of satisfactory results in any attempt of ultimate analysis, when we have it in view to identify any particular substance. From this cause, an easy and convenient cloak is afforded to the purposes of empirical compounders of medicine*.

* In order to compare the sensible properties of the eau medicinale, with the several medicines which have been supposed, as discoveries, to represent it, I instituted a series of experiments which I now subjoin, adding some particulars to my former report upon the subject.

In the examination of the following medicines, I used a very moderate temperature (not exceeding 120°) in bringing them to the consistence of syrup, leaving them finally to spontaneous concentration. After the first evaporation, I added distilled water to the mass, and re-evaporated; performing this process twice, so as to prevent any fallacious inferences which might arise from the difference of menstrua in the several preparations. The following were the results, the wording of which I have a little varied from my former statement; having conferred with an ingenious chemist on the best comparisons of the sensible properties of the medicines.

EXP. 1. *Eau Medicinale*.—(The menstruum, apparently, a French wine) colour, similar to the extract of gentian; smell,

The Peruvian Bark has been recommended very strongly by Dr. Tavares, a Portuguese physician, as pos-

very similar to it, inclining to the treacle lozenge; in taste, has some of the bitter of gentian, and slightly the flavour of opium; soon deliquesces after being dried.

EXP. 2. *Mixture of Wine of White Hellebore, and Vinous Tincture of Opium*.—Colour similar to the Eau Medicinale; smells slightly, and tastes strongly of opium; deliquescent.

EXP. 3. *Tincture of Colchicum*.—Colour, light brown; smell, rather similar to that of gum guiacum; taste, slightly but distinctly bitter, and entirely different from that of the Eau Medicinale; deliquescent.

EXP. 4. *Wine of Colchicum*.—Colour, deep orange; deliquescent.

EXP. 5. *Acetic Preparation of Colchicum*.—Colour, pale lemon; deliquescent.

EXP. 6. A very concentrated watery decoction of Colchicum.—Colour, bright yellow brown; only partially deliquescent, and so gummy, as to crack like dried mucilage of gum arabic. All these preparations of Colchicum, after the treatment described, resemble each other in smell and taste.

EXP. 7. *Tincture of Hedge-hyssop*.—Colour, almost black; no distinct comparable smell; taste, very bitter, like Taraxacum; dried and exposed in a damp apartment, very slowly and scarcely deliquesces.

EXP. 8. *Wilson's Tincture*.—(The menstruum, apparently, diluted spirit); colour, pale brown; smell, resembling that of extract of gentian mixed with orange peel and an aromatic; taste, similar to what is afforded by the substances just mentioned mixed together; deliquescent.

EXP. 9. *Reynolds' Specific*.—(The menstruum, evidently, rum) colour, blood-red, resembling compound spirit of lavender evaporated; smell, slightly similar to benzoic acid; taste, at first sweet, but afterwards imparts to the tongue a slight taste like that of benzoic acid; deliquescent.

In order to compare the quantities of solid residuum to be

sessing the power of cutting short the paroxysm, and rendering its returns less frequent. He quotes further

procured from the different preparations; and the relative solvent power of the menstrua for colchicum; one hundred grains of the following substances were inspissated by a gentle heat, brought to the same state of consistence, and then accurately weighed:

		Grains.
EXP. 1.—	100 grains of Ean Medicinale yielded.....	5 $\frac{3}{4}$
2.	————— Wilson's Tincture.....	11
3.	————— Reynolds' Specific.....	9
4.	————— Spirituous Tincture of Colchicum	2
5.	————— Acetic Preparation of Colchicum...	$\frac{3}{4}$
6.	————— Wine of Colchicum (so called, from Fisher's, as directed by Sir Everard Home).....	8
7.	————— The pure wine itself before dis- tillation.....	41 $\frac{6}{10}$
8.	————— The liquor obtained from the wine, the spirit being drawn off by distillation.....	61 $\frac{5}{10}$
9.	————— A watery decoction, concen- trated to the same strength as Sir Ev. Home's preparation...	9 $\frac{1}{4}$

Hence it appears from the preceding experiments, that if conclusions were to be drawn from the sensible properties of the different medicines, we should say that the quack preparations are unlike each other, and also different from any preparation of colchicum; but when we again reflect, that active substances may be entirely disguised by admixture with those which are inert, we cannot form any positive inference upon such grounds, and a more solid basis seems to present itself for the determination of the question, in estimating the agency of the several remedies on the human subject. In the medicines Exp. 2 and 3, it seems probable that extraneous

in its favour the authority of Dr. Held, and his remarkable praise of the remedy, in the following words: “*Uno verbo, cortex peruvians in podagra divinum est remedium**.”

Dr. Small also, in the paper already quoted †, offers a

matter is added for the purpose of disguise ; for the eau medicinale, although yielding less residuum, is much the most active substance. In regard to the preparations of colchicum, the watery decoction contains the largest proportion of solid matter, but, I must observe, that it is a turbid mixture, and contains much mucilage. In the vinous preparation, Exp. 6, when the residuum of the liquor itself is deducted, rather less is to be referred to the solvent action of the menstruum, than in Exp. 4, but more than in Exp. 5. The spirit (alcohol) therefore may be considered to dissolve most of the principles of the colchicum ; and with respect to the acetic preparation, I may observe, that if it were desirable to make it more concentrated, it could be rendered so by adding more of the roots than the Pharmacopœia directs ; but for myself, I must say, that I find it sufficiently active, according to the present formula.

The acetic acid, as in the example of the black drop, and still more in the acetate of morphia, appears to effect a useful modification in the medicinal properties of many substances of the vegetable kingdom. I must again remark, that it is from accurate practical observations, we are to form our conclusions as to the nature and comparative strength of these medicines. I have already delivered my general opinion ; and, in order to a further elucidation of the subject, I am now engaged in a series of experiments with the administration of the different substances to the dog. I shall relate my results in an Appendix to the present Treatise.

* Observationes et Epicrisis de Corticis Peruviani salutari et proficuo usu in Podagra.

† Medical Observations and Inquiries, vol. vi.

favourable testimony of the effects of bark, freely administered in the intermissions of pain and fever, and when the local inflammation has abated.

I have not hitherto been induced to make trial of this medicine in a paroxysm of gout; both from the perfectly satisfactory success of other treatment, and from an unfavourable idea which I entertain of the propriety of bark, in the circumstances under which this disease occurs.

Some patients inform me that they have made trial of the cinchona (*cordifolia*) in ample doses, as soon as the fever in the paroxysm has subsided; but it was not attended with good effects; and in a few instances, I learn that it very much disagreed. Guilbert and Hallé appear to think favourably of the use of bark in gout, to be administered on the same principles as in treating an intermittent; with great freedom therefore in the remission of the symptoms: but they candidly observe, that they do not consider the propriety of the practice as yet sufficiently established. M. Guilbert, when speaking of this remedy, gives a fresh proof of his caution with respect to purgative treatment; suggesting that some bitter should be combined with the bark, to prevent its action on the bowels.

Sudorifics.—Medicines of this class, as tending to debilitate the stomach, should be given with some caution. Antimony, in small doses, and conjoined with opium, in order to lessen the stimulating action of that medicine; and also with calomel, when it is used as a purgative, or as an alterative, is an important remedy, and has always appeared to me very useful. I consider, however, that the skin should not be made a channel of evacuation, either for the removal of the fulness of the

habit, or for the diminution of the local inflammation. When the treatment is such as to relax the skin very powerfully, an obvious disadvantage follows; namely, that of an increased susceptibility of the surface of the body to changes of the atmosphere, at the period of convalescence; and the consequent danger of relapse. A hot and dry skin will be much relieved by free sponging with tepid vinegar and water; by cool drink; and by a well-ventilated apartment, kept at a moderate temperature. All sources of stimulus to the circulation being avoided, the heat of skin soon yields to the action of the purgative, diuretic, and sedative medicines.

Narcotics.—Opium, under proper management, is a remedy no less advantageous than powerful for the relief of the most distressing of all the symptoms, *the pain of the disease.*

Although Sydenham, upon the humoral principles of practice, considered the pain to be “the disagreeable remedy of Nature,” and “a security to the patient’s life,” he permitted, in the event of violent pain, a dose of laudanum in the evening.

Warner expatiates warmly on the comforts which laudanum afforded him. He employed, with great propriety, a watery solution* of opium; and occasionally also an anodyne elixir, the formula† of which is complicated, and constitutes too heating a preparation. The good effects which opium is capable of affording, are dependant on the manner of its use, both as to the preparation and dose, and other points which I shall men-

* A full and plain Account of the Gout, p. 166.

† *Ibid.* 164.

tion: accordingly as it is administered, it may either aggravate or relieve the severity of the sufferings. M. Guilbert enters into the sentiment of Sydenham, and, in general terms, advises patience rather than any active medicine; but grants that *Dover's powder* has sometimes proved very serviceable.

Dr. Cullen observes*, "The opiates give the most certain relief from pain; yet, when given in the beginning of gouty paroxysms, they occasion them to return with greater violence."

As a rule of practice, always requiring careful consideration, any excessive state of inflammatory diathesis, and a constipated state of bowels, should be removed previously to the administration of opium.

With respect, however, to the first part of this statement, we have to estimate the genuine effects of the stimulus of pain on the action of the heart and arteries, as well as upon the immediate vessels of the affected part. Sir Everard Home, in an interesting paper†, "On the Influence of the Nerves upon the Action of the Arteries," has related several experiments which unite very well in the support of my present conclusion, that in any inflammation dependant on local and general irritation of the nerves, our rules of treatment must often be varied from that which we observe in the primary excessive action of the vessels in *common* inflammation, from which the nerves become affected in a secondary manner. On many occasions, in the gouty paroxysm, when the patient has described the pulsatory throbbing of the inflamed part, to resemble almost the successive blows of a ham-

* First Lines, part 570.

† Phil. Trans. 1814, part ii,

mer; when the heart has been in inordinate action; and when the inflammatory diathesis has appeared altogether urgent, I have stood by the bed-side and witnessed the happy power of a free administration of opium, in causing an abatement of the action of the vessels, and producing universal tranquillity in a short time*. Nevertheless, under the circumstances which I have just stated, we must employ united means; and although the use of general bleeding may be dispensed with on most occasions of this kind, we must avail ourselves of other modes of relaxant practice, in addition to the sedative influence of opium. In this intention is comprehended the action of purgative and diuretic medicine†, and the procuring of a cool and soft state of the skin.

In an ordinary state of the bowels, and when pain is urgent, the full action of a purgative is not essential as a preliminary; and under these circumstances, I have met with success in the exhibition of the draught, p. 186, at regular intervals, conjointly with the opiate at bed-time in adequate doses; so that, by the united effect of each remedy, relief has been obtained without delay—the influence of the mercurial purgative being added or not, as the particular symptoms have required. Under this regulation, comprising a due action on the bowels and

* The effects of opium in the agonizing pain of tooth-ache, when that severe affection of nerve produces high sympathetic fever, also illustrate the present point of pathology.

† I have often found that opium, and especially the black drop, has itself acted as a diuretic. I may add, as a general position, that it has had this effect only with patients of the nervous temperament; or, with those whose nervous system has been in a state of morbid sensibility.

kidnies, I have not found the observation of Dr. Cullen, just now quoted, to be confirmed; but I have several times experienced, that if I have trusted to the purgative, or the opiate, singly, a re-action of the circulation and violent return of pain have taken place at night.

In reference to the method of employing the remedy now under consideration, I shall quote the following excellent remark of Dr. Sutton:—"In the use of this medicine, also, it must be observed, that the benefit is not connected with a small dose of opium; but the quantity is defined by its producing a complete cessation of pain*."

I have myself found the use of this medicine remarkably successful in its crude state, and when joined with a small dose of antimonial (in preference, *James's*) powder. The patient being furnished with twelve pills, each containing one grain of *crude* opium and half a grain of *James's* powder, may be desired to take one, two, or, if pain be very severe, even three at bed-time, as the first dose, and repeat one every hour or two afterwards, according to the degree of pain; this being the only regulation as to the quantity to be employed, when no contra-indications are present.

It is worthy of consideration, that so powerfully does pain modify the influence of opium on the nervous system, in every kind of disease, that it may be given in the boldest doses, without hazard or ill effect, when pain is intense†; and in no other way than by the active repe-

* Tracts, &c. p. 216.

† Lately a striking example of this fact occurred to me. A young woman, of delicate constitution, suffered severe torture from some branches of the fifth pair of nerves supplying the cheek; the pain being of equal intensity with that of the *tic*

tion of such doses, can it be really efficacious, when the occasions for its employment are urgent.

It has long been a desideratum that this medicine should be rendered as free as possible from its stimulating and heating qualities, which prove always more or less unfavourable to its anodyne effects, and often forbid its employment, even when pain demands its use. The Lancastrian or *black drop*, which is a concentrated preparation of opium procured from boiling and digestion in a vegetable acid*, is found to agree with many indivi-

douloureux. Even delirium ensued. Three grains of crude opium, and one grain of antimonial powder, were administered as the first dose; and of pills, containing a grain of opium and half a grain of antimonial powder, one or two were desired to be repeated every hour, until pain should be overcome. Twelve grains of opium were thus taken in twelve hours; the pain was removed, and not the least inconvenience was experienced. A fortnight after, a little pain threatening the return of former suffering, she had recourse to one pill; but this now very sensibly disagreed, and caused sickness, and confusion and uneasiness of the head. Abundant proofs of the truth of the position in question I might state, both from my own experience, and on the authority of authors.

* Dr. Armstrong, in his late valuable publication on "Typhus and other Febrile Diseases," gives the following account of this medicine: "The black drop was originally prepared, upwards of a hundred years ago, by Edward Tonstall, a medical practitioner of Bishop's Auckland, in the county of Durham, and one of the Society of Friends. The recipe, passing into the possession of a near relative, John Walton of Shildon, who also prepared that medicine, was found amongst the papers of his brother, the late Edward Walton of Sunderland, and by the permission of my much respected friend Thomas Richardson,

duals much better than any of the usual forms in which it is prepared; and of this fact I have had many convincing proofs. The chief advantage which it has appeared to possess, has been in much less disturbing the stomach during its immediate operation, and the head on the following day, than either the tincture or wine of opium prepared according to the *Pharmacopœia*. It is perhaps imagined that the vegetable acid does not dissolve the resin of the opium, which appears to be the most heating of its component parts, but this is not a correct opinion; and although the black drop, both from its menstruum, and from the palatable ingredients mixed with it, proves rather grateful to the stomach, I must consider it a heating preparation. The resin of opium, apparently from the influence of extractive matter united with it in crude opium, is in a considerable degree soluble even in

senior, of Bishop's Wearmouth, one of his executors, it is here inserted

“ Take half a pound of opium sliced;—three pints of good verjuice;—one and half ounce of nutmegs;—half an ounce of saffron. Boil them to a proper thickness, then add a quarter of a pound of sugar, and two spoonfuls of yeast. Set the whole in a warm place near the fire for six or eight weeks, then place it in the open air, until it become a syrup: lastly, decant, filter, and bottle it up, adding a little sugar to each bottle.”

Dr. Armstrong quotes the authority of a friend, in estimating the strength of this preparation, to be as one to three, compared with the ordinary tincture of opium. He considers from his experience that this view is just. I was formerly of the same opinion; but my later and more careful trials have led me to conclude, that the difference is rather more than as 1 to 2, but not so much as 1 to 3.

water. M. Orfila quotes some experiments*, to shew that the watery extract of opium is of all its preparations the most active; but that each of its component parts possesses more or less of an anodyne power; and of the unessential principles (if I may be allowed this expression), the resin, the most. He adds, that the resinous part, although separately and freely administered, does not inflame the mucous membrane of the stomach†. Mr. Battley, the chemist in Cripplegate, whose useful labours in preparing and preserving all vegetable medicines in their most pure and active state, amply merit the thanks of the profession, appears to have succeeded in forming a preparation of opium more perfectly sedative than any which we have before possessed. He states that he has entirely separated the resin and the salt, by methods of decomposition which he promises to make known, when he has confirmed his several opinions respecting opium, fully to his satisfaction. I have made trial of the fluid preparation, to which he gives the name of liquor opii sedativus, and must in justice declare, that it has, in a few instances, produced very satisfactorily all the effects of an anodyne‡ and a sedative; namely, in relieving pain, and procuring tranquillity and sleep, with less subsequent inconvenience to the nervous system, the skin, and the action of the bowels and kidneys, than from any other form of opium which I have used with equal freedom.

* *Traité des Poisons*, partie iii. p. 144.

† I have made some experiments on the dog, with opium, which I shall state in my appendix.

‡ By the term anodyne, I speak of a medicine relieving pain; by sedative, of a medicine more particularly allaying simple nervous irritation.

A gentleman, suffering from rheumatism affecting the fascia of the thigh, and still more from extreme nervous restlessness, so that without the aid of a narcotic, sleep was entirely absent from his eyes, took, in comparative trials, the black drop and the liquor opii sedativus. His preference was decidedly given to the latter medicine, which calmed his nerves and left him free from inconvenience on the following day. On the contrary, the black drop, although agreeing well with the stomach, produced heat of skin; and on rising in the morning, he was unrefreshed, and felt as if he had been too much excited. In one most urgent case of pain from gout, attended also with excessive nervous irritation, I found it necessary to give in divided doses one hundred drops in the course of three or four hours; and this, for two or three nights in succession. No inconvenience followed, and the patient was much satisfied with its superiority over other preparations of opium, which had been variously tried. In this case, I may add, the extract of stramonium (from the seeds) had also been given in free doses without success. So far as I have judged of the relative strength of this medicine, I should direct it in similar doses with the tincture of opium of the Pharmacopœia; and I am convinced that it may be much more relied upon as a sedative. In extreme pain, however, opium in any form deserves our full confidence; and should the symptoms be of intense severity, it will sometimes be desirable to choose it in the liquid state, on account of its more immediate power of acting*; but under the most usual cir-

* I was called in haste to a lady, suffering torture from the state of the gall duct obstructed by a biliary calculus. I never witnessed greater signs of agony. The features were distorted;

circumstances which occur, I derive more satisfaction from the administration of crude opium joined with a small dose of James's powder, for the relief of pain, than from any of the fluid preparations which I have employed. The small addition of the antimony materially serves to prevent the opium from restraining the action of the bowels, as I have found in comparative trials with the same patient; and certainly it also improves the sedative operation of that medicine. The *pulvis ipecacuanhæ compos.* is also an excellent form of opiate; and, when joined with the use of saline medicine, sometimes proves more useful than any other preparation. I have found small and repeated doses, much to be preferred to the larger at more distant intervals. When relief from weariness, restlessness, and uneasy feelings, rather than pain, demand the use of sedative medicine, we may choose between the *liquor opii sedativus*, and the next class of narcotics presently to be mentioned; or, if feverish action be present, and the symptoms first stated are merely sympathetic, we shall often be more successful with the conjoined use of saline and antimonial medi-

she hallooed with exclamations for relief; the pulse was scarcely to be distinguished; the extremities were quite cold; the whole appearance of the patient was frightful. All the alarming symptoms of the case were overcome by opium. I gave instantly 100 drops of the tincture in hot brandy and water, and repeated doses of 30 drops every quarter of an hour until its due action was evidently established: 190 drops were necessary to the desired effect. The first dose seemed of no avail. In one hour sleep took place, and pain never returned in any degree. The hot bath was used in conjunction with opium; and purgative medicines were freely administered, when the dreadful paroxysm of pain was ended.

cine, and aperients, than with any species of narcotic. In concluding the subject of opium, I shall take a brief notice of its insulated principle, morphia, stated to be the characteristic constituent of opium, and not very long since announced* by M. Sertürner. I have made trial, in two instances, with that combination of this principle with the acetic acid, which has received the name of acetate of morphia. It was prepared by Mr. Garden of Oxford-street, according to the formul† advised by M. Robiquet, and found to be much superior to the previous method of Sertürner. In the cases in question, I was driven to the necessity of choosing a new narcotic, for all the ordinary forms of opium had ceased to relieve; and the sufferings of the patients were intense. I had the satisfaction to find this new medicine agreed remarkably well; and it proved highly anodyne. With such very limited experience of this remedy, I forbear, at present, from any further observations respecting it.

Notwithstanding, however, these stated improvements in the mode of administering opium, there are some instances of particular idiosyncrasy of constitution, in which this medicine cannot be borne in any form. So repugnant is the nervous system to its influence in these cases, that, sometimes, even a moderate dose of the syrup of the white poppy produces a state of temporary delirium. This I have witnessed on two occasions; and in less aggravated examples of disagreement, the smallest portion of opium confuses the head, and deranges the stomach in a distressing degree. The exceptions to the advantages of employing opium, when pain, not depending on *common* inflammation, is urgent, fortunately do not often

* See Thomson's Annals of Philosophy, No. liv.

† This is stated in the French Pharmacopœia, 1818.

occur; but with such persons as do experience from it, in any dose or form, an increase of nervous irritation, so that even the relief from pain is an advantage too dearly purchased, it is incumbent on us to make trial of other narcotics; respecting which, I shall give a brief account.

The extractum hyoscyami, as a sedative and weak anodyne, is entitled to some reliance; but it has appeared to me to vary in its powers, more remarkably than any other in this class of medicines. Thus, with some individuals, three or four grains twice in the day, prove a quantity as much as can be taken without disagreement; while, in other instances, a drachm in the twenty-four hours is given without inconvenience: but when the system is not soon sensible to the influence either of this medicine, or of *conium*, which I should place in the same scale of power as a sedative, it follows, according to my experience, that a perseverance in the dose is quite useless, and tends only to weaken the stomach. The very opposite degree of effects produced by all vegetable medicines, does, in most instances, unquestionably depend in a great measure on their different mode and state of preparation, and consequent virtue*; but with respect to

* It not only happens, that the extracts of the vegetable substances are sometimes ill prepared in the first instance, but also that they undergo decomposition from long keeping, and particularly if in a damp situation. In order to have a vegetable medicine most active in its properties, the first step is to collect the plant in its best state of vigour; the next is the drying it by a proper heat; and the last, to reduce it to powder, and put it up for use in a closely stopped bottle, covered with dark paper to secure it from the influence of light; according to the method which the College directs. In this statement, however, I admit,

the statement just given of henbane, I have made reference to cases, in which the same extract was employed.

The humulus lupulus was strongly recommended by the late Mr. Freake, as a remedy in the paroxysm of gout; but from very sufficient trials which I have made of this medicine, I can assert, that it is quite undeserving of dependance. It is in any case a very weak sedative. This opinion is also supported by the experiments of Dr. Bigsby*.

The soporific medicine prepared from the inspissated white juice of the lactuca sativa, or common garden lettuce, introduced to our notice by Dr. Duncan, sen. is well entitled to good report; and I am happy in this opportunity of bearing testimony to its virtues. The dried inspissated juice exuding from the plant when wounded,

that an extract well prepared, and carefully preserved, is also fully entitled to our confidence.

M. Orfila observes (Waller's Translation, vol. ii. p. 50), with regard to the medicinal power of vegetable substances, "1st. That certain extracts prepared by expressing the juice of fresh plants, and evaporating them in a water-bath, are incomparably more active than those obtained by any other process. 2nd. That they are of a yellowish colour, which forms a singular contrast with the black colour of those found in commerce, which without doubt depends on the decomposition which these undergo by caloric. 3rd. That the extracts of those plants sold in the shops, differ greatly amongst themselves in regard to their action on the animal economy; that a great proportion of them possess no virtue; which circumstance, we are of opinion, ought to attract the attention of those Professors, whose duty it is to visit the shops."

* See Medical Repository, vol. iv. p. 287.

evaporated to the consistence of extract, to which the Professor has given the name of *lactucarium*, is the preparation of which I have made trial; and from my experience of its effects, upon a very extensive scale, in miscellaneous cases, I can much recommend it as a mild sedative, calculated to tranquillize without stimulating; to allay a cough; to promote sleep; and, in a *slight degree*, to relieve pain. But certainly the dose should be more freely administered, than I have understood to be directed; and indeed a smaller quantity than four or five grains at bed-time, to be repeated also at convenient intervals, in a similar dose, as occasion may require, would, according to my experience, be insufficient to produce any satisfactory result.

In comparing the efficacy of lactucarium with the *lactuca virosa*, I find the most decided preference to be due to the former medicine. I consider it indeed a very valuable remedy under certain circumstances; and have only to regret the great difficulty and expence which necessarily belong to the preparation*.

The use of the *datura stramonium*, or thorn-apple,

* Mr. Probart, of Great Portland Street, by my recommendation, made an extensive plantation of lettuces for the purpose of preparing lactucarium, which he keeps in great perfection. The milk was collected by means of cotton, and inspissated merely by the action of warm air; lest by any other mode of applying heat, the delicacy of its properties should be injured. He found the plant to afford most of the milky juice when in seed. It scarcely contains any portion in the early state in which it is brought to table. Being desirous to form an extract from the lettuce which should prove efficacious if used in larger doses than the lactucarium; be less expensive, and therefore more accessible for general use; he made some experiments, the result of which I shall state in his own words.

“When the plants are cut down, the milky juice for the

prepared and formed into an extract from the seeds of the plant*, has lately been revived to the attention of the profession, and much recommended as an active anodyne by Dr. Marcet†. I have, on several occasions, made trial of this extract, in free doses, in the severe pain of gout; but certainly with a most inferior effect to that obtained from opium employed in the manner I have described: and now on every occasion of prescribing for the relief of severe pain in gout, I almost confine myself to the use of crude opium, in union with James's Powder. But, to return again to those cases of exception, in which

most part, assumes a concrete form, and is concentrated chiefly in the rind and in the old leaves, such as are about changing to a yellowish colour. These parts, therefore, I select for my extract, rejecting the substance of the stalk and the young sprouts; and having macerated them for twenty-four hours in water, I boil them for two hours; and, without pressure, allow the clear decoction to drain through a sieve. I then evaporate it as far as it can be done with safety, and finish the process in a number of shallow dishes for the purpose, in the air."

From the success which attended the use of this extract, in the case of a gentleman who suffered severely from an irritable state of bladder, I have reason to think that it may prove an useful medicine. In the case to which I allude, the symptoms were extremely urgent, and the dose employed was ten grains three times a day. I must add, that various other sedatives had been tried with much less relief than was afforded by this extract.

* Mr. Battley informs me, that in his opinion the far more active properties of stramonium reside in the rete mucosum of the capsule.

† See a paper on the subject, *Medico-Chirurgical Transactions*, vol. vii.; to which, for further particulars of the properties of this medicine, I refer the reader.

opium is inadmissible from the idiosyncrasy of the patient, I must very conclusively give the preference to the stramonium over all the other narcotics which I have tried. It has most succeeded in relieving that kind of pain which is dependant upon, or immediately connected with, spasm of the muscular fibre; and, given at bedtime, it is well calculated to counteract the tendency to cramps. I have never found it necessary to commence the dose in a smaller quantity than a fourth of a grain twice a day; and, as to the maximum, I should say, that when used to the extent of ten grains in the twenty-four hours, in divided doses (of course gradually brought to this amount), if it do not produce decided advantage, its use should be entirely relinquished. In conclusion of the subject, I think it important to add, that I have obtained more decided advantage from employing the stramonium extract and lactucarium, in conjunction, than from the former separately; and in reflecting upon my experience with these medicines, I am tempted to affirm, that the former acts most as an anodyne, the latter as a sedative; and that where both effects are desired, they will be most favourably procured from the union of the two preparations. One gouty patient in particular, who experiences much inconveniencce from the use of opium, derives the happiest effects from this combination, and finds it competent to relieve even severe pain.

The *atropa belladonna*, or deadly night-shade, is a narcotic of well known activity, and in cases of pain and spasm, which have resisted other anodyne and sedative preparations, it is eminently entitled to trial*. In hoop-

* It is not imaginary to state, that all the narcotic vegetables have their own peculiar mode of acting, and which may be called *specific*. Hence the propriety of change, when any one narcotic disappoints our expectations.

ing-cough, and in one case of spasmodic asthma, I have been much satisfied with its effects; but in the gout, having succeeded in soothing by other narcotics, I have never yet been induced to employ it.

The *aconitum napellus*, or wolfs-bane, is quoted by Barthez (tom. i. p. 152), as efficacious in relieving the chronic pains of gout. From such little experience as I have had with this medicine, I am not disposed to give it confidence.

That I may not be misconceived in my opinion of the value of the narcotic class of medicines, in relieving the symptoms of pain and irritation which the gout produces; I must, in conclusion, state, that in the great importance which I do attach to them, I view them still as subordinate to the more radical means of treatment; as auxiliary only to the primary objects, of removing inflammatory action; of rectifying all the secreting functions; and adjusting the circulating system to a healthy balance.

Diet.—It is in this important particular, most generally, that the greatest errors are committed by the gouty patient. This fault is sometimes founded on his own love of habitual indulgence; but it is also connected with the false doctrines which he has imbibed concerning the disease, and with mistaken notions of the prevalence of debility.

The avoiding of every circumstance with regard to food, both in respect to its quantity and quality, which can produce hurtful excitement; and still more especially, the shunning, or very cautious employment of wine or any spirituous stimulant, are obviously considerations of the utmost importance. Not only, are the symptoms of the paroxysm materially aggravated and prolonged by errors of this nature; but the additional evil of erysipelas sometimes arises as a consequence, and

takes place either in conjunction with the gout, or as an immediate sequel to it. Of this result I have seen some very striking examples. Under very acute symptoms, the nourishment should be wholly fluid, and not stimulating. Boiled bread and milk is much to be recommended, and usually agrees perfectly well. A debilitated stomach, with weak constitutional powers, may sometimes require a diet rather cordial and supporting, even in the paroxysm; but this matter should be managed with much discretion. We should be careful not to support the disease, at the same time that we assist the powers of the patient; and on this principle, any stimulating article in the nourishment, should be subservient chiefly to the curative employment of medicines.

On the few occasions which authorise the use of wine in the paroxysm, it should, as a general rule, be given in dilution; and its cordial powers are, in general, most usefully obtained when mixed with some article of nourishment, such as, arrow-root, sago, or gruel. These nourishments, or panada agreeably spiced, will often alone prove sufficiently cordial during the urgent symptoms, although much languor and depression do prevail.

It is incumbent on me here to mention, that I have met with instances of such great peculiarity of nervous temperament, that even during the height of the painful symptoms, a little animal food and two or three glasses of madeira undiluted, have been really necessary in the daily plan of diet. Such material support will, however, seldom be proper, until the convalescent stage is fairly commenced; and the exceptions to the general rule, of slight and cautious nourishment, are of rare occurrence.

As a diluent beverage, rennet-whey is equally agreeable and useful. Sometimes the thirst of the patient is

so urgent, that he desires the coldest drink, and solicits even that the water should be iced. I have not opposed an inclination thus dictated by disease; nor have I seen any harm produced by the occasional use of such a refreshing indulgence. Yet the action of the medicines on the bowels and kidneys must be aided by frequent draughts of warm diluting fluid, as tea, thin gruel, or barley-water; or very warm water alone. Acidulated drink is usually unfriendly to the gouty stomach, and especially during the paroxysm; but I have met with some instances, in which even at that time the free use of lemonade has perfectly agreed. I need hardly mention, that, during the action of the mercurial purgative, acid matter should be avoided in every shape; and the warm fluids already mentioned ought alone to be in free use.

With the exception just expressed, the subacid fruits in season, as oranges, and grapes of a good quality, and apples roasted, may be included in the proper gratifications of the palate; provided they be used with moderation; and that they perfectly agree with the patient's stomach.

Bodily Exertion.—Sydenham advised that the patient should take daily exercise in a carriage, even in the beginning of a fit, except when in excessive pain. Such a degree of resolution as this would require, is not easily practised, nor can I assent to its propriety. It would indeed be incompatible with the attentive treatment suggested in these pages. The opposite extreme, however, of the entire quiescence and relaxing influence of the bed, continued for a longer period than is really necessary, is to be equally opposed. Practice affords us

abundant proofs in how great a degree a stiffness and debility of the limbs may be counteracted by moderate and early efforts of exercise.

Van Swieten relates * with pleasantry the story of the dancing-master, whose large family did not allow him to be long idle in nursing his gout.

Some few, on being threatened with an attack, have removed the symptoms by the spirited exertion of a long walk; but it is a doubtful experiment, and is much more commonly followed by serious disadvantage.

Dr. Small walked abroad as soon as the inflammatory action had ceased; and expresses his opinion, "that nine in ten of gouty cripples owe their lameness more to indolence and fear of pain, than to the genuine effects of the gout †."

In cases of great suffering, and of neglected treatment; and when the patient has, for a long continued period, been entirely confined to the bed, or by choice indulges in it; it is probable, that the excessive secretion of mucus in the urinary bladder (already described as accompanying the irritating quality of urine of high specific gravity) may indirectly, by its cementing quality, tend to the production of calculous concretion; and thus lead to the double tortures of the stone and the gout.

This statement, I wish however to be understood, belongs exclusively to the greatest mismanagement of the case; and to the consequent occupation of the bed for weeks and months.

Except under extreme circumstances, it may be said, the patient should every morning leave the bed for the

* Commentaries, § 1261.

† Med. Obs. and Inqu. vol. vi. p. 200.

couch or the chair, having his legs raised and supported in the most easy position; and, in proportion as pain and inflammation abate, should gradually employ such further exertion, as relieves, rather than produces irritation. But I must add, that, in a severe paroxysm, and during the height of the symptoms, the patient is of necessity confined to his bed in the most helpless state; and, in this case, so far from using any forcible attempts to move the affected limbs, thereby producing great increase of pain and irritation, I advise that they may be relieved and supported in the most studied easy position, by means of small pillows placed so as to cause the greatest relaxation to the muscles. A further comfort is also afforded by the use of a cradle*, applied so as to keep off entirely the weight and pressure of the bed clothes; still using sufficient immediate covering to the limbs to produce comfortable warmth.

I have witnessed many instances, in which the too early exertion of the limb, has, by the over-action of the weakened parts, produced relapse. This part of the management of the last stage of the fit, often requires a very delicate exercise of medical judgment.

The Passions.—Many marvellous cases of the immediate cure of the paroxysm are related, by ancient authors, to have been effected by the sudden influence of the strong passions of the mind, and more particularly by *terror*. Dr. Falconer very justly observes, that these

* This is easily constructed, by fastening with nails two or three segments of hoops, bent in a semi-circular form, to two parallel pieces of wood.

narratives “are rather matters of curiosity than utility, and what we can make no application of in practice*.”

It should, indeed, be our care to allay rather than to excite any violent emotions, which are so much the prolific offspring of the disease itself. We cannot too freely introduce hope, cheerfulness, and tranquillity; and the attention should be agreeably diverted. Fortitude and true philosophy are more valuable aids to the welfare of the patient, than the superstitious charms of the ancients; in proportion as it is better to improve the reason, than to deceive the imagination and subvert the judgment.

LOCAL TREATMENT IN THE PAROXYSM.

THE present consideration is truly a point of great importance, and offers in its details much useful opportunity of novelty. The inflammation of gout has never been treated upon fixed and regular principles. It has most commonly been left to its own injurious course, unchecked and unrelieved. By one kind of management, it has been nurtured, increased, and prolonged; by another treatment, it has with hazard been at once dispersed. Amidst such irregular proceeding, it becomes a valuable desideratum, to establish a settled mode of practice, which may at once be safe and useful. This view of my subject I shall now endeavour to offer, in a short notice of the chief topical remedies which are in use at the present day, or such at least as are not wholly discarded; and in describing the particular method of treatment which I am induced to recommend.

Leeches.—The immediate emptying of the vessels under inflammatory action, may appear at first sight very

* Upon the Influence of the Passions on the Disorders of the Body.

consonant to the just doctrines of pathology; and the propriety of the practice has been advocated by authors in general. Some practitioners prefer the more prompt method of opening one of the distended veins by a lancet, expecting in this manner more effectual and speedy relief*.

From an attentive experience of the effects of local bleeding, however performed, I am led to consider, not only that such treatment is unnecessary, but that it is in most instances injurious, and, even when apparently useful in relieving the immediate symptoms, is very seldom attended with permanent advantage.

It has always appeared to me to be a correct and sound principle of practice in local inflammations of every kind, that, whenever their violence is such as to influence the action of the heart and arteries in any considerable degree, the abstraction of blood should be made from the arm, rather than from the part affected; but that when the inflammatory action is almost entirely local, the depletion of the vessels should be local also. With regard to the inflammation of gout, however, we are to reflect, that it forms only a part of a constitutional disease, and that it is often of a more fugitive nature than any other kind of inflammation.

In a few instances after the application of leeches, I have seen that the inflammation has very suddenly transferred itself to the other limb; hence pointing out, that the constitutional causes were in active force, and not to be relieved by local loss of blood. I do not argue that it is hazardous to the constitution; nor can it be liable to such an imputation: but I must add, that the indiscriminate use of leeches in gouty inflammation is by no means innocent in its local consequences. An increase of inflam-

* The use of leeches, when pain and inflammation are violent, is much recommended by the French physicians.

matory irritation and pain now and then follows their application; but as a more serious, because more lasting, evil, I have often seen the debility of parts increased in this manner; and when much blood has been removed, a troublesome œdema of long standing has followed, attended with a corresponding incapacity in the action of the nearest joints. In three instances, I have witnessed the effect of taking blood from one of the distended veins near the foot, when violent gout was situated in the toe; and I was by no means satisfied with the result. An increase, rather than a diminution, of pain, was the consequence in two of the examples; and in the third, much local weakness seemed to be the result. The affected parts during the inflammation are in a state of such high irritation, that it is requisite to exercise great delicacy in the local remedy which is used, both as to its nature and the mode of application.

It has invariably happened within the range of my experience, that active constitutional treatment, as already detailed, together with the employment of local evaporation in the mode presently to be described, has entirely superseded the necessity of more doubtful and complicated means for the removal of the inflammation.

Vesicatories and Irritants.—I have not sought for any experience in this class of remedies in acute gout, from the objection which I conceive to their most probable agency; and from my being satisfied with the efficacy of other means.

Cullen (Par. 565, 566) speaks of blisters as effectual, but hazardous; and expresses the same opinion of *stinging with nettles*. The burning with *moxa* (the Chinese mode of cauterizing) must be considered as a species of painful blistering. Sir William Temple relates the cure

which he received from it*. Hippocrates employed the burning with raw flax in the neighbourhood of the affected joints. Of *ludicrous* applications, a copious list is furnished in the following amusing extract from the curious *τραγοποδαγρα* of Lucian, which I copy from M. Guilbert.

Terunt plantagines, et apia.....
 Et folia lactucarum et sylvestrem portulacam.
 Alii marrubium ; alii potamogeitonem ;
 Alii urticas ferunt ; alii symphytum ;
 Alii lentes afferunt ex palustribus lectas ;
 Alii pastinacam coctam ; alii folia persicorum,
 Hyosciamum, papaver, cepas agrestes, mali punici cortices,
 Psyllium, thus, radicem ellebori, nitrum,
 Fœnum græcum cum vino, gyrinum, collamphacum,
 Hyarissimam gallam, pollen hordeaceum,
 Brassicæ decoctæ folia, gypsum ex garo,
 Stercora montanæ capræ, humanum oletum,
 Farinas fabarum, florem asii lapidis :
 Coquunt rubetas, mares-araneos, lacertas, feles,
 Ranas, hyænas, tragelaphos, vulpeculas.
 Quale metallum non exploratum est mortalibus ?
 Quis non succus ? Qualis non arborum lacryma ?
 Animalium quorumvis ossa, nervi, pelles,
 Adeps, sanguis, medulla, stercus, lac.
 Bibunt alii numero quaterno pharmacum :
 Alii octono ; sed septeno plures.
 Alius vero bibens hieram purgatur :
 Alius incantamenti impostorum deluditur, etc.

Warmth.—The various contrivances of warm covering to the affected part, for the purpose of producing pers-

* See his Works, vol. iii.

piration ; or, in the fanciful language of humoral pathology, *to invite the deposition and discharge of the morbid matter*, although apparently full of prudence, and seemingly warranted by the custom of ages, do indeed belong to the worst part of the ancient practice. Combed wool, or socks and bootikins, have been favourite modes of fulfilling this plan of treatment*. One gentleman, who gave ample trial to the use of bootikins, informs me, that his ankle joints have never recovered the excessive weakness which they occasioned. From several patients I have obtained the same information, of much permanent debility being produced by this treatment. Some have, in a more partial manner, applied silk oil-skin to the affected part, with relief to the inflammation by means of the perspiration produced, as in the effect of bootikins ; but which was not a sufficient compensation for the subsequent weakness to which it led. *Flannel and patience* still form the adage of many, whose caution is greater than their judgment. It is obvious that the confinement of morbid heat by the warm covering, on the one hand, must serve to increase pain, and prolong the disease ; or if, on the other hand, perspiration be much induced, the debility which is consequent on this mode of evaporation becomes a secondary evil of great magnitude. The most tedious and intractable cases which I have seen, have been those in which the relaxing practice has been carried to its fullest extent, both by local accumulation of warmth, and by the influence of oppressive covering of the bed, together with corresponding errors in the regimen throughout. Even in those occasional exceptions to the general course of the complaint, in which a fit of the gout has taken place, rather as a remedy than a disease, it is proper that only moderate warmth of cover-

* Gardiner on the Gout.

ing should be employed; and I may add, that, in the paroxysm, the patient should at all times prevent the heating influence of the fire in his apartment from being received on the inflamed parts.

The Pediluvium, Fomentation, &c.—Hot bathing of the feet is inadmissible while any inflammation remains; and I have even seen the symptoms reproduced by its employment very remarkably, where no hazard of such an occurrence had appeared to exist. Those who have adopted this practice, as a remedy in the paroxysm, have, for the most part, informed me of very unfavourable results. A lady of great nervous sensibility, on the fourth night of the paroxysm, immersed her feet, one of which was much inflamed and swollen, in hot water for ten minutes. Almost instantly the gout quitted the foot; and such was the progress of its transference, that, during the night, it affected the knees, the elbows, and the wrists; never again, in the course of the fit, which was of unusual length, returning to the foot. A gentleman, having gout severely in each ankle joint, immersed the feet in hot water with bran, which gave him present ease, but seemed to occasion a quick removal of the inflammation to the knees and to the elbows, which followed in a few hours; and a tedious fit ensued. I have known, in other instances, that this treatment has materially aggravated the local inflammation and pain; in which case, no removal of disease has taken place. Poppy fomentation, and the vapour of hot water impregnated with aromatic herbs, have been patiently tried for the relief of pain and inflammation; but seldom with any good result, and more commonly with manifest disadvantage.

Also when the cleanliness and softening of the skin

of the surrounding parts, together with that immediately affected, are desired, during the paroxysm, the use of free sponging with tepid water is much to be preferred to immersion. *The muriatic acid bath*, first adopted in France, is still occasionally employed in this country. I have witnessed its effects in one patient, who found from it an aggravation of pain and inflammation. Some inform me, that they have obtained relief from this remedy. It appears to me an injudicious practice, and to be either too exciting, or too relaxing, according to the degree and continuance of temperature at which it is employed.

I know a gentleman of great intelligence, much subject to gout, who is partial to the use of heated air as a remedy, which he obtains by burning alcohol at the extremity of a tin tube, bent in its form, and connected with a wooden cradle, to be received under the bed clothes, so that the air has free circulation. He finds that, in about twenty minutes, it begins to produce perspiration, which soon becomes profuse. He states that he has materially reduced the duration, and the degree of pain of his paroxysm, by this treatment; and is convinced that he has also, on several occasions, effectually carried off the gout by the skin. I had the opportunity of witnessing the operation of the remedy in this gentleman's case, and came to the following conclusions. It appears to me that this process constitutes a very ingenious mode of procuring to the body a medium of warmth, in which, from the blood being universally determined to the skin, general perspiration, under favourable circumstances, becomes speedily excited. I think it calculated to answer a very useful purpose in the cold fit of any fever which is forming; and in the chilled state of body after any hazardous exposure to wet and cold. As it regards the gout, notwithstanding the favourable report which

this gentleman makes of its effects, I do not consider it an appropriate remedy on a general principle; although there may be circumstances, under which, in the first invasion of the fit, its use would be highly rational. I have lately stated my objection to the plan of excessive perspiration in the cure of a gouty paroxysm; and, in addition to what I then observed, I may again advance the argument, that as this disease has its seat only secondarily on the surface of the body, and is, in its true character, radically depending on an obstructed and vitiated condition of the digestive organs, and the liver especially, with a disordered state of the nervous system; so, in conformity with this view, the use of diuretic purgatives, of mild alterative medicine, and of sedatives, forms the most direct and efficacious method of cure.

This is the statement regarding the present remedy which I formerly offered; since which, I have had an inducement to make trial of it in a very urgent case of gout, which had been excited entirely by exposure to cold. Lumbago was joined with acute gout in the feet, knees, and hands. The patient was so heavy and corpulent, that there was unusual difficulty in carrying into full effect the general plan of treatment which I have been studious to recommend. I employed, therefore, this apparatus, and not without some good effect. The heat immediately produced was more than comfortable, and indeed rather distressing; and very copious perspiration followed, which proved useful. But, as I expected, it was only auxiliary to the other treatment, which was indispensable; and which, to the utmost extent that it could be conveniently employed, consisted of the means already laid down. I found, in this case, great advantage from the use, at short intervals, of Dover's powder and saline medicine; and had indeed great cause to approve this seda-

tive part of the treatment. I administered, occasionally, calomel and James's powder, with colocynth, and the draught, p. 185, every day in the early part of the paroxysm, with more or less frequency. The lotion in the form of poultice also proved highly useful.

I have no doubt that there are many circumstances of disease, under which, it being difficult to effect perspiration by ordinary means, the use of this apparatus will be found particularly serviceable; and the facility with which it is used is no slight recommendation of the remedy.

Poultices, as a mode of evaporation, either simple or variously medicated, have been generally considered useful. Sydenham speaks of having derived some occasional benefit "from a cataplasm made of white bread and saffron boiled in milk, with the addition afterwards of a small quantity of oil of roses." Observation has convinced me, that a free employment of relaxing poultices, made in the ordinary way, has the disadvantage of increasing œdematous swelling, and subsequent debility. I have, however, found great cause of satisfaction in the occasional use of a poultice, made with bread which has been scalded with boiling water, pressed through a strainer to dryness, and again rendered of sufficiently soft consistence by means of the lotion which I shall presently describe. It is then to be applied, thick in quantity, very moist, just tepid*, and without any intermediate covering between it and the affected part. When the hands and feet only are the seat of complaint, I have for the most part been in the habit of using this poultice at night only;

* So rendered by adding to the lotion enough of boiling water. A hot poultice is very injurious in its effects; for, in addition to the debility to which it leads, it renders the part very liable to be affected with rheumatism.

a time when the patient wishes for quietude, and cannot allow of the same frequency of attention that is offered in the day; and which is necessary to the effect of evaporation by means of the lotion. But for the relief of the knee and elbow, I have usually found the poultice to be very much more useful than the lotion applied by compresses; and have directed it to be repeated twice or thrice in the twenty-four hours, according to circumstances. Many patients have extolled the effect of this poultice in very high terms; and my late experience with it, authorises me to recommend it as an excellent remedy. I was consulted by a gentleman, ill with acute gout, who had previously been applying poultices of bread and water without the least relief; but on using the one now under description, he experienced prompt and decided benefit. For a short time after the discontinuance of the poultice, the part should be covered with a single layer of flannel; for the obvious reason of preventing the supervening of rheumatism.

M. Guilbert offers the formula of Pradier's cataplasma, and enters at some length into its *modus operandi*; recommending it as a remedy in the paroxysm, unless "inflammation runs high, with a great sanguineous determination to the parts." He advises it also when gout is hovering about; and says that it is desirable by this means to develope its action. I quote the following translation from Dr. Johnson.

"Pradier's Cataplasma.—The registered composition of this remedy stands thus: Take balsam of Mecca six drachms, red Peruvian bark one ounce, saffron half an ounce, sage an ounce, sarsaparilla an ounce, rectified spirit three pounds. The balsam of Mecca is to be dissolved separately in one third of the spirit; the other ingredients being macerated for forty-eight hours in the

remainder. The two liquors are then to be filtered and mixed together. When wanted for use, this tincture is to be diluted with two or three parts of lime water, and as a precipitate takes place, the bottle is to be shaken when the contents are poured out. A poultice of linseed meal is now to be prepared in sufficient quantity to envelope the foot and leg up to the knee. It should be of a good consistence, and spread an inch in thickness, on a napkin of proper size. If both feet are affected, it requires about three pounds of linseed meal. When the poultice is spread even, and ready for application, about two ounces of the above-mentioned liquor are to be diffused over the surface of the cataplasm, and then the foot and leg are to be immediately enveloped with it; and over all, another covering of flannel, secured by a roller. It is not to be changed for twenty-four hours."

This composition is very complicated, and most probably would be found too stimulating in acute gouty inflammation. With the omission of some of the ingredients, which appear to be quite insignificant for external use, it may deserve trial in obstinate cases of chronic gouty action; provided the more simple form of poultice, which I have prescribed, should fail.

When the knee has been still affected with uneasiness and tenderness, either as consequent to the acute form of the complaint; or altogether from being under the influence of chronic gout; I have sometimes, and with a very good result, found it convenient to apply the soap plaster of the Pharmacopœia, spread on linen or soft leather; using it at night only, and the lotion in the day; or continuing it to the part wholly, as the only local remedy. I have to conclude with observing, that, during the paroxysm, if the affected parts are cool and free from pain at the period of bed-time, all local treatment may be

omitted till the following day; and no unusual covering of any kind need be used.

Of other Modes of Evaporation.—The practice of Dr. Kinglake (for so it may be designated, although originally derived from Hippocrates*, and other ancient writers) seems, and most justly, to have fallen into disuse among most of those who were its first promoters. Dr. Kinglake introduced this treatment on the narrow principle of considering the gout as a local disease, and as analogous, in this respect, to simple inflammation. Such confined and erroneous pathology requires not any serious refutation. That the gout is sometimes *comparatively* very much a local complaint, is a position not to be denied; and the *cold water* practice in the paroxysm has occasionally been successful. It appears to me, however, a more safe and correct conclusion, to consider that gouty inflammation is always more or less intimately connected with the system; and that it requires a treatment in conformity with this principle. When we reflect on the facility with which gouty inflammation is transferred from one part to another, the sudden repelling influence of intense and continued cold must appear, even in theory, full of hazard; and in practice, its bad effects are now notorious. Indeed, from all that I can learn of the practice of applying cold water, the relief is never so certain as the danger; and I could enumerate many instances, in which the patient has, in the very beginning of the treatment, received a timely alarm from a sudden spasm at the stomach or diaphragm; and well-attested cases of danger might be quoted without number;—a few also, certainly, of speedy death as the consequence.

* Aph. xxv. sect 5.

The immediate abatement of pain is often followed by numbness, increased swelling, and much continued uneasiness; and in muscular and tendinous parts, its influence is by no means favourable to the free return of motion. Even in its most successful operation, the effects are too suddenly produced. The cure is rather local than constitutional; whereas we should consider that we have a two-fold object to accomplish.

In this disease, indeed, it must be steadily kept in view, that our first and sovereign object, is the constitutional treatment; and that local treatment is of secondary importance:—but that this last consideration is nevertheless a point of great magnitude, cannot, upon any reasonable grounds, be denied. The certain consequence of *neglected* gouty inflammation is, sooner or later, permanent debility and lameness; and, as we see by many melancholy examples, the patient, eventually, is often disqualified, by the crippled state of his limbs, from the degree of exercise which is equally essential to his health and to his comfort, even in such his limited state of enjoyment.

It had long since appeared to me, that a more gradual exhaustion of the inflammation, and a more soothing mode of effecting this than can be obtained by active cold, might be free from all the objections and disadvantages of the treatment in question. It seemed also not unreasonable to expect, that a moistened evaporating surface would procure very different sensations to the patient, from those which accompany the dry and burning heat of skin; and which, gouty inflammation so certainly produces.

I have now the satisfaction to state, that in about one hundred and thirty cases, I have made very free use, and with the best success, of a lotion composed of one part

of alcohol and three parts of *Mistura Camphoræ*; applying it to the affected part by means of linen rags, first rendered just agreeably luke-warm, by the addition of a sufficient quantity of boiling, or very hot water. In this manner, a prompt and convenient method is afforded of using the lotion, on the principles on which I recommend its adoption. The rapidity with which the alcohol alone would evaporate, is advantageously restrained by the dilution with the camphor mixture; and the warming it, by the addition of hot water, preserves it from that escape of the volatile parts, which the sudden heat of the fire would occasion. In using the lotion, if it be applied either hot or cold, the intention of the remedy is considerably frustrated; and I have observed, that from being made too warm, its operation has been injurious, rather than beneficial. If the temperature be measured by the thermometer, I may state that it ought not to be less than 75°, nor more than 85°. I consider, however, that the expression of *just agreeably lukewarm*, is a secure and sufficient direction to the patient. The linen compress, constantly kept wetted with the lotion, should consist of six or eight distinct folds, one laid upon another; and the slightest and coolest covering only should be used in addition. The effects of this lotion, when it has been attentively employed, have been most satisfactory, and have really answered my warmest expectations.

In the extensive opportunity which I have possessed of putting this local method of treatment fully to the test, I have only met with three instances in which I found that the lotion had been laid aside from disapprobation. In the one case, the nervous system of the patient was peculiarly sensible; and at the same time, the gouty inflammation shewed an unusual disposition to quick transference, and to fix severely in every part.

The lotion appeared to irritate rather than relieve, and was on this account discontinued. In the two other cases, it produced this last-mentioned effect only in a slight degree.

In the third example, the substituting of water for the camphor mixture in the dilution of the alcohol, was successful in preventing its causing any irritation. On a subsequent occasion, the first patient to whom I now allude, was again ill with acute gout; and I had the opportunity of making trial of the lotion by a little variation in the mode of its application; namely, the parts were freely sponged with it every hour, and then covered with a thin layer of flannel. In this manner it did not disagree, although I must confess that it was not so useful in its effects, as it commonly proves. From that time, however, I have been in the habit of directing this method of using the lotion, when the gouty inflammation is taking its leave, so that the constant application of it is no longer necessary. In such instances, I seldom find it necessary to use any particular covering to the parts, after the sponging process.

I have not in any one instance discovered, from the most unlimited free use of this lotion, that it has had the smallest tendency to produce retrocession, even with patients who have on other occasions suffered actual transference to internal parts, from the influence of exposure to cold.

Some who have formerly applied cold water according to Dr. Kinglake's plan, with more or less cause of alarm from its effects, have used this lotion without the least disagreement. I may add the same account, in the instance of a gentleman, who had, in a former fit, made use of a lotion of liquor ammon. acet. and alcohol quite cold. In a few hours after its first application, being seized with extraordinary giddiness and confusion of the head, and numb-

ness in the affected foot, he desisted from the treatment ; and with much terror of the result of his proceeding, he resorted to the readiest means of restoring the warmth of the limb.

In the cases to which I have just now alluded, the disagreement of the lotion was only local. No internal part was in the slightest degree affected. It will be readily imagined that its operation cannot prove equally beneficial to every patient, in a sensible and immediate manner ; or to every part under inflammation, in the same degree. The majority of persons, however, have praised even its immediate palliative effects in very strong language of satisfaction, and have called it soothing and delightful. It is an agreeable circumstance that its odour is pleasant and refreshing. In order to procure fully its good effects, the linen compresses should never be suffered to become dry ; and one set of them should be alternated with another, when the part is much heated, for the advantage of a cooler medium of application*.

I have here to observe, as a particular point of instruction, that when the gouty inflammation is so far overcome ; or, from peculiarity of constitution in the patient, it happens, that the application of the lotion by compresses as just directed, ceases to produce comfortable sensations of relief ; and, on the contrary, rather occasions chilliness, and possibly some numbness ; the method is to be changed, and the parts should be sponged with the lotion frequently, and left damp ; with or without subsequent covering of a light kind, as circumstances may direct.

* As a plan of great convenience for the application of the compresses, I usually direct that the foot of a large stocking be cut off from the remaining part, so that it is removed without difficulty for the purpose in question.

In order briefly to exemplify the effect of this evaporating treatment, and how much it is capable of relieving the painful symptoms, I quote the following report from two patients, given in their own language : and I can with truth add, that these statements are by no means too partially chosen from amongst my cases.

A gentleman was severely attacked with gout, which affected in succession the right hand, left foot, ankle, and knee, right foot, ankle, and knee, and lastly the right hand. He stated of the lotion, "I must unequivocally declare, that the benefit I received from the application was great beyond what I could have expected, and infinitely mitigated the usual throbbings consequent on gouty inflammation. My right hand first yielded to its influence, the pain and swelling gradually subsiding; and in a few days, all vestiges of either were removed in it. The other parts affected, nearly I believe in the order in which they had been attacked, exactly followed the hand, the swelling *gradually* disappearing, and leaving the joints unaffected by the least hardness."

Another gentleman much subject to violent paroxysms, was attacked with the usual severity in both feet. He began some purgative diuretic medicine late in the evening, and went to bed not having made use of the lotion. The pain became intense; the heat and throbbing were excessive; and it appeared to him that the parts "were tearing and separating in the sinews." At about six in the morning, having passed a sleepless night, and still in agony, he applied the lotion, and, in less than an hour, the pain so much subsided, that he fell into a refreshing sleep. He awoke free from pain. His medicine did not act until the middle of the day; and he praised the lotion as the surprising source of his com-

fort. Two days after, the hand was affected, and became equally relieved by the free application of the lotion.

It is necessary to observe, in reference to the narration of the first case more particularly, that the internal means of treatment were adopted with very active attention; and that the patient's expressions of approbation of the lotion, must be received in connexion with the effects of the still more important influence of internal medicine.

To continue my account of this remedy, I may state that in slight inflammation and pain, its powers of affording relief are soon discovered; and very sensibly acknowledged by the patient. Under symptoms of great suffering, although its influence is very beneficial, it is inadequate to procure ease; and it is in *opium* (the proper indications, as already stated, being fulfilled) that our superior and most necessary confidence must then be placed.

The good effects of the lotion are most remarkably seen, when gouty inflammation is situated in the superficial ligaments, tendinous structure, and common integuments; and it will be readily supposed, that in proportion as the seat of disease is remote from the surface, its operation will prove less efficacious: as, for example, when the deep-seated ligaments, and the bursæ mucosæ, particularly those belonging to the knee-joint, are the parts affected. In such cases, it has of late been my practice, chiefly, to confine myself to the use of the poultice made with the lotion; in general, directing it to be renewed thrice in the twenty-four hours. I have derived the best effects from this treatment. When the inflammation has been almost overcome, I have discontinued the application of the poultice, and resorted to the frequent sponging of the part with the lotion, in the manner lately described.

Upon the practice of effecting slow evaporation from the inflamed part by means of this lotion, I may be allowed to indulge in a few further observations.

The application being used at a heat only just tepid, it has a pleasing soothing influence; and its more active operation is obtained so gradually and securely, that I have never been able to lay any bad effects to its charge. Of its advantages, much might be said. In the same degree that we mitigate the local symptoms, we tend to relieve the general irritation of the constitution, and help to abridge both the violence and duration of the whole disease. It is also a fair subject of consideration, that present relief is not the only benefit that is gained. This lotion, independently of its powers in removing the inflammation as an evaporant, stimulates the absorbent vessels to useful action; and, further, acts as a tonic to the parts, greatly counteracting the ordinary debility which follows the inflammation of such important textures, as the gout affects; and which debility is especially produced by the peculiar action of this disease. In our successful treatment of the local inflammation, we accomplish much towards the preservation of the organization of the affected parts; and I do not hesitate to affirm, as a general position, that, with *timely* management of gout, a crippled state of limbs may with certainty be prevented.

With regard to the wandering character of gouty inflammation, I have not observed that this local treatment has increased the disposition of the disease to change its situation, unfavourably; and in no case, I must repeat, has any injury followed its most free employment. That the inflammation quitting one part, should immediately affect another, is an action of the disease not to be imputed to the influence of the lotion. This is the well

known character of gout; and we are to pursue our treatment with the same confidence that we began it.

In conclusion, I have further to assure my reader, that the interruption of the paroxysm, by the combined method of practice which I have endeavoured fully to lay down, has not been attended with the serious objection to which some remedies are liable, namely, its inducing an earlier return. On the contrary, however, I can with truth declare, that if the patient adhere steadily to the continued rules of treatment which it is the duty of the physician to prescribe, this first abridgment of the paroxysm by active practice, is at once the safe and direct method of striking at the root of the disease.

The occasional transgression of the patient, in taking too early an advantage of his amendment, or in abuse of rules after recovery, is not a fair impeachment of the propriety of the past treatment in the paroxysm.

The constitutional remedies, as already mentioned, having been employed to the necessary extent, and the evaporating treatment having fully produced its auxiliary effects, in the entire removal of all inflammation, we arrive at the next distinct stage which requires consideration.

THE CONVALESCENCE.

IN the early and least severe visits of the disease, the state of convalescence, as to constitutional management, may now and then require no other attention than what is prompted by good sense, and the consequent exercise of correct habits. In older and violent paroxysms, the

continued assistance of the physician is of the utmost importance.

Those tedious fits which run a neglected course of many weeks or months, and even those which have been correctly treated, but yet shew a strong disposition to relapse, may always be considered as radically depending on the presence of visceral disease of more or less magnitude. It is incumbent upon us to be convinced that the patient is really convalescent. It is not sufficient that our treatment has been active in the paroxysm. We have a great and two-fold duty remaining to be performed; the restoration of the healthy state of the digestive functions, and of due strength to the weakened limbs.

It occasionally happens, that the returning powers of the stomach demand, rather that the patient should exercise a degree of self-restraint, than that his appetite should be excited by medicine; and careful regimen, both as to diet and exercise, early hours, and a due regulation of the bowels, may constitute in such case all that is necessary to restore the general health. *Festina lenté* should be the constant motto, both of the physician and the patient. We should restrain the tendency of the vessels to acquire again the plethoric state; and keep in our recollection, that excess of blood in a weakened circulation will lead to many evils even worse than the gout; and the gout itself, may, or may not follow as a consequence. In judging of the propriety of employing bitters, or other stimulant tonics, simply, we should carefully ascertain that the secreting functions are become regularly healthy; and this inference must be drawn from the appearance of the tongue, the fæces and urine: and where it is necessary to make the judgment still

more accurate, the specific gravity of the morning urine may be examined from time to time.

It does occasionally happen in certain constitutions, that simple debility alone exists; and in that case, simple tonics only need be employed; the proper attention being at the same time directed to the regulation of the bowels. With this view, I have now and then derived advantage from the use of a chalybeate; but I have not proceeded with its use beyond a short period, so much is it the property of chalybeate medicine to excite the circulation unfavourably, and lead to a recurrence of gout. Where the habit is plethoric, the administration of steel should be considered, as a general rule, to be quite forbidden. In the instance of a gentleman of the nervous temperament, of a slight form and not plethoric, his debility being very great after a protracted fit of gout, the legs œdematous, and the urine albuminous, I prescribed the ammoniated iron and soda pills with great advantage; regulating the bowels with the pulvis aloës compositus, formed into pills with the decoction of the same, and a little soap.

The *tinctura ferri ammoniati* is a delicate preparation of iron, and may be conveniently given in water, cold or warm, in a dose from twenty to sixty drops, it being gradually increased.

Dr. Cullen, in speaking of tonics under the head of gout, remarks, "The most effectual medicine for strengthening the stomach is iron, which may be employed under various preparations; but to me, the best appears to be the rust in fine powder, which may be given in very large doses." I am well persuaded, that this very insoluble preparation of iron is the most objectionable of all its forms; and, as I have remarked in my

Analysis of the Chalybeate Water of Tunbridge Wells, the most soluble preparations of iron may certainly be esteemed the most active and useful. The tincture of muriated iron is one well deserving recommendation, though less delicate than the ammoniated tincture, and therefore not so generally admissible; but if suitable, it may be usefully joined with the cascarilla, or cusparia bark. And certainly, when from great debility of the constitution, the skin is so relaxed, that on the slightest exertions, perspirations, with faintness, take place; the muriated tincture may prove a very useful astringent tonic.

The case of a gentleman presents itself to my recollection, in which, from the error of leaving the paroxysm to its own spontaneous course, such a state of debility resulted, that for a considerable time most profuse perspirations took place at night, joined with great general debility and much œdema of the lower extremities. Here, the muriated tincture, with cascarilla, proved very useful. Advantage was also derived from sponging the skin very generally, night and morning, with vinegar and water, slightly tepid; local treatment was used for the recovery of the limbs; and the diet was made restorative, more especially by the use of solid than fluid nourishment.

It is important to observe, that a paroxysm of gout, correctly treated, never leads to the debility which is here described. It arises only out of the fruitless efforts of the constitution to get rid of the disease; and its consequent protraction to a period almost indefinite: or it may be the consequence of injudicious empirical treatment. It most commonly happens, that, although the patient be left by the paroxysm in a state of debility, there is that condition of the constitution which rather

requires the use of corrective and tonic medicine in combination, than of tonic astringents simply. It may be stated as a general rule of practice, that in the convalescent stage, our treatment should be such as is calculated to answer a part or all of the following intentions.

The stomach is to be restored to its proper energies; the liver to its healthy state of functions both as an organ materially concerned in the just balance of the circulating system, and as furnishing by its secreting action an important stimulus to the digestive process; the bowels are to be brought to their due and regular action; the kidneys to their share of duty in separating from the blood itself what is intended to be excrementitious; the skin to an equal and uniform office of insensible perspiration; the animal heat of the body to a proper standard, and an equal distribution. If these points be accomplished, the nervous system will require no other assistance by medicine; and equally useless and improper are stimulants given to the nerves, when the condition of the various functions just mentioned is neglected.

Sydenham, after many excellent observations on general rules of management, says, "It is clear, from what has been delivered, that whoever undertakes the cure of this disease, must endeavour to make a thorough change of his habit of body, and restore it to its former constitution, as far as age and other circumstances will permit."

It may be laid down as a very general axiom, that in every case, in which the constitution has been much under the influence of gont, the liver is more or less obstructed, and under error in its secreting functions; and also that such a condition of the digestive organs prevails, that, however the appetite may seem active at a favourite meal, the digestive process is imperfect, and the assimila-

lation of the chyle is not completed in a manner favourable to perfect health. This, therefore, is the period at which the skill of the physician is to be exercised with the greatest attention; and the patient is to consult the future welfare of his constitution, by the careful observance of all the rules which are laid down.

The particular method of treatment which the convalescent period may require, must obviously vary in its shades of difference, as much as every case itself varies in some of the particulars, of the age, constitution, temperament, and habits of the patient; and the peculiar individual features of complaint. And unless we make this discrimination in practice, we render ourselves truly empirics. It is with this provisional restriction that I now enter on my further details.

In the large majority of my cases, I have seldom found occasion to make use of bitters, or any description of tonic medicine; and I have on the other hand preferred to pursue a mild course of aperient and alterative medicine; in conjunction with a regulation of the diet and general regimen.

In those instances, in which, a languid state of stomach and deficient appetite do suggest the propriety of using a stomachic medicine, I can much recommend the following combination:

℞ Calumbæ radicis concisi ℥i ad ℥iss
 Cascarillæ corticis contusi ℥ii ad ℥ss
 Rhei radicis concisi ℥i ad ℥ii
 Cardamomi seminum (capsulis demptis) contrit. ℥ss
 ad ℥j
 Aquæ ferventis octarium dimidium.
 Macerâ per horas duas, et cola.

℞ Hujus infusi ʒxi ad ʒxv
 Tincturæ Aurantii ʒj
 Sodæ Carbonatis gr. x. ad gr. xv. M.
 Fiat haustus bis quotidie sumendus.

It is sometimes an advantage to increase the proportion of the carbonate of soda in this draught, and to direct a dessert spoonful of lemon juice to be mixed with it; so that, taken in effervescence, it becomes a more agreeable, and often a more useful medicine. When the palate of the patient is more to be studied, or the delicacy of the stomach require it, the rhubarb should be omitted, and pills, consisting of pulv. rhei, pulv. aloës compos. et sapon. dur. to be taken at bed time, may be substituted for it. As another form of tonic stomachic medicine, which I have found to agree particularly well with the gouty and dyspeptic stomach, I may mention the union of decoct. aloës compos. infus. gentian. compos. et mist. camph. with a moderate addition of spir. ammon. compos.; to be given once, twice, or thrice in the day, accordingly as the languid powers of the stomach and nervous system may require its use.

As an alterative pill, to be given with the view of exciting healthy secretions, and administered in a dose of five grains every other night, the pilula hydrargyri submur. compos. has appeared to me more decisive in its good effects than the pilula hydrargyri*.

* It may at first sight appear immaterial, whether corresponding doses of the pilula hydrargyri, pilula hydrarg. submur. compos. or hydrarg. oxydum ciner. be employed; but experience teaches us that each of these preparations possesses considerable

In those examples of convalescence, when an attention to the bilious secretion and the action of the bowels is alone required, it will often be sufficient, with regard to medicine, to administer the alterative and a purgative pill on alternate nights. This latter pill may consist of suitable proportions of gum. gambog. pulvis aloës compositus, et sapo durus. The patient, under this course of medicine, should receive particular injunction to avoid any careless exposure to wet and cold, at all times a matter of some hazard to a gouty person; and such caution is the more important, when we consider that the gout chooses that season of the year for inflicting its worst pains, when wet and cold almost constantly prevail. Still the necessary treatment must not be deferred; for I can scarcely conceive that any one, however circumstanced in occupation, can find a real difficulty in observing those moderate rules of care in clothing, and general means of security, which common prudence itself points out.

Some individuals have a peculiarity of constitution; and, also, the constitution of any individual may be in such a peculiar state; that the administration of a mercurial alterative, even though given occasionally only, and in a small dose, must be a matter of careful consideration. A very white appearance of the tongue, joined with much nervous irritation of the system, comprise a strong contra-indication to the use of mercurial medicine,

difference in its action. We see also that, in some individuals, one or other of these preparations will agree, when the other forms of the medicine have disappointed our expectations.

It is a circumstance always demanding careful consideration, that the quicksilver which is used in the preparation of the *pilula hydrargyri*, should be perfectly freed from lead or other contamination, by distillation.

as a general rule; and should make its continuance questionable, if arising as the evident or suspected result of its influence, when it has been employed.

In some cases it may meet the wishes of the practitioner to employ a vegetable tonic diluent, as the decoctum sarsaparillæ compositum; the effects of which were much praised by Sydenham; but on the particular form of restorative medicine, it would be both tedious and superfluous now to enlarge.

The peculiar kind of diet, which may be most appropriate, must, as with regard to medicine, be suitably varied for different persons; but to the prudent, a few plain rules will prove a sufficient guide. I think it, however, necessary to observe, that the general rule of moderation in diet at the convalescent period, which is so important a point of observation, is sometimes carried to an injudicious degree of abstinence; and much debility of the system becomes the consequence. I saw lately a striking instance of this result, in a gentleman, who pursued unnecessarily a course of low diet, when all the inflammatory action of the paroxysm had been perfectly removed. He brought on so much constitutional relaxation, that petechial spots (*purpura simplex*) appeared in various parts of the body, associated with œdema of the ankles, excessive languor and depression, and a disposition to perspire profusely on slight exertion. All the symptoms soon yielded to invigorating treatment.

While on the one hand, therefore, we avoid that repletion of vessels which too much indulgence would produce; we should, on the other, equally guard against incurring a state of inanition and debility, by an extreme restraint of regimen.

The use of ass's milk, as a mild nutritive diluent, is particularly deserving of recommendation, when it is our

object to pursue the restorative plan of treatment in the most efficacious manner. Half a pint should be taken every morning early; and, if convenient, should be repeated at night. It often has the property of acting gently as medicine both on the bowels and kidneys; and has a very favourable influence on the stomach itself*.

The advantages of a good air, and of a change of air and scene, with a suitable portion of exercise, are points of obvious importance towards the perfect establishment of health. As the present subject of general management will again come under our notice, in the consideration of *prophylactic regimen*, I proceed now to discuss the treatment of the weakened limbs.

The permanent œdema, and excessive debility, almost amounting to paralysis, which are sometimes consequent on the paroxysm, are more especially the effects of warmth and relaxing treatment, and may be prevented by correct proceeding; but present care cannot always repair the evil of former error; and it becomes sometimes a difficult task to invigorate the enfeebled limbs.

When, in addition to the ordinary debility of the limbs remaining after the perfect removal of inflammation, œdema takes place; the use of rollers, which may consist either of calico or flannel, according to the season of the year, is of important service. In any instance in which the veins are varicose, this treatment is indispensable. Also, when remarkable weakness of the ligaments prevails, so that the patient feels the want of support in

* Ass's milk was much extolled by Hippocrates, Celsus, and Pliny, in the cure of the gout.

the ankle joints, the bandage will be found a source of comfort and advantage.

I have sometimes seen, that, from nervous irritability, the patient at first objects to the sensations of restraint and pressure from a bandage; but a little resolution may cause these soon to be disregarded; the degree of tightness being duly regulated.

At this period, the patient should adopt the practice which I shall more particularly describe at a future page, of sponging the affected parts with salt water, rendered slightly tepid, every morning. The skin being wiped carefully dry, diligent friction with the hand, or flesh brush, should be continued, till a comfortable glow in the skin is produced. This method will supersede the necessity of immersing the feet in warm water, which is commonly practised from motives of cleanliness or comfort; but which I disapprove, as tending to increase debility.

If the paroxysm has, in all respects, been correctly treated, it seldom happens that additional remedies to this plan are required for the restoration of the limbs; but when, from any cause, an extraordinary degree of debility exists, so that there is great deficiency of energy both in the blood-vessels and nerves, together with the relaxation of the ligaments and muscles; as indicated by œdema, coldness, paleness of the skin, inability to bear the weight of the body, difficulty in walking, and an evident feebleness and shrunk condition in the muscular fibre; we must, in addition to the morning plan of sponging, apply, with a due share of friction, a stimulating liniment. The following formula I have found remarkably successful. It is to be made more or less exciting, as the individual case may require; this being a point deserving most particular attention.

R Tinct. lyttæ ʒss.

Linim. camph. compos.

—— saponis compos. āā ʒiss. M.

Fiat linimentum, quocum partes affectæ diligenter fricentur semel vel bis quotidie.

In urgent cases of the present description, the systematic method of friction, which I shall have to mention when speaking of chronic gout, will be found necessary to complete the plan of treatment; in order to hasten the recovery, and give a vigour to the limbs, which they may not probably have possessed for many previous months, or even years. In further illustration of the principles of practice which I have advanced, I shall now offer a series of cases, transcribed from the register which I kept of them. A few of these I shall state with minute detail; but others, I shall condense for the sake of my necessary limits; endeavouring at the same time, to retain such particulars as appear the most important and instructive; and, for the more complete elucidation of my subject, annexing to the narrative occasional commentary remarks.

CASE I.

W. W. a coachman, aged 60; tall, of stout make, was originally robust and vigorous, but now his muscles, and particularly those of the lower limbs, are small and weak; has a circular chest; of sanguineo-nervous temperament, plethoric habit, and irritable disposition. He is occasionally nephritic, and once, a few years ago, suffered very severely with retention of urine and severe spasms of the bladder. He is frequently, of late years, dyspep-

tic; his tongue is always more or less furred; and his nose and face exhibit the *gutta rosacea** in a great degree. With such exceptions, has never had any other disease than gout. This disease quite unknown in his family. When young, he was fond of athletic exercise. Having always lived in good places as coachman, and for a part of his life kept an inn, he has had very constantly the means of indulging in free living. He describes himself "as a moderate eater, and for many years past, particularly, not indulging freely in liquors;" but acknowledges, that he has regularly been in the habit of taking porter and mixed spirits. When he has occasionally exceeded in a glass of raw spirits, has experienced much consequent heat of stomach, a feverish state generally, and now and then a paroxysm has followed such irregularity. His first attack was at the age of thirty. He says he was in full health at the time. He had been bathing in the sea, and, when returning to shore, contused his foot severely on some rock. He reached home without much inconvenience, and went to bed well; but in the middle of the night awoke with pain in the great toe of this foot; it was decided gout, and continued a fortnight. It returned in two years after, first in the same foot, and then in the great toe of the other foot. This second fit lasted a month. He soon became a confirmed gouty subject, seldom escaping a whole year; more commonly affected twice a year; but his chief fits were in the beginning of spring, and were remarkably periodical in their return. He imputes his attacks most commonly to wet and cold; but some of the severest have followed blows and strains; and the most

* *Acne rosacea*.—Bateman.

violent fit which he ever had, succeeded immediately to a bad strain of the ankle. On that occasion other parts became affected, but in some of the fits excited by local injury, the disease has confined itself to the part so inflamed. He wears flannel socks, and commonly perspires much in the feet; and he remarks, that a little previous to an attack, this perspiration is often suspended. His usual premonitory symptoms are, depression of spirits, much listlessness and yawning; flying cramps, especially affecting him at night before his first sleep; a cough, with nauseous expectoration. A fit usually makes its invasion about one or two A. M.; but this happens with some exceptions, which are more remarkable when local injury has been the exciting cause. Some of his fits have continued twelve or fourteen weeks. Three years ago he was bled at Christmas (from choice only, being then as well as usual), and entirely escaped gout the following year.

Progressively he has had gout in every part of each foot, in the knees, hands, and elbows. Has *uric-concretions* at the outer side of each heel, from which, painful ulcerations have now and then formed. A little finger, which was inflamed in his last fit, exhibits the whitish appearance of incipient concretion. Has large distended veins, and those of the legs are varicose. The bursæ mucosæ, both in feet and hands, are either much puffed, or in the state of *ganglion*; the tendons of the fingers are rigid from their distended and thickened sheaths; and from such change of surrounding structure, the phalanges have the appearance of bony enlargement. He has long suffered from constant lameness and uneasiness, both in the feet and hands.

Intending to make this patient the subject of some comparative examinations, in regard to the secreting

action of the kidneys, I examined his urine on the 3rd of May 1815, when he was not complaining. Having already detailed the particulars of this investigation, at p. 120, I shall, in this journal, for the most part, only state the specific gravity of the urine on the relative days, and the proportion of phosphoric acid. The morning urine of this day was of specific gravity 1009·4; and four ounces afforded of phosphoric acid ·8 grain.

On the night of the 3rd of May, he was exposed on his box, a considerable time, to wet and cold; and remained several hours in wet clothes. On the 4th, although he went about with difficulty, the invasion of a paroxysm had begun. He was chilly, feverish, and his right arm was painful. On the 5th (Friday) I visited him, and found him with severe gout in his elbow and hand. In each part there was swelling and some pitting; but most at the back of the hand. The skin was vividly red. The usual sensations of acute pricking, tearing pain, burning, throbbing, and weight, were present in an intense degree. He experienced the greatest degree of suffering from the elbow; and the slightest motion of the arm was infinitely distressing. To use his own words, he felt also, "spasms in the limb, and hot glows all over; qualmish and thirsty; sudden perspirations, as suddenly leaving him; heat and spasms in the stomach; spirits very low." Pulse frequent, and rather full; skin hot; tongue much furred; much thirst; bowels costive; the urine passed with frequency and irritation, and in small quantities at a time, as is usual with him under gout. It deposits, on cooling, a copious mucous and lateritious sediment. Its specific gravity 1·0201. Four ounces afforded of phosphoric acid 5·36 grains.

With a view to form some opinion of the comparative quantity of urea in this urine, with what should

be found afterwards, at the restoration of the patient's health; and also to judge of its relation in this respect to the urine of other persons in health, I evaporated a given portion to one-third of its bulk, and added concentrated nitric acid freely (about an eighth part). Crystals consisting of urea and the nitric acid, were produced in much greater abundance, than I have ever found in a similar experiment with healthy urine.

From the persuasion of a fellow-coachman, he was taking a quack medicine, *Bateman's drops*; and on this account I did not see him again until Tuesday, May 9th, when I began the following journal:

He reports that the drops have occasionally procured him sleep, and caused much perspiration; but he felt heated by them. Bowels open daily, the last two days. Last night was the worst which he has had; quite sleepless. At five this morning, his agony was very great. Pulse now 84, and full; skin hot; much thirst; no appetite; tongue much covered with brownish white fur. Very restless and nervous; sore to pressure at the stomach, and in the right hypochondriac region. Bladder very irritable; passes much less urine than natural, with frequency and some difficulty. Relates that at all times his bladder is rather irritable; yet in general he has a free and easy stream. I had not the opportunity of seeing this day's urine; but he states, that its appearance was similar to that of the 5th. The catarrhal cough, with which his gout commenced, is much abated.

The elbow is improved since last report; the hand still much affected; thumb the most inflamed part; middle finger much inflamed; vivid redness; no pitting. The whole of the external part of the left foot much inflamed; vivid redness; much bursal distension around the ankle joint, but no pitting; veins very full, and par-

ticularly all those ramifying from the inflamed part*. Two of the small toes red and swollen; much subsultus tendinum. He was seized first around the heel, with numbness, and sudden entire loss of strength in the foot. Last night severe throbbing; sense of great weight; foot feeling as if not belonging to the leg; tightness; burning heat. The following treatment was adopted:

R Hydr. submur. gr. iv.
 Pulv. antimon. gr. ij.
 Extr. colocynth. comp. gr. x.
 Saponis duri, gr. iij.—M. fiat pilulæ iij. statim
 sumendæ.

R Magnes. Div.
 Sulph. magnes. ʒvj.
 Aquæ menthæ viridis, ʒv.
 Aceti colchici,
 Syrupi croci aa ʒss.—M. Hujus misturæ capiat
 partem quartam, 6tis vel 8 vis horis, prout alvus soluta
 fuerit.

The *Diet* to be limited to gruel, tea, and bread, barley water, or rennet whey. The constant application of the evaporating lotion to the affected parts, according to the formula, p. 254; a sufficient portion of hot water being added to it, to render it just agreeably tepid.

Wednesday Evening, the 10th, 9 o'Clock.—The medicines have agreed perfectly. The bowels copiously

* The veins of the right leg, at the lower part, very full in the line of the inflammation; and altogether fuller than in the other leg. No difference in the veins of either upper arm. In the affected one, the fulness begins at the wrist.

affected twice, at twelve last night, and at eight this morning; the evacuations very green and foul. The pain was very soon much mitigated by the lotion. As soon as the bowels yielded, his skin cooled; he became easy, and passed a comfortable night. He has not perspired. The urine *first* passed in the morning, of an amber colour, with slight pink sediment, and less mucus than yesterday; its specific gravity 1·0099. Four ounces afforded of phosphoric acid 3·17 grs.

He continued easy and comfortable till five this afternoon, when the paroxysm was renewed in the right ankle; and he is now suffering *such agony*, that he repeats an expression which he has before made use of during the sufferings of this attack, "*that he would be thankful to any one to shoot him.*" The inflammation in the newly-affected foot is deep seated, as the skin is scarcely discoloured; much bursal and thecal distension; pain shoots through both ankles; violent throbbing and burning; sense of immense weight, and extreme tightness; has frequent spasms, pricking and shooting; subsultus tendinum, and convulsive state of gastrocnemii muscles; receives only momentary intervals of relief. Pulse 72, rather full, but not inflammatory; excessive thirst; skin temperate. The application of the lotion has been neglected this afternoon: now to be used freely and constantly. The bowels having been freely emptied, and constitutional inflammatory irritation being much abated, the indication for direct soothing treatment appeared clearly manifested. I directed the following pills:

R Pulv. opii crudi gr. xx.

Pulv. antimonialis gr. x.

Confect. rosæ canin. q. s.—M. et divide in pilulas xx, quarum capiat iii. statim; et repetatur i. omni hora, donec dolor sublevetur.

Thursday, May 11th, 1 P. M.—He felt immediate comfort from the lotion; but the great relief, which he very soon obtained, was certainly due to the opium. Even in a few minutes he was a little composed; in twenty minutes, a numbness, rather pleasing, of the affected parts, with an abatement of heat, throbbing, &c. took place; and the convulsive action of the muscles, and the subsultus tendinum ceased in about half an hour. He did not find it necessary to take a second dose of the opiate. The night was passed in tranquillity and dozing. Now has slight perspiration, and no feverish heat of skin; thirst is abated; tongue rather more furred than yesterday, but he has some appetite; head rather aching, yet has not experienced any confusion; an excellent pulse at 76. He lies upon the bed; and his apartment is kept temperate. Uric deposit appears to be taking place immediately under the cuticle in two of the fingers. Both feet almost easy, and but little discolouration remaining. Bursal distension lessened; there is some pitting of the integuments; fulness of veins much removed; temperature at the right ankle, at present most the seat of complaint, 97°. Has taken the mixture regularly every six hours; it has acted twice freely; the fæces very foul, with much acrid bile, very hot to his sensations, less green than before. Much diuretic effect also from the medicine.

The urine is passed both more copiously and comfortably than at any period since his attack. It deposits dirty brick-coloured sediment, some reddish crystals, and much mucus. Its specific gravity before filtration, 1.0242; after, 1.024. Four ounces afforded of phosphoric acid 3.88 grains.

To continue the mixture every six hours; the opium pill as the pain may require; and, every night, five grains

of pilul. hydr. submur. compos. The lotion constantly. Diet as before, fluid and wholly unstimulating; may take bread and milk, or milk porridge, at discretion.

Friday, 12th.—Much better. Has not had severe pain; slight accession of symptoms about nine last evening, continuing till twelve; but he did not suffer enough to be induced to take the opium; a small dose of which, however, I should have approved. His night was rather restless and feverish. The gout, about four A.M. affected one middle finger, but not severely. I find it swollen, hot, and partly red. All inflammation removed from the left foot; a little remains in the right ankle, with slight redness; very little pitting in either foot; little toe in right foot still red and painful; he expresses great ease and benefit from the lotion; has taken the mixture at the regular intervals of six hours. Pulse natural; spirits improved; surface temperate; tongue less furred; nose, which has been *glowing* red, now getting pale; the heat in the stomach and bowels, of which he has been complaining, is removed. The discharges less heating, but still foul and of yellowish-green colour; no preternatural thirst; no nausea; moderate appetite. The urine has a slight mucous cloud, and no other sediment. Specific gravity 1.012. Four ounces afforded of phosphoric acid 1.02 grains. To continue the treatment.

Saturday, 13th.—Going on most favourably; passed a good night; no return of pain; every part almost free from inflammation; and the parts most affected yesterday, to-day bear considerable pressure; feels uneasiness through the foot at the little toe, and there is a slight spot of redness on the upper surface at this part; suffers most tenderness from the old concretions near the heel, but derives much relief from a simple bread poultice

begins to walk tolerably. The medicines have produced their usual effect. The urine similar in appearance to that of yesterday, with the addition of a few minute crystals; specific gravity 1.0105. Four ounces afforded of phosphoric acid 1.47 grains.

Sunday, 14th.—Did not pass so good a night on account of pain in the last phalanx of the little toe, and along its metacarpal bone; the spot of redness before described continues; the pain was not sufficient to render the opiate pill necessary; has slight spasms from the foot to the knee; pitting of each ankle in a small degree, with paleness of skin; pulse and skin natural; tongue still rather furred; three stools in the last twenty-four hours, much improved in appearance, thin and of a light yellow colour, with but little accompanying heat. Urine abundant, of a light straw colour, with a fine flocculent mucous sediment, and a small portion of minute uric crystals; specific gravity 1.0106. Four ounces afforded of phosphoric acid 3.47 grains.

Has lived hitherto on diluents; but on this day he has eaten bread-pudding, with appetite and relish. The last two nights, has taken the alterative pill at bed-time, till last night; has continued the lotion to all the parts yet affected with any tenderness.

Tuesday, 16th.—In all respects doing well. The urine of specific gravity 1.0085. I discontinued my experiments for the present. The mixture now to be taken twice a day, with ℥j. only of sulphate of magnesia as a dose, and the other ingredients as before; the alterative pill on alternate nights. All inflammation being removed, apply rollers from the foot to the knee; sponge the extremities with water containing salt dissolved, and the chill removed so as to feel pleasant; wiping the skin afterwards perfectly dry, and using diligent friction.

He expresses benefit from using a horse-hair brush through the stocking. Much caution to be used in diet; a little animal food of light digestion, once a day, with mashed potatoe, and some fresh table beer. The weather being fine, to be much in the air, and to use gentle walking exercise.

I may here observe, that each day's urine has reddened litmus; has precipitated reddish-white or brownish-white flakes, more or less abundantly with oxy-muriate of mercury; brownish flakes with infusion of galls*;

* I have invariably found, that the first urine of the morning, such even as is secreted in good health, *immediately* becomes turbid with infusion of galls, and with the solutions of oxy-muriate of mercury and of alum. The precipitate which subsides, is more or less abundant, and dense and coloured, according to the proportion of saline and animal matter present, and the consequent high specific gravity of the urine. The galls occasion the darkest coloured precipitate. That from the oxy-muriate of mercury is usually most dense, and is often coloured of a reddish hue. It has the appearance of pus mixed with water, and a little coloured. The precipitate produced by the solution of alum is more white and flocculent, and contains also uric acid deposited in crystals.

Mr. Cruickshank has erroneously described the effect of the oxy-muriate of mercury, as being an indication of disease, in the following words (Rollo on Diabetes, 2d edition, p. 443): "The corrosive muriate of mercury is a very useful re-agent, as it has no immediate effect upon recent healthy urine; but in every case of increased action of vessels, more particularly of the inflammatory kind, a greater or less milkiness, and a whitish precipitate is instantly produced." This incorrect statement is copied by Berzelius (View of Animal Chem., p. 98). That the effect is *greater* in urine of high specific gravity, and therefore in certain states of disease, I have already granted; but the very material

has not suffered any change of transparency from nitric acid ; or, from the application of heat.

Sunday, May 21.—Has daily improved in health ; appetite good, and digestion comfortable ; tongue has still a whitish fur, but it lessens. Walks with much more strength ; ankles swell a little after exercise, but the swelling is on the decline ; the veins have acquired a healthy appearance, and are even appearing in a better state than before this attack. The bursæ are smaller. The temperature at the ankle, which in the paroxysm was 97° , is now 86.5 . The bowels have been kept in a free state by the mixture, suitably abated in the frequency of dose. The fæces, though much improved, are not yet healthy. The urine is quite healthy. Discontinue the mixture ; take the alterative pill every third night ; and the following purgative pill occasionally :

R Pulv. alões compos. ζj .

Pulv. antimon. gr. v.

Saponis duri gr. x.

Decoct. alões comp. p. q. s. M.

Fiant pilulæ xx. quarum capiat ii. vel iii. h. s. albo astrictâ.

Disliking exceedingly the sensation of the rollers, he may be allowed to discontinue them ; but must regularly pursue the plan of sponging and friction in the morning. Having now to return to his work, may take

mis-statement to which I have referred, deserves, I think, to be pointed out. Dr. Blackall (on Dropsies, p. 15) speaks in doubtful terms of the effect of the infusion of galls, as to the nature and extent of its indication in disease, or, “ in some circumstances of *apparent* health.” Here, the powers of this re-agent also are not correctly conceived.

half a pint of porter at dinner ; but on no account to have animal food more than once a day, and that of the most digestible kind. Did this patient's station of life permit it, I should much prefer allowing him two or three glasses of good wine daily, rather than porter.

Saturday, May 27.—Convalescence most favourable. Tongue almost clean; appetite and spirits good; healthy strong pulse. Urine passed very freely, but the long acquired irritability of the bladder remains in a considerable degree; bowels open each day; discharge not yet appearing quite healthy; lameness almost removed; some tenderness at the seat of the uric concretion at the side of the heel. In order to excite absorption of the uric matter, apply freely night and morning, by means of friction, the following lotion :

R Liquoris potassæ

Misturæ amygdalæ aā ʒij. M.—Fiat lotio.

To wear over the part, a soap plaster spread on soft leather. Continue each pill a fortnight, as before directed; and then lay aside all medicine, except the occasional employment of the purgative pill. For several days past, he has taken a pint of porter or upwards. On no account to exceed a pint and a half in the day; and to eat animal food once a day only. To guard very carefully against wet. To sponge the feet and ankles thoroughly every morning, in the manner described at p. 270*, as a constant habit; wiping them carefully dry; using subsequent friction till a comfortable glow in the skin is produced; and finishing the process with one foot, before the other is begun.

*—See also under the head of Prophylactic Regimen.

July 4th.—He assures me that he has pursued attentively all the regulations prescribed. He is now restored to good health. Tongue clean; countenance clearer than I remember to have seen it; appetite and digestion good, and alimentary secretions healthy; spirits cheerful; limbs much stronger than they have been for years; the uric matter very much absorbed, and the parts where it was deposited almost free from tenderness.

The morning urine of this day of amber colour, with a slight mucous cloud; specific gravity, 1.0172. Four ounces afforded, of phosphoric acid, 2.8 grains.

I now made a comparative experiment with reference to the proportions of urea and uric acid, and found that these principles were also, comparatively, in much smaller quantity than had been afforded by the urine voided during the paroxysm.

June 1816.—This patient has entirely escaped gout since the last report, notwithstanding he has been regularly employed as coachman, and consequently exposed to the changes of weather, both by day and night. He has, also, during the whole period, enjoyed unusual health and spirits. An example of a habit more completely gouty, could scarcely be selected. Under the disadvantages of exposure, which are attached to his situation in life, and of occasional inattention to regimen, it cannot be presumed that the gout will not return; but it is shewn, both that the treatment in his last violent paroxysm (similar in its degree to his former fits, which neglected, or slightly treated, always ran a course of twelve or fourteen weeks) was speedily successful; and that it has not, by shortening the disease, or by a quick removal of its most urgent symptoms, induced any disposition to a more early return; but quite the reverse, as the present date abundantly shews.

June 1817.—He makes at this period a very favourable report of the improved state of his constitution. In the course of the last winter and spring, he had two or three very slight attacks, in consequence of continued exposure to wet and cold, and living in a damp situation; but they proved tractable and of short duration. He expresses himself now to feel in good health and spirits.

September 19th, 1818.—Has been free from gout since the last date, with the exception of one attack, which continued only a few days, and readily yielded to the former plan of treatment. Has also been free from gravel. He has carefully observed my instructions to regulate the bowels; to use the daily washing of the head; and, with as much regularity as in his power, the sponging of the feet with salt water. He has, at three or four different periods, taken the mixture (p. 276) once a day for about a week, with an alterative pill each other night. Hence, when every circumstance is considered, this must surely be regarded as a strong example, both of the controul obtained over the paroxysm itself, and of the success of prophylactic treatment.

CASE II.

April 1815.—J. C. aged 38, a plumber and glazier; short in stature, has a circular chest, is thin, but, according to his statement, was formerly corpulent, both before and some time after he became gouty; of sallow complexion; of nervous temperament; has great sensibility of nerves, but is apparently of mild disposition; perspires easily on exertion; suffers much from the cold of the winter, and particularly in the feet; for two years past

subject to copious hæmorrhoidal discharge; and says, that if this occur when the paroxysm of gout is present, he experiences from it immediate and very considerable relief; previously to the flux, has local heat, weight, and fulness; habitual state of bowels costive, since he has been gouty; before this, they were regular. Has pursued his trade in London nineteen years. Has always drunk porter and gin daily, and formerly in excess; also, till of late, that he has been debilitated, ate much of animal food. Not subject to other diseases. Gout unknown in his family. He was first attacked nine years ago, at the latter part of spring, when the weather was hot, in the great toe of one foot only. The fit continued ten days. The second attack was at Christmas following; first in the same part as before; next, on its subsiding there, in the great toe of the other foot. This fit continued three weeks. In subsequent attacks, the knees and hands have been severely affected. Exposure to cold the most common exciting cause. The time of the invasion of a paroxysm generally about twelve or one in the morning; sometimes, when having gone to bed with the usual feelings of good health. When he has had precursory symptoms, they are the following: unusual fulness of the veins; prickings at the ends of the fingers, if the hands be threatened with the disease; but if the feet, flying pains in them; feels getting lame; restless at night, with little and unrefreshing sleep; depressed spirits; loss of appetite; suffers cramps also severely, at the side of the hand, and in the fingers themselves, if the hand be about to become the seat of gout; in the calf of the leg, if the foot; commonly has rigors on the invasion of the fit, and sometimes afterwards, very irregularly; always suffers most pain in the night; most in an old part again affected. In some fits, the pain has continued

a fortnight with little intermission, even in the day; the longest duration of a fit, eight weeks; the shortest, ten days; the longest interval, fourteen months; the shortest, three weeks or a month.

The present paroxysm began on Monday, April 3d. I saw him first on the 15th. He relates that he was exposed for several hours on the Monday, to a cold wind and rain, in cleaning windows; went to bed, however, without complaint; was awoke between one and two in the morning with pain in the middle finger of the right hand, which rapidly inflamed and swelled. The whole hand was also soon affected. Gout continued in these parts for eight days; then seized the left hand; first in the middle joint of the fore finger, and subsequently in the metacarpal joint of the same finger, the inflammation spreading also to the back of the hand. Describes his pain as most excruciating, and sometimes to the degree of producing delirium. To use his own language, "he screams from agony; the affected parts feel as if in the hottest fire; sweat issues from the ends of the fingers in the pain; often feels as if dogs were gnawing the flesh from the bones; violent throbbing; sometimes it seems as if pins and needles had been forced in under the nails, and were again forcibly drawn out of the flesh; heavy as if a ton weight were hung to the foot." He says that this last sensation, though very strong in the hand, when it is the seat of complaint, is still more violent in the foot; does not speak of the tightness which is so often complained of in gout.

At present (15th), the wrist and back of the left hand, and middle and metacarpal joint of the same, are the most affected parts. An abscess has formed over the middle joint of the finger externally. Part of the skin covering the matter, is black from effused venous blood;

part exhibits the appearance of white chalk-like spots: The abscess was opened, and a considerable quantity of white curdy puss was discharged*.

The inflamed parts are vividly red †; the back of the hand pits much on pressure; the adjacent veins, and particularly those in the line of the inflammation, are in the fullest state of distension. The great toe of each foot is much enlarged from the swollen bursæ; the skin faintly red, and there is much bursal distension at the external part of each ankle; suffers great uneasiness in walking, from the weakness of the ankle joints; and particularly on uneven ground, either foot sometimes turning suddenly with severe pain.

Pulse 104, full, and rather throbbing; irritable, with an unequal beat; tongue moist, and not very foul. Some days ago, vomited sour mucous matter, of a greenish yellow colour; is liable to this in the gout. Has some tenderness in the right hypochondrium on pressure, but none at the epigastric region; not costive; has a foul state of bowels, much slimy, clay-coloured discharge from them; the bladder very irritable, urine being passed frequently, and of a deep orange colour, but without any sediment, except a slight mucous cloud. That of the morning was found to possess the following properties: Its specific gravity 1.014; it reddened litmus in the usual degree; was rendered slightly milky by heat; and on cooling deposited an albuminous precipitation, appearing in whitish coalescing flakes. Four ounces afforded of-phosphoric acid 1.02 gr. (see Experiment v. p. 120). In the treatment of the case, the pills of calomel, colocynth, &c. (p. 185) containing of calomel three

* See Appendix to the Case.

† See Exp. p. 149.

grains, with two of antimonial powder, were directed immediately; the draught (p. 186) containing magnes. sulph. ζ iss. acet. colch. ζ j. &c. every six hours; the evaporant lotion constantly to the inflamed parts; and a bread poultice to the seat of abscess. The diet, gruel, tea, and bread.

16th.—Much better. Has taken the pills and three doses of the draught, without nausea; two copious evacuations, foul and dark, watery and hot. Urine much more abundant, and passed with less frequency and irritation than yesterday; its specific gravity 1.0105; affected as yesterday by heat. Pulse 84, with diminished irritability; has passed a good night, sleeping comfortably from ten to six without the assistance of an opiate. Previously, his nights have been sleepless, painful, and most distressing. Found great relief from the poultice, and assures me that the lotion gave ease to the inflamed parts in an hour. The appearance of the knuckle affected with abscess much improved; skin now of pale red; œdema lessened; finger less swollen, and can bend it a little. Still much tremor of muscles; often has palpitation, and is indeed altogether in a very nervous state. Continue all the treatment; add light broth and bread to the diet.

17th.—Feels much amendment. Had rigors yesterday afternoon, with a sensation of cold water down the back, not followed by heat of skin; still has tremors; complains of thirst and want of appetite; pulse 72, more equal and natural; tongue moist, but furred. Gout exceedingly relieved in the parts yesterday affected. The little finger of the same hand has inflamed suddenly within the last two hours; the skin is vividly red, burns and throbs. No further apparent secretion in the abscess. The skin appears as if inflamed from a scald; some vivid

redness remains. He mentions that three days ago he observed a red line beginning from the abscess, and passing round the elbow up to the shoulder, with much tenderness in the whole course. Was rather faint and weak on rising this morning. Has not passed urine so abundantly in the last, as in the preceding twenty-four hours, but with ease and freedom. It was affected as before by heat, and I now observed that it became turbid at 180°. The draught has acted freely four times; the stools hot and watery, and without griping: yesterday they were almost black; this day rather lighter. Repeat the pills, and all the other remedies.

18th.—Much improved. Has passed a good night; pulse natural; scarcely any redness remaining in any part; can move the finger having the abscess, without pain; its appearance improved; it bears some pressure; the swelling of the hand almost removed; expresses great satisfaction from the lotion. Free from rigors; appetite returns. The bowels have continued to be much affected; the discharges becoming natural. The urine still flaky from heat, but in a less degree. Continue the draught twice a day, and take five grains of the compound calomel pill each other night. Dress the sore with simple dressing. To increase the diet gradually, but to observe a correct regimen. To sponge the limbs and use subsequent friction, on the plan before mentioned, every morning.

June 20th.—He reports that he continued the medicines about a week. He rapidly recovered after the last report, and has remained well; feeling better, he assures me, than for two or three years past. This morning's urine possesses every healthy character, and is not affected by heat, or nitric acid. Its specific gravity 1.0137. Four ounces afforded, of phosphoric acid .91 gr. (See

EXP. vi. p. 120.) A small speck of uric concretion appears under the skin ; but the part is free from tenderness, and the use of the finger is perfect.

In addition to the injunction of a careful regimen, and the morning practice of sponging and friction, he is desired to regulate his bowels by means of pills, containing pulv. aloes comp. pilul. hydrarg. gum. gambog. and sapon. dur. prescribed in suitable proportions.

*Appendix of the Chemical Examination of the Matter
from the Abscess.*

Having collected a sufficient portion of this curdy pus, and exposed it to dry spontaneously, I submitted it to chemical examination*, with the following results :

This matter, exhibiting the appearance of *chalk-like* particles intermixed with pus, seems to be a mixture of albumen and urate of soda. Before the blow-pipe it decrepitates slightly like albumen, emitting an ammoniacal odour, and burning with flame. When urged by increased heat, the white residue fuses like carbonated alkali, and, like it, has a powerful effect on turmeric paper. The residue, exposed to the influence of a damp atmosphere, did not deliquesce ; whence it may be concluded, that the alkali was soda.

Like urate of soda, on being treated in the usual manner with nitric acid, it exhibited distinctly the rose hue.

March 1816.—This patient has continued perfectly

* My friend, Dr. John Davy, was present, and obligingly joined in the experiments.

free from gout, and has not experienced any interruption of good health. He has, however, lately relapsed into irregularity of habits; and consequently his future security will be very precarious.

June 1817.—No opportunity of discovering his present place of abode; nor of learning his subsequent history.

CASE III.

J. W. aged 42; of small stature, chest circular; thin; of nervous temperament; of active habits, walking much daily; has habitually drunk porter and spirits, but he says never in excess, and eats moderately. His mother was severely gouty, and died at the age of 46; he says, from the gout. He is not subject to other diseases. Was first attacked at the age of twenty-eight in one great toe only, from continued exposure to cold on a snowy night. This fit was of short duration. Second attack in seven months after, and in the same part only. The third fit about July in the succeeding year, affected the following parts in succession: the great toe of the same foot as before, each hand, one elbow. Continued to have gout annually, and often twice in the year; and in one year, after the free use of the eau medicinale, four times in the year. This medicine gave him present ease, but he felt, long afterwards, extremely nervous, and local weakness in a greater degree than he had ever before experienced. His bowels were left costive, and his skin appeared yellow and sallow for many months. Some of his fits have lasted twelve or fourteen weeks. One attack, a year since, affected various parts in the following order; right foot, right elbow, right hand, left elbow,

left hand, left foot, and also both knees. The diaphragm was frequently affected with spasm. His most remarkable premonitory symptoms are, cramps in the muscles of the lower limbs, affecting him severely as he is getting his first sleep, for several days before the attack; and sometimes for a week before, he has much scalding of water and sense of strangury. Also is dyspeptic, with great depression of spirits. The present fit (Dec. 1813) in which I saw him, was excited by cold and wet from walking in thin shoes on snowy ground. As is usual to him, he was attacked soon after midnight. The fit began three days before my visit, which was in the evening. I found the gouty inflammation just abating in one foot; and attacking one hand, which was inflamed and swollen, and in exquisite torture. His bowels were costive, and his urine deposited much pink sediment. I filtered it, and found that it coagulated with considerable firmness from heat at 180° ; and gave a very dense whitish precipitate with nitric acid*. I prescribed the purgative pills

* Dr. Wells observes (Trans. of a Society for the Improvement of Medical and Chirurgical Knowledge, p. 208), "I never, but in two instances, saw a pink-coloured sediment in urine which contained the least quantity of serum." In addition to the phenomenon in my present cases, I have met with it in a female who had ascites. The pink sediment was considerable, and the urine was strongly affected, both by heat and nitric acid.

The *precipitation* which is produced by nitric acid, or other re-agents, is commonly but very improperly expressed, as that of *coagulation*; a term which relates only to the agency of heat. The error in question was first pointed out by Dr. Bostock, in his interesting paper on the "Nature and Analysis of Animal Fluids."—Medical and Chirurg. Trans. vol. iv. p. 56.

and draught already mentioned in the two preceding cases, and the use of the lotion. This patient lived at a considerable distance in the country, and I did not see him again; but in a few days after I received a circumstantial account of his progress. The bowels were freely purged in three hours, and he was immediately relieved, so that comfortable sleep followed. The fæces were dark and slimy. The lotion proved very comfortable, and assisted much in giving ease. He has not required opium. The draught has been continued regularly, twice or thrice a day, with much purgative and diuretic effect. The urine still deposits pink sediment, and is in a slight degree affected by heat and nitric acid. A week after, the report was quite satisfactory. Appetite, sleep, and spirits returned; fæces of healthy appearance; the urine clear, and no longer affected by heat or acid. From the severity of this attack, he believes, from former experience, that without assistance, it would have continued two or three months, and attacked various parts. From his long fits, he has always experienced much constitutional and local debility for many weeks after their going off. Now his strength and energies return daily.

The only circumstance on which I shall further remark in this case, is, the albuminous state of the urine, which was more remarkable than in the preceding case. This phenomenon has been more particularly presented to our attention than before, by Dr. Blackall, in his valuable publication on Dropsy*. In thirty gouty cases, in

* Dr. Wells, in the paper to which I have already referred, read to the Society June 1811, has considered the subject of serous urine in a truly elaborate manner, and has presented an

which I have examined the urine with a view to this circumstance, I have found it albuminous in seven. I now offer the following additional examples :

CASE IV.

J. M. a strong man ; of sanguineo-nervous temperament ; has gout at present in the feet, but the paroxysm is abating ; slight œdematous swelling remaining ; pulse natural ; nerves irritable ; tongue furred ; bowels costive ; urine not abundant ; of specific gravity 1·0196 ; of orange colour, depositing much pink sediment ; filtered and heated it became flaky ; and with nitric acid also afforded a copious flaky precipitate. Two doses of the pills, with calomel, colocynth, &c. and the daily employment of the draught, p. 186, soon restored the health of the patient ; and even in a few days, the urine, when examined, was found to possess every healthy character.

CASE V.

E. L. of middle age, and originally of strong constitution ; of nervous temperament ; has lately had a severe paroxysm of gout, in both feet and one knee. The feet are very œdematous ; the skin is pale. The pulse is rather weak ; he is extremely nervous, and complains of

extensive series of examples.—Mr. Cruickshank, in the second edition of Dr. Rollo's Treatise on Diabetes, in 1798, appears to have been the first author who gave notice of the phenomenon in question.

much lassitude. Occasional spasms affect both the upper and lower limbs. His urine is clear and light, with only mucous sediment. I had not the opportunity of ascertaining its specific gravity. It coagulated with considerable firmness near the boiling temperature, and gave a dense precipitate with nitric acid. This patient was soon restored to health by a steel tonic, an invigorating diet, and by the local plan, of friction with a stimulating liniment, and the use of circular rollers. The urine, examined at the expiration of a week, from the beginning of the treatment, was perfectly healthy.

CASE VI.

T. W. aged 64 ; of sanguineo-nervous temperament ; frequently dyspeptic, and subject to bilious obstruction. A *martyr* to gout. He has numerous uric concretions both in hands and feet. No part of his family has had gout, except an aunt, who suffered from it severely. I first examined his urine under a paroxysm of gout, before any œdematous swelling had taken place, and found it albuminous in a great degree. Its specific gravity was 1·0141. It became turbid at 120°, and, as the temperature advanced, formed heavy flakes. It afforded a dense precipitate with nitric acid. This urine contained a very small proportion of saline matter, yielding from four ounces, by means of nitrate of lead, not more than 4·6 grains of precipitate. It possessed slight traces only of urea or uric acid.

At the patient's restoration to his usual state of health, I was much surprised to find similar characters of urine remaining. At several distant periods up to the present time (January 1819), when he has been in his best health,

and when his appetite and digestion, though never entirely correct, have appeared to be very near the standard of health, I have examined his urine. It has always been secreted in abundant quantity, and I have, without one exception, found it more or less strongly affected, as described, by heat and nitric acid. Its specific gravity in these trials has varied from 1.0041 to 1.0076. These specimens of urine have scarcely afforded the least traces of urea* or uric acid, and a very small proportion only of phosphoric acid; but it is worthy of observation, that of each of these principles, the gouty specimen afforded comparatively the strongest evidence, though still only slight. On each occasion the urine reddened litmus.

At my request, Dr. Prout obligingly examined a portion of this patient's urine, when he was least complaining of indisposition. The following is his statement of the results: "I found the specific gravity of this urine at a temperature of 45° to be 1.0084. It coagulated at a temperature of about 130°, which is considerably below the coagulating point of albumen. I could not satisfy myself so well as I could wish of the existence of uric acid, though I believe it contained a small quantity. It contained also some urea, but less than natural. After having been kept some days in the bottle, it acquired the smell of sour whey, and very strongly reddened litmus; evidently from the developement of acetic acid. The animal matter present differed somewhat from albumen, and approached in its properties to curd, though it is evidently a distinct substance from either." It seems to

* Oxalic acid was used in the examination of this urine for urea, as also the nitric. It is a more delicate test of this principle, although acting more slowly, than the nitric acid. For this fact I am indebted to Dr. Prout.

me probable, that the peculiarities of this patient's urine are connected with the extensive secretion of uric acid, which is continually going on in the hands and feet. He has several ulcerations, from which the inspissated discharge has furnished me with white concretion ; and this, treated with nitric acid, has produced the rose hue.

In my present digression, I purpose briefly to consider the pathological character of a serous state of the urine. Dr. Blackall conceives, that an inflammatory disposition of vessels "prevails in those cases principally, in which the urine is coagulable." In this observation, the reference of the author is made to dropsies alone. Not intending to discuss the accuracy of this opinion, or to animadvert on his practical statements, which are numerous and highly instructive, I shall offer only a few remarks further in connexion with this subject.

This anomalous function of the kidney has appeared to me sometimes associated with nervous irritability, and general weakness, rather than with an inflammatory state of the circulation. I examined the urine of a gentleman under inflammatory excitement from mercury, and its transparency was perfect at the boiling temperature. Some time after, when debility only was present, it coagulated very considerably from heat.

Both in *dropsical* and other urine, which I have found to be more or less albuminous, the kidney has seemed to be sustaining an irritable and hurried action ; the patient, in most cases, passing urine with considerable frequency and irritation, and in some of the examples very abundantly. It must not, however, be understood, that I intend by this observation to ascribe this peculiarity simply to *nervous* disturbance. As a general position, I believe it may with truth be contended, that most of the morbid actions of the kidney, which we find, are derived

from some error in the functions of the digestive organs ; and hence, probably, the source also of the present anomaly. In eight examples of albuminous urine, procured from patients not dropsical, I have found, by careful examination, a remarkable deficiency of urea and uric acid, and also of the usual saline principles. Thus it appears that the secreting action of the kidney, under these circumstances, is very imperfectly performed ; and I conceive, that, in proportion as the urine bears the characters in question in the greatest degree, it is not so much the product of regular secretion, as of *percolation*. As I have above noticed, we do not meet with the usual principles which distinguish the regular secretion of this important fluid. The probability may therefore be further suggested, that urine, which by examination affords the evidence of being serous in a slight degree, does not contain much true albumen ; but that this albuminous principle is in such instances chiefly a modification of the animal matter, which the kidney, at all times, secretes from the blood ; and is not, therefore, necessarily to be regarded as an extraordinary drain of serum from the circulation ; which, by the common theory on the subject, it might be considered.

Proceeding, however, upon the supposition, that urine which is affected by heat and acid, as I have described, is always containing a portion of the serum of the blood, it becomes an important desideratum, to possess the means of forming, with some facility, a presumptive calculation of the quantity which it bears to the urine in any given measure, and to rate the consequent estimate of what may be discharged in the twenty-four hours. Dr. Wells, by inference from comparative experiments, adopted as a criterion, the apparent degree of coagulation, which the urine presented from the application of heat. I have re-

peated the processes which that author has detailed (in the volume before mentioned, p. 224), but I could not arrive at any satisfactory results.

The test of nitric acid, employed as I shall now describe, appears to me, a process at once simple and instructive. I found that one part of serum diluted with five hundred parts either of urine or water, instantly yields to nitric acid a slight precipitate, perfectly white. Diluted even with seven hundred parts, after some time, a sensible precipitate is afforded; and this is the *maximum* of the power of the test. The *immediate* result, however, produced from the first mentioned proportions, is the most convenient evidence. Accordingly, therefore, as the urine under examination will suffer dilution with distilled water, and continue to furnish an *immediate* sensible precipitate with nitric acid, is the proportion of serum to be estimated, as so much *exceeding* 1 to 500. For example, urine that exhibits this effect, after dilution with four parts of water, may be considered as containing 100th of serum, or rather, of albumen. The nitric acid should be added in about the proportion of a 6th, as albumen is soluble in very dilute acids; and without this precaution, the method which I propose will not succeed. I ascertained that the power of nitric acid as a test, was superior to that of heat, which renders the fluid (urine or water) faintly milkish, when the proportion of serum is 1 to 600; but when extended as 1 to 700, does not impair its transparency. Nitric acid produced as much effect on the proportion of 1 to 500, as *muriatic acid* did on that of 1 to 100. The specific gravity of the serum which I employed in these experiments, was, at 60°, 1·0285.

In the further cases which I shall relate in a condensed form of detail, I purpose to hold in view some

different particulars in every communication ; and begin with offering an account of some examples, in which a relapse shortly followed the recovery from a paroxysm ; and of quick returns of the paroxysm in other instances ; in consequence of causes admitting of clear explanation, and affording some instructive lessons both to the practitioner and the patient.

CASE VII.

J. L. aged 46, a mechanic ; has a circular chest, and is corpulent ; of sanguineo-nervous temperament ; rather plethoric ; irritable in disposition ; often dyspeptic and subject to hæmorrhoidal discharge ; habits of living sometimes free, and formerly excessive. Neither father nor mother had gout, but his grandmother on his father's side was afflicted with the disease. This patient was first attacked at the age of thirty-five, in the toe of each foot in succession. The ankles, knees, hands, and elbows, have been affected in subsequent fits. The disease has gradually increased both in severity, and in the frequency of its recurrence.

I visited him first in March 1815. He had been suffering a severe fit, five weeks, and had still active inflammation in the right ankle, right hand, and right elbow. Other parts had been previously affected. He had chiefly employed sudorific medicines, rather a supporting diet, and flannel covering to the parts. His tongue was furred, his pulse irritable, the skin relaxed, countenance sallow, urine of natural colour, but depositing pink and mucous sediment ; the bowels costive, and the fæces blackish and slimy ; the nervous system ex-

tremely irritable. On most nights his gout was severely painful. I adopted the mode of treatment mentioned in the former cases, purging him twice with pills as described at p. 185, and with daily doses of the draught, p. 186; at night tranquillizing him with the pill p. 224. The lotion afforded its auxiliary benefits most satisfactorily. At the end of a week, he was convalescent, and I discontinued my attendance, enjoining a strict attention to diet, care against exposure, &c. and prescribing as internal medicines a mild alterative course of *Plummer's pill*, and a daily bitter aperient; with the use of bandage, liniment, and friction to the weak parts.

Flattered with his rapid amendment, he too soon and too far presumed upon it. He had neglected the methodical treatment which I had prescribed, and carelessly exposed himself to a cold east wind with wet.

In May, I found him again suffering in both feet and one knee; and, together with pain and nervous depression, the stomach was deranged, and the secretions were very unhealthy, indicating, in concurrence with a *lead*en complexion, and some hypochondriac tenderness, that the liver was much deranged in its functions. No alteration of structure was discoverable. The first-mentioned treatment being repeated, he soon again became convalescent; and I now arranged a strict plan which I should superintend. The alterative employment of mercury with the bitter aperient, p. 262, and regulated diet and exercise, were pursued until the secreting functions became healthy, and the complexion indicated the constitutional improvement. Finally, I directed that he should, carefully, at all times, regulate his bowels by the pills, p. 280; take from two to four glasses of sherry, or old port, daily, accordingly as fatigue might require it; avoid

malt liquor, and take animal food once a day only; and every morning pursue the plan of sponging and friction of the limbs, with strict attention.

He has found full advantage from these means. The present period, April 1816, is already a longer interval from gout, than he has enjoyed for a few years past; and his general state of health, together with the improved vigour of his limbs, afford us complete satisfaction.

In this case we see it exemplified, that an unhealthy state of the digestive organs, and particularly of the functions of the liver, may be the foundation of a severe relapse. An exposure to cold, which, in a state of truly recovered health he might have borne with impunity, again excited the gout.

The obvious inference presents itself, that the physician should never pronounce his patient *cured* from the fit, or secure from its early accidental return, until the digestive functions, in the comprehensive extent of the expression, are permanently restored to health.

CASE VIII.

D. S. aged 38; has a circular chest; is tall, robust and corpulent; of a nervo-sanguineous temperament; very plethoric; has been of indulgent habits, drinking wine and other liquors indiscriminately. Father had gout severely: first attacked at the age of thirty-four, in the knee, in the month of June; he thought it was a strain. In the autumn of the same year he suffered another fit, the ball of the toe in the same limb being the part affected. From this period to the date at which I saw him, his gout returned frequently, and with in-

creased severity. Each knee and foot became affected. I visited him first in December 1814. Severe gout was in the feet. He had been much distressed with cramps, and stated that these commonly occurred in the legs, after much exertion in walking or dancing, or from wearing thin stockings, or standing in slight shoes on a damp ground, or exposure to a cold night air. From the same causes he suffered occasional spasms at the stomach. *Punch* was a liquor that did not sensibly disagree with his stomach; but he accused it of sometimes exciting a fit. In some of his former attacks he had taken hellebore and laudanum regularly, and purgative doses of calomel occasionally. From this treatment he procured ease in the affected parts, and shortened the paroxysms; but sometimes he suffered much inflammatory heat of stomach from the hellebore; and, as I have related, the gout recurred quickly.

On the present occasion he had much visceral derangement; the tongue was furred; the bowels were inert without medicine; and such discharges as had been procured, were foul and highly bilious; the urine was of deep colour, and with much pink sediment. I adopted my usual treatment with speedy success; and took leave of him with many injunctions of caution and forbearance. In February 1815, after exposure to cold without a great coat, he became chilly with subsequent heat of skin and head-ache: and the throat became slightly sore. Gout in one foot soon followed, but not with its usual severity; yet the other foot did not escape. The former general and local treatment was repeated. I saw him first on the 6th, the second day of his attack; and on the 16th he felt perfectly recovered. During this period of my attendance, I found that his secretions had not become permanently healthy in the interval of the two fits. I now,

therefore, pursued the usual plan of corrective tonic remedies, till this result was obtained; and pointed out a strict prophylactic regimen, as essential to his permanent cure.

At the present date, April 1816, he informs me, that he has escaped gout and all complaints. He has duly regulated his bowels with the pills, p. 282; and, on being occasionally heated from accidental circumstances, has also taken a portion of a solution of sulphate of magnesia in mint water. He has practised the morning sponging and friction with considerable regularity; and ascribes to this method much improvement in the power of the limbs, and a diminished susceptibility to the impressions of variable atmosphere.

In this example of youth and robustness, an abstemious course of life and vigorous exercise should be steadily continued, in order to counteract successfully a fulness of habit, and secure an exemption from gout.

CASE IX.

J. S. aged 43; has a circular chest; is tall, robust, and corpulent, and very plethoric; of nervo-sanguineous temperament; costive habit of bowels; has hæmorrhoids, and occasionally gravel; perspires freely on much exertion; has been a free liver, and indulged in meat suppers, in the use of porter, wine, and other liquors. His exercise has been irregular. Has lived in London seven years, and his gout has been more frequent and severe since he quitted the country*. He has suffered much from dys-

* The increased frequency of the disease in this patient must not, however, be ascribed to this circumstance alone; but rather

pepsia. His father had gout. This patient was first attacked at the age of twenty-eight, in the ball of the toe of one foot; but since, scarcely any part either in the upper or lower limbs has escaped. Twelve months have made the longest interval of exemption from a fit; and four, the shortest. The longest duration appears to have been ten weeks; the shortest, five or six days. He has often been warned of an attack by symptoms of severe dyspepsia, a very costive condition of bowels, high-coloured and scanty urine, depression of spirits, a numbness of the parts about to be affected; and just at the invasion of the fit, he suffers severely from cramps.

The fit in which I first saw him, in February 1815, had already existed for seven weeks, and was still severe, and affecting various parts. Wet and cold had been the exciting cause; and different situations had been seized in the following order:—left elbow; left knee; right shoulder, right elbow, and right hand. He was entirely in a helpless state; and suffered, each night, pain and fever, cramps and violent startings. He had employed continual means, both by medicine and warmth, to produce sweating; but with no marked good effect. and with obvious increase of the weakness and irritability of his limbs, attended with general languor. The skin of the hand had almost a *scalded* appearance, and the fingers exhibited the character of paralytic weakness. The secretions were in the most vitiated state. The tongue was foul; the urine scanty and much loaded with mu-

to the continuance, or probable increase, of improper habits, co-operating with the loss of country air; and, added to such causes, there is to be considered the sure tendency of gout to increase in the constitution, when unrestrained by true *prophylactic* care.

cous and pink sediment; and the fæces were slimy and bilious. His nervous irritation and depression of spirits could not be exceeded. The successful treatment of a case so long neglected, aggravated by much internal derangement, and attended by morbid sympathies established by repetition, became a matter of great difficulty. To the parts yet inflamed and tender, the lotion was applied with the utmost advantage. Five grains of the compound calomel pill were directed each other night; and the usual pills of opium, p. 225, to be taken as pain or restlessness should require. The bowels were freely acted upon by the draught, p. 186, repeated three times a day; and the diet, which had hitherto been too supporting, was now simply diluent. So early as the 8th of February, he appeared convalescent, and an improvement of diet was admissible. The urine had assumed a healthy appearance, but the bowels were not yet in a correct state. The alterative pill was continued; a draught of infused rhubarb and columbo with magnesia, and compound cardamom tincture, was substituted for the former medicine. The weak and œdematous limbs (together with the usual plan of sponging and diligent friction) were rubbed with linim. camph. compos. and linim. saponis comp. in equal proportions; and rollers were applied.

On the 14th and 15th, he took carriage exercise. At my visit on the 16th I found that he had ventured in the air, the wind being in a cold quarter, with too little caution, and he was beginning to feel its ill effects. On the 17th he complained that his night had been restless and painful. One knee and one foot were rather inflamed, and all the parts affected in the foot were tormented with shooting pains. The nervous system was exquisitely sensible, and his mind was depressed with all the horrors

of returning suffering. The first treatment, both general and local, was repeated; but more frequent doses than before of the opium pill became necessary to tranquilize pain and irritation; although the total quantity of this medicine was not much increased.

At this time, the pink sediment in the urine was again abundant, and the alimentary secretions bore an unhealthy aspect. Very light diet only was allowed, and the medicines were continued. In a few days the pink sediment of the urine was changed to one of whitish colour; and this I found, on trial, to consist chiefly of phosphate of lime with mucus, a little animal matter, and about a fourth of uric acid. The urine itself reddened litmus paper. The state of the bowels gradually improved.

On the 28th he was a second time convalescent; and appropriate treatment, in reference to the favorable change of circumstances, was renewed. The unhealthy state of bilious secretion still demanded attention; and with a view to this point, and the improvement of the whole digestive functions, the alterative calomel pill was directed every third night, and the columbo mixture, with cascarilla and rhubarb, p. 266, twice daily. The local management described on the 8th, was again adopted; but the liniment, p. 271, was now used; the parts so often affected, being extremely debilitated and requiring artificial excitement. I desired that he should take animal food of light digestion once a day, with any vegetable in season; avoid malt liquor and spirits; and drink, after his dinner-meal, a few glasses of old sherry.

These means were pursued with advantage; and the general health being restored, he was directed to regulate the bowels with pills, according to the formula, p. 282, and to be strictly careful in regimen. The limbs

still suffered from much weakness, which had been implanted in them by long neglect. Upon uneven ground the feet would turn suddenly*, with excessive pain, and he scarcely escaped immediately falling. At night the ankles were frequently œdematous, and affected with much aching; and cramps were not yet absent. Twelve months elapsed before I had the opportunity of seeing this patient again; and I had the pleasure to find that he had continued quite free from gout; but his limbs were not wholly recovered. It is true that he had not patiently persevered in friction and other treatment; and I enjoined, therefore, an employment of the means before mentioned, to be pursued with due attention; *tinctura lyttæ* being still added to the liniment.

Soon after this period, I found him gradually recovering the strength and ease of his limbs in the most favourable manner, with every prospect of a perfect cure.

CASE X.

The following case, exhibiting a most striking example of acquired gouty diathesis, appears to me peculiarly interesting in the many instructive views which it affords. It points out the disposition of this disease to increase its severe holds in the constitution, when indulged in its own course according to the doctrines of Sydenham; it exhibits the power of regular treatment; and at the

* He mentioned, "that on first rising in the morning, his right knee would feel as if slipping out of the socket; snapping also, with some degree of pain." Lately, on getting into a carriage, he inadvertently bore his weight for an instant on that limb, and suffered much consequent pain and inconvenience.

same time affords a strong warning as to the care required on the close of the paroxysm, however favourable the circumstances of convalescence may be.

B. M. aged 41 ; of middle height, circular chest, muscular, and of late years disposed to corpulence ; of nervo-sanguineous temperament, and of strong constitution. Not subject to other diseases. Gout unknown in his family. His habits of life for many years free, and without any attention to regimen. Exercise irregular, but strong and violent, particularly on horseback. Was formerly much subject to violent bleeding of the nose. His first attack of gout at 30, in the great toe of one foot only. Thought it was a strain. For the first five years had not either frequent or severe returns. During the last five years, the longest interval from a fit has been eight months ; the shortest, two. Has not noticed any distinct premonitory symptoms to precede his attacks ; but has found himself getting corpulent, and particularly in the abdomen ; and during the summer months to have acquired a very sensible fulness of habit.

For the last three years, the most violent paroxysms have taken place in November or December, and continued, with only occasional intermissions of suffering, till March. In some of his fits, he trusted to nursing alone, with the most simple expedients for the regulation of the bowels ; being taught to believe that the course of the gout should, in regard to the safety of the constitution, be encouraged, and on no account be disturbed. In other fits, however, like gouty patients in general, who are so much prone to extremes in doing too little or too much, he resorted occasionally to irregular and rash methods.

In the fifth year of the gout, the feet being highly inflamed and painful, he immersed them in cold water,

and by its continued use for half an hour at a time, and repeatedly in the course of the night, the symptoms were very much abated. In the night the stomach was affected with spasms; but with respect to the limbs, they were so much improved by the next morning, that he was able to leave his bed and walk. Much internal derangement soon succeeded, continuing throughout the remainder of the year; and the stomach and bowels became in a serious degree subject to flatulent distension, to pain, and the sensations of spasm.

Since that time the paroxysm has been more frequent and violent. In the year 1814 the attack began in the first week of December, after very severe exercise and irregular living during the two preceding months. On the day previously, had been in very strong exercise in hunting, and did not use the precaution of changing his clothes, although very wet. In this fit suffered much from general fever and irritation, with excessive local pain. The feet, knees, and elbows, were affected in succession. Profuse perspiration, and much head-ache. Common aperients, and laudanum, constituted the only means of treatment. Did not recover till the warm weather. In the spring, and through the whole summer, took a table-spoonful of cream of tartar in a tumbler glass of warm water; but notwithstanding this constant attention to the bowels, and considerably more care in the habits of living both as to diet and exercise, a violent fit ensued in the following year. Passed the summer, as usual, well, until September, when he had a slight return of gout in one foot, and was again severely attacked in November; so unfriendly was the disposition of a severe fit to prevent an early return. On the quitting of every paroxysm, severely affected with spasms, mostly seizing the knee, but also the feet.

In this attack in November, the symptoms became more than ever severe and untractable. He took laudanum, and also the black drop, in very large doses, with only slight relief. He underwent frequent and excessive perspirations without the smallest permanent good effect. To the knee, when affected with deep-seated pain, he applied a liniment, by means of strong friction, of sulphuric acid and oil of olives, and vesicated the skin. The parts were much relieved; but in a few days afterwards (not considered as the result of this application) the stomach and bowels became severely and dangerously affected with vomiting and obstruction, and much spasm. These symptoms soon yielded to medicines. In the following week, however, the muscles of the chest, and the collar bone, were seized with excruciating pain. Violent vomiting came on, lasting two days. Then some interval of ease. But in the following week, a painful diarrhœa commenced on the Saturday, and continued until Wednesday, producing excessive weakness. From this time his recovery progressive, but his debility very considerable. Came to London in the middle of February 1816, and in the first week of March was attacked with much pain in the chest, collar bone, and shoulders. Very soon the feet and ankles were inflamed and severely painful. Afterwards, the knees and elbows suffered equally. Purgatives, and some stomachic medicines, were taken during this attack. No return of internal complaint. The paroxysm was in every respect much milder than the preceding one of November. Recovery very slow. In the summer found himself in good health. Took regularly every morning through the whole summer a teaspoonful and a half of calcined magnesia, which acted well on his bowels, and he thought it was affording him a cure.

Such then was the determined character of the disease in this case, and in a constitution irritable in the highest degree. It was scarcely possible to have an example of more difficult and intractable complaint.

I saw this gentleman first in October 1816. He was soon afterwards severely attacked; the fit returning a month sooner than in any former year. On the first day, in the left ankle; on the following, in the right foot; on the third day, in the right foot; in three days afterwards, in the left foot with great violence. I adopted my usual methods of treatment with immediate success. The continuance of the symptoms in each part did not exceed two days, although the hand was seized more violently than on any former occasion. He used the lotion unremittingly. The internal parts were not in the smallest degree affected with spasm or pain. Almost in the first week he began to think himself convalescent. This promising amendment was now sadly interrupted. Eager to get abroad, he exposed himself a little to the atmosphere. He threw off a blanket from his covering at night, and on the very morning of taking a mercurial purgative, on one of the coldest days in November, went into the drawing-room very slightly clothed. In the most sudden manner, the stomach, bowels, and kidneys, were affected with intense pain and spasm, and with the utmost rapidity of symptoms. The pulse was low, and the attack was evidently spasmodic. It yielded to external fomentations, and to purgatives and antispasmodics. He soon recovered from all these serious inconveniences; but while pursuing a plan of alterative medicine, and of course on all accounts highly susceptible, again exposed himself towards evening to the most unfavourable state of atmosphere, a heavy fog with a piercing north-east wind. A severe relapse of gout consequently took place, which became very

tedious ; and my attendance, which before had been interrupted, was now regular. The severity of the symptoms was successfully controuled by the treatment ; and ease and sleep were procured under those intense sufferings, which on former occasions had failed to receive relief from three or four hundred drops of laudanum, administered in short intervals of time. In no case did I ever witness such constant and excessive deposition of pink sediment, or so completely vitiated a state of the alimentary secretions. That the deep foundation of this case of gout existed in a disease of the liver, was verified by the following indications ;—a pain in the right hypochondrium, increased on depressing the ribs by strong pressure ; by the symptoms already mentioned ; by a complexion considerably jaundiced ; and by a tongue always furred. The appetite, however, was scarcely ever interrupted. It would be tedious to detail the exact progress of the case. I treated the disease on the usual principles of conducting the cure of an unhealthy liver, having more than ordinary regard to caution in the use of mercury, for the reasons which I have before stated. The draught, p. 186, was regularly taken first twice, and then once a day, for a long period ; the dose of the acetum colchici being ζ iss. It did not once produce nausea or inconvenience. It will be readily imagined, that some chronic symptoms of pain in the limbs and difficult motion attended so determined a state of visceral complaint ; but a comparison drawn between the sufferings of all his former fits, and on the present occasion, notwithstanding the effects of accident and carelessness of exposure, was to his own mind so favourable, that he spoke of every remedy which was used in the highest terms of satisfaction. No case could be more demonstrative than this of the perfect safety of the lotion ; ap-

plied as it was with unlimited freedom; in a constitution which had so often shewn the tendency to retrocession, and indeed had suffered some actual attacks. After some difficulty and untowardness of treatment, the progress of amendment at length became perfectly favourable; and he can scarcely date the period when he has felt so much freedom from gout, as for some months past. He has, on several occasions, exposed himself to offending causes, with impunity. He is instructed to pursue every point of prophylactic regimen; and to enter upon a course of ass's milk, with a mild plan of alterative treatment, in order that the liver and connected functions may be as radically improved, as medical means can accomplish.

With attention to good rules, he is free from serious fears of relapse. If the gout return, he knows by experience that the painful symptoms may be readily controuled; but he is sanguine from his present satisfactory recovery, that he shall by care obliterate his tendency to gouty attacks.

I do not confine him to any severe restraints in his habits of living, and have allowed the regular use of the purest port, sherry, or madcira, in a limited quantity.

I myself entertain the confident expectation, that, with a due observance of prophylactic regimen, the future issue of this case will be quite satisfactory.

Such was my report of this case in June 1817. I have now, January 1819, to add the following account:

Not to make an unnecessary disclosure of private history, I must briefly state, that at various periods this gentleman has been incautious in his general habits; and often, indeed, has unavoidably been inconvenienced by excess of anxious occupation, without exercise, or any regularity of meals.

In January, 1818, the feet and one knee were affected with gout for one week, in consequence of exposure to cold. This attack was preceded by head-ach, drowsiness, epistaxis, and those sensations of fulness of habit to which he was formerly subject.

In March, two days after receiving a severe fall in hunting, he travelled in the mail to the north of England, and, on a very cold night, with the wind at east, he incautiously slept close to the window of the coach, with the glass down. Thus exposed, and so recently under the influence of a violent shock to his frame from the fall, he incurred another attack, which in two days after took place in both feet, and continued about eight days.

In August, during the extreme heat of the weather, which rendered him feverish, and, in conjunction with great fatigue, very much disordered his system, he was attacked severely in each hand and one wrist, the fit lasting ten days.

On these several occasions, my usual active treatment proved perfectly successful, and a speedy controul was obtained over all the symptoms, without the least threatening of retrocession, or any untoward occurrence. In the attack of March he travelled a great distance in a carriage, when suffering very agonizing symptoms of gout in the feet; so that it required all his fortitude to endure the torture which he experienced from keeping his limbs in a depending position, and from the concussion of the vehicle. He submitted himself to this painful and hazardous exertion from the desire of returning to London. His journey being completed, I visited him on the 23d, and found his legs frightfully swollen, with an inflammation of an erysipelatous appearance extending in patches from the feet to the knees. In addition to the immediate pain of the gout, his system was in a high

state of synpathetic irritation, and the spasmodic jerks which affected the limbs were unusually severe. So successful, however, was the combined method of treatment which I employed, consisting internally of the *saline* diuretic purgative, p. 186, with occasional doses of the mercurial pills, p. 185, and the narcotic pills of crude opium and James's powder very freely exhibited, the external remedies being the lotion in the day, and the poultice at night, that at the end of eight days he began to be favourably convalescent; and on April the 4th, felt so well and cheerful, that he was very desirous to engage in his favourite amusement of hunting.

Since August he has been free from gout. In his last communication, a few days ago, he says: "Upon the whole I feel much better in health than I have been for several former years, and I flatter myself that I shall be able to avoid any serious attack."

It is evident, that in this gentleman's case the gouty diathesis prevails so strongly as to demand unremitting attention in regimen and in all his habits, with an occasional course of medicine. He has of late constantly regulated his bowels with the draught, p. 186, and highly praises its effects. The attacks which he suffered in the last year were strongly provoked by occasional want of care, or accidental causes; such as in another individual must have produced illness of some kind, but in his constitution produced the gout. Yet the short duration of the paroxysms, and the immediate relief afforded by the treatment, affords a happy contrast to the extent of his former tedious fits, in which the severe symptoms were not on any occasion controuled by the ordinary routine of treatment. On one occasion, he relates that he gave trial to the *eau medicinale*, without deriving from it even temporary relief.

CASE XI.

The following case of gout is an additional example of a remarkable frequency of attack, even where the diathesis was wholly acquired, arising chiefly from the influence of a morbid state of the liver; and consequently pointing out, that in these instances our only method of cure for the gout is the adoption of that treatment which is the most suited to restore the action of the liver to health, and also to rectify the digestive functions in general. It further demonstrates that the action of gout, although not interfered with by any treatment, entirely fails to prove a curative remedy for such internal derangement; and apparently serves only to harass and depress the powers of the constitution.

S. J. aged 40, of middle height and bulk, and not very plethoric; of sanguineo-nervous temperament, and strongly bilious diathesis. Seventeen years ago had the yellow fever in a hot climate, the shock of which to his constitution he has never recovered. Once since he has been severely jaundiced. Of moderate habits, and usually very careful. In general sedentary, and disposed to retirement. Gout unknown in his family. First attack was in January 1814, in the great toe of one foot only. Has since had five regular paroxysms. Passed two winters in Paris, without any advantage from the climate of that city, as compared with a residence in London. His attacks always violent. In the fourth fit, both feet and hands were affected. He usually adopted only simple treatment; and in the first two attacks did not use any remedy, but merely covered the parts with flannel.

In one paroxysm took magnesia very freely, which

acted beneficially as a purgative; but on being discontinued left the bowels costive. Has occasionally complained of great debility of stomach, and, according to his own statement, “accompanied with shiverings, coldness of the extremities, weak pulse, very low spirits, total want of appetite, and, as it were by instinct (it being so contrary to his custom), a desire to take some spirituous liquors.” Also, when he has found himself dyspeptic and bilious, has complained of the sensation commonly called that of *pins and needles*, in the legs particularly, and in the arms slightly.

In the third fit, having severe pain, inflammation and swelling in one foot, he went into a warm bath at 95°, which for about a quarter of an hour very much relieved the intensity of the pain; but on his quitting the bath, it returned in an increased degree: and this fit proved of more tedious duration, and produced more œdema and weakness of limbs, than any which he has ever experienced.

I was consulted by this gentleman in January 1817, when the gout was affecting first the feet in the usual manner, and afterwards the hands. In union with this attack much jaundice was present, a furred tongue, loss of appetite, depression of spirits, the urine of a deep colour and depositing lateritious sediment, the fæces of a dark green colour and in every respect indicating bilious derangement; and the right hypochondrium was also tender on pressure. I directed in this case, repeated purgatives of calomel and colocynth, together with the colchicum draught, p. 186, and use of the evaporant lotion, until all the acute symptoms were removed. Except on one night, the aid of a narcotic was not required. Of the lotion he stated, “that it quickly relieved the pain, heat, and throbbings of the parts, and he soon

found his feet sensibly strengthened from its use." He did not suffer the subsequent œdema and debility of the ankle joints, which, on previous occasions had rendered him very infirm. He had recourse to the tepid salt water sponging in the morning, with bandages during the day, and a stimulating liniment at night, with the best effects, in the convalescent stage. He pursued a mild corrective course of mercurial medicine, in conjunction with the stomachic draught, described at p. 262, and took ass's milk night and morning.

By these means, steadily pursued, in unison with a strict attention to general regimen, he gradually lost his jaundiced complexion; no longer was sensible to any pain on pressure in the region of the liver; acquired a good appetite, cheerful spirits; and, in short, a general renovation of energy and comfort. Very soon afterward, this gentleman went to the Continent; and I have not had the opportunity of hearing of his health since.

CASE XII.

Of the following case I shall give a brief sketch of the particulars, with which the patient has furnished me, of the general history of his gout; together with such details as appear to me to set forth, in an interesting point of view, the decided efficacy and importance of active treatment in a severe paroxysm. The account also embraces other valuable facts.

C. R. aged 48, tall, corpulent, of plethoric habit, and the mixed temperament. Not subject to any other disorder than gout and rheumatism. The gout not known in his family. First attacked at the age of 21. In the first six or eight years, the paroxysm took place about

twelve or one in the morning, without any premonitory symptoms; but subsequently it was preceded by dyspepsia, and cramp-like twitchings of the legs and feet. In some of the fits he was bled freely; which served to abate the inflammation, but did not shorten the duration of the paroxysm, nor materially relieve the pain. The longest fit which he ever had, followed the practice of bleeding to the extent of 90 oz. in the space of four or five days. From leeches, used with great freedom, no good effect resulted; and sometimes they served to increase the inflammation, irritation, and pain. Seven years ago, made an attempt to abridge his diet; eating meat only each other day, and abstaining from wine entirely; but allowing himself table beer. The gout returning, he resumed his usual habits of living, which of late years had been guarded, but not severely restricted. His fits after a time becoming more than usually severe, he resolved upon a determined plan of rigid abstinence. For the space of sixteen months, he abstained wholly from animal food and wine. He allowed himself some table beer daily; but in solids confined his diet to vegetables, milk, and plain puddings. By means of this plan, the state of his system became materially altered. The bowels were uniformly regular; the hæmorrhoidal discharge, to which he had been much subject, did not occur; the urine, which from every accidental excitement to the circulation deposited the pink or lateritious sediment, was invariably transparent, and light in colour. These changes began to take place in about six weeks from the commencement of the plan. In the course of this probation, he had some attacks of gout, with much less inflammatory action than before, as may be supposed; but, in opposite balance to the advantages which I have stated, his nervous system was affected in a distressing manner;

he was always *comfortless*; very continually suffering general aching pains, which he indifferently called nervous, gouty, or rheumatic; and sometimes had short attacks of acute gout. His legs very commonly became much swollen after slight exertion. In short, he had in this way so little cause of satisfaction, that he returned to the use of animal food daily, and wine in a very restricted quantity, occasionally. This improvement of diet proved very favourable to his constitution; and, with the great attention which was paid to the action of the bowels and the state of the secretions, he was not more affected with gouty feelings than before, during many succeeding months.

In regard to the frequency of attack, and the treatment which had on most occasions been adopted, during a series of years, I shall further cursorily mention the most interesting circumstances. In 1810, when suffering from a severe paroxysm in December, which had been preceded by an attack in October, he took the eau medicinale. The relief was so immediate, that he went out on the following day. A relapse took place in less than a fortnight. The eau medicinale was repeated with the same relief. In the spring of the following year, two or three slight attacks occurred, but a very severe one in November. The eau medicinale was taken early in this fit, and it seemed only to suspend the disease for a few days. It was then repeated. The effect was rather more evident; but it produced distressing vomiting, and the fit was lingering. In 1812, when under a second attack in the year, he reluctantly repeated the eau medicinale. The pain was somewhat relieved, but the sickness was insupportable, and the stomach would not retain either liquids or solids for many days. In 1813, he laboured under a paroxysm for three months, taking ape-

rients and sudorifics regularly with very little relief. In 1814, he pursued the plan of abstinence. Suffered very troublesome pains in the wrists and hands, which continued many weeks. In 1815, he placed his dependance on the use of *vinum colchici*, which relieved the pain; but the fit lasted a month, and the recovery was tedious. In 1816, he suffered from gout as usual. In 1817, a fit took place in September, lasting a fortnight; but, after the interval of a week, he relapsed and was confined to his chamber between three and four months. In this paroxysm scarcely any medical treatment was used, and never more than slight expedients, under the supposition, that, if Nature were permitted to relieve herself, the future intervals between the attacks would be longer, and the general health would become improved. In this expectation, however, the patient was disappointed. The fit finally left him tormented with chronic pains, which sometimes appeared to be gouty, sometimes rheumatic. Under these circumstances my opinion was consulted; and I advised the combined use of sedatives, an alterative pill, and the draught, p. 186, to be continued as circumstances should direct, in conjunction with the regular use of the vapour-bath*. He gradually got into better health, and derived very sensible relief and evident benefit from this last remedy. He stated, indeed, that it

* He made use of a portable bath, which was constructed after a model obligingly lent to me by Dr. Park. It is truly convenient in its application. I have engaged a male and female attendant to render their services when wanted, for the conveyance and administration of the bath to a patient; and shall be happy to extend its easy employment for other practitioners, by any communication and explanation in my power.

produced a more capable and comfortable state of limbs than he had experienced for many years.

In 1818, a fit rather severe, induced by cold, came on in September, after his having enjoyed more than usual health and exemption from pain for a considerable time. It continued nearly three weeks; but the symptoms were less acute than usual. He was not confined to bed. He resorted to the vapour bath with much relief. The poultice, p. 249, was remarkably useful, and the other treatment which was occasionally adopted, also succeeded. On the following occasion, however, my combined plan of internal and external treatment was conspicuously successful; and allows of a flattering comparison with the results of the empirical practice; the ordinary remedies of the physician; and the influence of Nature left to her own operations.

On December 26th, after a most unfavourable exposure to cold in the middle of the night, he was attacked severely in the knees; and almost every part of the upper and lower extremities became quickly affected in succession. I never witnessed a seizure of more determined violence. The symptoms were of the highest intensity. The deposition of bright pink sediment in the urine was more copious than I had ever before noticed. The skin was rather jaundiced, and the state of the bowels also proved the surcharge of the bile and obstruction in the proper channel. Recollecting the tedious sufferings of 1813 and 1817, he apprehended a similar continuance of confinement. To avoid prolixity of detail, I shall merely observe, that the remedies already amply described were administered in full force. During the symptoms, the kidneys were attacked with excruciating pain; at which moment the limbs were either more easy, or the attention was absorbed in the greater suffering. A copious bleed-

ing from the arm gave immediate relief; and when ease was obtained, a small uric calculus was voided. The blood was sizy and much cupped. To conclude—the patient obtained so satisfactory a removal of this paroxysm, that, in rather more than a fortnight, he took walking exercise out of doors; and has not, on any one day since, been confined to the house. He follows up the treatment upon a mild alterative plan. When, from the exertions of the day, he experiences aching and some sense of heat in the ankles so recently affected, he applies the poultice at night with the happiest effect. He uses the morning sponging with salt water; and occasionally in the day, applies the lotion to the hands and wrists by sponging. His appetite and spirits are quite recovered.

February 3rd. He continues well.

CASE XIII.

A gentleman, aged 48, of sanguineo-nervous temperament, and of very bilious diathesis; affected with hereditary gout, and having suffered frequent fits in the course of the last ten years, was attacked with severe acute symptoms in each foot, and one knee. A few days previously, the second toe of one foot had accidentally suffered tight pressure from a new boot, so that pain and swelling soon followed as a consequence. At the moment when the inflammation of the other parts was at the height, the integuments of the toe were evidently in the state of abscess; and, on a puncture being made, a considerable quantity of curdy matter, mixed with particles of chalk-like substance, was forcibly pressed out. This, on being submitted to chemical examination, pre-

sented results precisely similar to those which I have described at p. 291. One of the fingers also, which had for a long time past been thickened in its tendinous sheaths, in this paroxysm, without the influence of external injury, became more than usually swollen, and so distended near the surface on the palmar side, that some fluid was distinguishable to the touch. On being punctured, much blood, and a small portion of the curdy pus, were discharged. This pus agreed in all its properties with what was furnished by the abscess. The thickened finger, which had long been deprived of free motion, was treated with the empl. hydrarg. cum sapon. in the day, and a mercurial poultice at night, with the best effects; and the toe was soon restored by simple treatment. The gouty inflammation of the feet and of the knee yielded very favourably to the combined influence of purgative mercurials, the draught, p. 185, pills of crude opium and antimony, and the free use of the lotion; but I have chiefly introduced the short narrative of this case, as affording a striking instance of gouty inflammation proceeding to specific *suppuration*. In this, as in the former example, Case ii, the common integuments were the seat of the abscess. In the toe, which was injured from violence, we may conceive that common inflammation and gouty inflammation were mixed; but it certainly appears, that in gout, when the capillary vessels are from any cause excited to that increased action which is analogous to the suppurative, a secretion of urate of soda is the consequent result.

In the present case, during the whole continuance of the inflammatory symptoms, the urine deposited an excessive quantity of the bright pink sediment, and it was much loaded with urea. The fæces abounded with vitiated bile; pain; and tenderness to pressure in the right

hypochondrium, were felt considerably; and indeed every indication of an unhealthy state of liver, requiring an alterative course of treatment, was strongly marked.

It deserves to be noticed, that this gentleman had, a short time previously to this attack, made regular use of Dr. Scott's acid bath for five weeks; at first with seeming benefit, but eventually without the least sensible or apparent advantage; and certainly I found him labouring under bilious obstruction in a great degree. This, however, readily yielded to the purgative and alterative treatment which I employed; and, in its decisive result, might fairly be contrasted with the truly equivocal operation of the bath. I may also mention, that a few years ago he made trial of the eau medicinale; which, in the dose of half a bottle, affected the bowels considerably, and quickly subdued the symptoms, but left the stomach very dyspeptic. The gout returned in three weeks, and assumed a chronic form. He now took the eau medicinale in doses of twenty drops daily, by which the pain was relieved; but his stomach became materially weakened, his legs much swollen, and the nervous system was affected with many distressing symptoms. He soon therefore relinquished the medicine.

He had never before experienced any signs of uric concretion; but has always noticed the pink sediment to be abundant in the urine during the paroxysm; and it was so in an extreme degree in this attack, during the whole continuance of the inflammatory symptoms; marking an extraordinary disposition in the system to the secretion of uric acid.

He stated, that in former fits not commencing with more violence than this, he had suffered far more severely, and for a much longer time. At the period of convalescence in this case, I directed an alterative course of medicine to be pursued for several weeks, with the free use

of ass's milk ; and that the empl. ammon. cum hydrarg. spread on leather, should be worn on the tender part of the hypochondriac region.

May 1818.—Has continued quite free from gout ; but having been on the Continent, and lived on foreign cookery, his stomach has become unhealthy ; and on this account, and with the view of preventing his usual annual return of the gout, he adopted, by my advice, a course of medicine for three weeks with perfect success. I have reason to believe that he has remained free from gout up to the present time (January 1819), and that he is in good health.

CASE XIV.

This, and the following case, are additional examples to the foregoing ; shewing with how much certainty a secretion of urate of soda takes place, when the gouty inflammation does, as a very rare occurrence, proceed to the extent of suppurative action ; and it is probable that, in such peculiar constitutions, this result may happen in whatever texture the inflammation may be seated ; although it may be expected to take place most frequently in the common integuments.

A gentleman, aged 37, tall, robust, and rather corpulent, was first attacked with *hereditary* gout in the great toe at the age of 24 ; and from that period has seldom passed a year without more or less of severe gout. In his last fit, both feet, both knees, and one hand, were violently affected with inflammation. An abscess formed in the bursæ at the ball of the great toe, and also at the heel. Each was opened, and a large quantity of the uric secretion, with pus, was discharged. That from the heel was of a beautiful rose colour, the only instance of the

kind which I have seen. The parts were treated with simple bread poultice. The bursal membrane continued to secrete this peculiar matter for a few days, and then healed favourably. In all the previous fits of gout, the inflammation had not proceeded to this termination; nor was any concretion to be discovered either in the upper or lower extremities.

In this case, as in the former, the urine gave a copious deposit of pink and lateritious sediment in the course of the fit. However much, therefore, this extraordinary action of the capillary vessels may be vicarious to the natural function of the kidney, in its secretion of uric acid, under the influence of chronic gout, as I have suggested in the way of hypothesis at p. 145; it is evident, that when *acute* symptoms are present, the occurrence of this phenomenon does not supersede the function in question; which, on the contrary, itself takes place in a preternatural degree.

CASE XV.

A gentleman, aged 39, tall, of middling bulk, of bilious diathesis, was first affected with *acquired* gout at 34, in the great toe, and has since been attacked in almost every part of the upper and lower extremities. Has a considerable quantity of uric concretion in the cellular membrane at the elbow, in the bursæ, at the ball of the great toe, and also in the lobes of the ears. Lately, a heavy person trod on the toe, and excited so much inflammation, that an abscess formed. The skin over the bursa being much distended and very thin, a puncture was made, when a large quantity of the old concretion in the state of thick paste, and of the new puriform curdy secre-

tion, was pressed out. A simple poultice was applied ; and some pressure was used daily to remove any fresh portion of secretion. The wound shortly healed. Soap plaster, spread on leather, was worn over the part, and changed every second day. It promised to be perfectly cured. As the constitutional treatment in these cases was, in essential points, similar to that which I have employed in ordinary cases of gout, I have, for the sake of brevity, avoided such details. I shall here conclude my account of Acute Gout, and proceed to the next division of my subject.

CHRONIC GOUT.



THIS species of gout, according to its characters, which I have offered at p. 14, occurs most commonly as an ultimate consequence of the acute form of the disease; and appears when the paroxysm has not formed a crisis; or, when repeated attacks have so much enfeebled the constitution, that strong inflammatory action no longer takes place. In this case, the former severe paroxysms, which occurred with distant intervals, become exchanged for those which are milder, but which are more frequent and irregular.

This chronic state of gout is more commonly blended with wandering pains than *the acute*; and these pains have now and then the rheumatic character, and are at other times indefinitely nervous. It occasionally occurs in subjects, in whom the acute form has never existed; but examples of this kind are comparatively rare, and are found more frequently among women than men. In this mode of attack, it seldom happens that the great toe is the part affected; but sudden swelling and pain, having but little of the regular character of gouty inflammation, fix in preference on the hand or wrist, or instep, and about the ankle.

When the chronic gout occurs as a sequel to the acute, the various parts which have been inflamed in the paroxysm, continue affected alternately or in conjunction.

The following may be offered as a further description of the local and constitutional symptoms in both the states here mentioned, viz. when original and when consequent.

The sensations of the affected part are rather those of heat and coldness alternately, than of the more continued *burnings* which takes place in the acute disease ; but, as in the acute, the night time is most the period of active pain. Frequent numbness is present, and an uneasy sense of fulness, bulk, and weight. The muscles, tendons, and ligaments appear, from weakness, unequal to their office ; and cramps, which especially affect the lower limbs, occur even in the day, but mostly at night, when the patient attempts to take his first sleep*. Starting also, in the imperfect sleep which is procured, take place in a very distressing degree, both when chronic gout is present, or when it is threatened. If any redness appear on the surface of the pained parts, it is of a pale colour and usually transient : sometimes it is of a purplish hue ; but often, indeed, the skin retains its natural colour, and especially when the knee is the part affected. The bursæ and the sheaths of tendons are much more the seats of complaint in the chronic, than in the acute gout ; and these textures acquire a very continued state of puffiness and distension, producing, at the wrist and instep, the matted feeling of parts to which I have adverted in a former page. In situations more cellular, œdema is very permanent ; and this is attended with a preternatural fulness of the adjacent veins. In the most favourable state of the general health, the ankles are affected with

* Sydenham observes, “ The tendons of the muscles of the tibiæ are sometimes seized with so sharp and violent a convulsion, or cramp, that if the pain it occasions were to last only a short time, it could not be borne with patience.”

aching and a distressing sense of heat after every unusual exertion in walking. The tenderness of the parts on pressure; the shooting pains of different nerves; the painful difficulty of motion, and entire want of energy in the limbs, under any effort of the will, belong also to the local debility produced by the disease. The state of the constitution in chronic gout embraces a great variety of symptoms, which are modified by the temperament and the habits of the patient; by the situation and degree of local disease; and also by the seat and nature of the internal visceral derangement. Indeed so many anomalies often arise in this impaired state of the health, partly depending on internal causes, and partly on the painful or uneasy state of the affected textures, that probably no description would be adequate to include all these shades of sympathy; and it may be sufficient to delineate a general outline.

It most commonly happens, that the patient is severely dyspeptic, and that the stomach is distressed with various uneasy sensations. A craving desire of food, and nausea, are often felt alternately. Oppression after an incautious meal, and flatulent distension, attend the gouty dyspepsia in an urgent degree; and to this may be added *heart-burn*; or, in some individuals, a coldness at the stomach of a peculiar nature and intensity, compared occasionally to that of marble, or even of ice. Fugitive spasms affect the muscles of the abdomen, or ribs; or cramps occur, which are of more distressing continuance.

In many examples in which the appetite seems natural, the patient has not the feeling of benefit from his diet; and when this is of too stimulating a kind, the irritation produced by local uneasiness grows into feverish action. An exceeding irritability marks the temper. The mind is hypochondriacal; imaginary evils disturb the

judgment, and shake the resolution on trifling occasions. Palpitations affect the heart; and the sensations, described as flutterings, are still more frequent. Either from pain or uneasiness, the sleep is disturbed and unrefreshing. I have met with female gouty patients in particular, who are so exquisitely sensible to the vicissitudes of atmosphere, that instantly on the change of the wind to a cold quarter, and especially if accompanied with moisture, they feel wandering pains in the limbs; and indeed are so susceptible, that their nerves are true barometers.

In the worst instances of the disease, a general *cachexy* takes place; or, the former bulk of health is partial; so that the lower limbs are wasted and weak, and the abdomen becomes large. The secretions are more or less vitiated. The bowels are in opposite states, but for the most part costive; and the bilious secretion is deficient and unhealthy. The hæmorrhoidal veins are often painful; and blood, either arterial or venous, is occasionally discharged; but, when in much quantity, it is venous and very dark. The urine is variable, being influenced by many circumstances. In some instances it is in sparing quantity, and much concentrated; in others it is abundant and dilute. In that of the morning, there is usually much mucous deposit; and under excitement of the circulation from disease or diet, the uric sediment (pink or lateritious) is almost certainly present.

A gentleman, much afflicted with chronic gout, informs me, that he notices his fæces to be particularly clay-coloured, and that the pink sediment appears in his urine, when he is most complaining of weakness and pains in the limbs. He readily suffers dyspepsia in a distressing degree from slight causes.

A sallow appearance of the complexion; sometimes a

yellowness amounting to slight jaundice ; occasional pain in either hypochondrium ; a torpid action of the bowels, and an evident deficiency of bile in concurrence with much irregularity in the quantity and appearance of the urine, more especially when it deposits the pink or late-ritious sediment ; comprise a condition of the constitution which frequently attends upon chronic gout.

A chronic cough is not unusual ; and in some cases it is of a primary nature ; but more frequently it is truly sympathetic, and depending on the faulty state of the digestive organs. I have repeatedly seen instances of this kind, in which the patient and his friends have been alarmed for the safety of the lungs ; but the cough has entirely yielded to the treatment best adapted to the threatening symptoms of gout ; and in these cases, I have not forbidden the moderate use of wine. I have described the nature of this cough at p. 17.

In other instances, an habitual excess of mucous secretion in the trachea, proves a very troublesome symptom, even when no cough is present. This, which is represented as a viscid nauseous phlegm, accumulates during the night, and causes much difficult expectoration in the morning.

Another example of chronic gout may be stated, in which the natural functions appear, at the first view, little disturbed from their healthy action ; and irregular pains, and transient inflammation at the ordinary seats of the disease, with consequent lameness, constitute the patient's chief complaint.

In this case, however, it will be found on accurate investigation, that the internal functions are not healthily performed. Although the tongue may bear a healthy appearance, and the appetite may be good, the state of the secretions is more or less faulty ; there is probably

congestion of vessels below the stomach ; the general balance of the circulation is interrupted ; and the nervous system, as a certain consequence, is much disturbed.

Such are the forms of broken health and frequent sufferings, which chronic gout produces. When neglected and left to its natural progress, its inroads on the constitution are so sure and constant, that the afflicted patient, in many cases, is indebted to the genial summer alone for a state of partial ease.

In this state of the frame, any internal disease which is casually produced, assumes a modification more or less remarkable, in consequence of the gouty diathesis ; and, as I have formerly observed, such occurrences have been the fruitful source of error and confusion both in theory and practice.

I shall endeavour to give a more exact and familiar description of the nature of chronic gout, than I have here attempted, in the several cases of the disease which I shall presently offer.

CAUSES.

WHEN it is considered that the chronic state of gout is truly a modification of the acute, and that it is essentially the same disease with a difference of degree and circumstance ; it follows as a consequence, that the same general pathology, with an exception which I shall presently offer, must be applicable to each. I trust, therefore, that in my remaining pages, I may with propriety adopt a convenient conciseness of discussion ; and depart from that close analytical arrangement which I have hitherto endeavoured to observe.

It appears to me that the difference in the agency of the remote causes, either as producing the acute or chronic forms of the disease, must be wholly referred to the particular state of constitution. If, when the gouty diathesis prevails, the system possess considerable energy, and the action of the heart and arterics be in sufficient vigour, the acute symptoms will be produced by the free application of the remote causes.

If, on the contrary, languor and debility possess the frame, chronic symptoms and various anomalies arise.

In the theory which I have offered of the *first* paroxysm of acute gout, as it usually appears, I have considered that a redundancy of blood, chiefly affecting the circulation of the vena portarum, induces, in persons of a certain predisposition, temperament, and diathesis, the peculiar inflammation of gout. In a favourable case, Nature with her own powers relieves the system by this process, and health returns.

The circumstances of the chronic disease, as the term itself expresses, are very different; and the constitution is involved in a more complicated manner, than in this theory of the first fit of *acute* gout. To the returns of the acute attacks, however, a similar principle of pathology may apply, as to the chronic; observing the modification which arises from the different state of the powers of the circulation and of the nervous system, as I have so lately explained.

It remains that I should briefly describe, in a more particular manner, the chief causes which influence the production of chronic gout.

It sometimes happens, that the gout not appearing till the patient has arrived at an advanced age, it is connected with a plethoric state of system which is joined with much debility; and then, chronic symptoms alone

take place. The disuse of former exercise, and increased indulgence in the general habits, introduce the disease in this manner.

Examples of the gout assuming in its first invasion the chronic form, as early as the middle age of life, most frequently occur in females of weak constitution, whose parents have been, one or the other, gouty. It may be added, by way of explanation, that the remote causes are applied more weakly in the female sex; and hence, in women, the gouty action is often less completely evolved than in men. The mismanagement of the paroxysm is probably the most frequent source of the chronic symptoms. The neglect of purgative medicine, and the injudicious employment of stimuli, produce irregular inflammation, indirect debility, and excessive nervous irritation. The improper use of mercury powerfully contributes to produce these effects. Hot clothing of the inflamed parts, as is usually practised, prolongs the symptoms, and increases the tendency to relapse from slight external causes.

In the examples which are most obstinate and untractable, the cause will certainly be found to exist in some visceral disease. The liver itself is most usually the chief organ affected; but I have sometimes been persuaded that the spleen has also betrayed strong symptoms of disease; and which have concurred to support the gout. It is in the chronic form of gout that we often meet with the worst instances of dyspepsia, and that the urgency of flatus is so remarkable. This state of complaint is certainly existing in connexion with a vitiated condition of the bile, and of the whole alimentary secretions. A deficient or irregular action of the kidneys is also an usual attendant. In the majority of cases of chronic gout, it will be found, according to my observa-

tion, that a state of unhealthy function in some department of the œconomy of digestion is the great supporter of the symptoms; and that the irregularity in the general circulation, and also the train of nervous symptoms which so commonly supervenes, are entirely of a secondary nature. An obstruction in the vessels of the vena portarum, giving rise to the features of complaint described at p. 81, and subsequently, is the most ordinary deviation from health in the digestive organs, which is found to occur with gouty persons.

The *eau medicinale* has been the fruitful source of many cases of chronic gout, by enfeebling the nervous system; and occasioning, together with irregular pain and obscure inflammation, a degree of despondency and languor never before experienced. The same observation applies to the use of other gout specifics, as Wilson's and Reynolds' tinctures, and the *vinum colchici*; and I can with truth affirm, from troublesome experience, that no chronic cases of gout are so tedious and untractable as those which have been empirically treated by these medicines. On a former occasion I have fully explained their merely palliative and temporary influence; and that although they subdue the present symptoms, they leave the inherent seeds of the disease in stronger and more radical force. By the frequency of relapse, there becomes added to the original difficulty of cure, that which belongs to the *habit* of diseased action; in the same manner that an intermittent fever of long standing resists the same remedies, with more obstinacy than one which is recently contracted.

In proportion as the gouty diathesis is strong, and the constitution is unsound, so will the power of all the remote causes be more strongly marked. Thus, where the susceptibility is established in a great degree, every

accidental exposure to wet and cold, and even the influence of the east wind alone, will more or less give rise to flying pains, and inflammatory action of a slight and transient kind.

Distress of mind acts powerfully in keeping up the symptoms of chronic gout. I have seen very strong proofs of this fact; so that, in some constitutions, while the feelings are anxious and much afflicted, it is truly difficult to accomplish a cure.

DIAGNOSIS.

THE distinction of chronic gout from *chronic rheumatism* is seldom very difficult, when all the circumstances, both general and local, are carefully considered. Chronic rheumatism, not preceded by the acute form, occurs much more frequently than chronic gout*. Our opinion will be assisted by reference to the constitution of the parents. The presence or absence of dyspeptic sympathies is not a certain criterion of distinction; but it may be stated as a general proposition, that the natural functions are much more deranged in chronic gout, than in chronic rheumatism, and invariably, according to my observation, when the disease originally fixes itself as chronic gout; but when the occurrence is subsequent, the distinction in question is less remarkable. I have seen some examples, in which, the pains of chronic gout affecting those who have had frequent acute fits, have very much assumed the character of rheumatism, and

* Dr. Haygarth, in his *Clinical History of Disease*, mentions, "that out of 470 cases of rheumatism, only 170 had the rheumatic fever."

have not been attended with any disorder of the appetite or digestion. The structure, constitution, and temperament of the patient; his age, and his habits of living; together with a consideration of other remote causes, must be brought under review. Dr. Clerk, with great confidence, founded his distinction of the two diseases, as he says*, “on white ropy filaments floating in the urine, which when taken out of it are pellucid, and when dried turn to a kind of calx.” This cannot be justly considered as a pathognomic symptom. Such an appearance of the urine (produced by mucus and saline matter) occurs in various kinds of disease, in connexion with disordered digestive functions.

The character of the local appearances, in chronic gout and chronic rheumatism, is an important point of distinction. Although the gout in its chronic form is still more fugitive and uncertain in the part which it occupies, than when acute; and in this respect approaches nearer to the nature of rheumatism; yet it is much more disposed to attack the hands and feet than the last disease, and also to be more solitary in its situation.

The circumstance of rheumatism affecting tendons at their insertions, the aponeuroses of muscles, the ligaments, and branches of nerves, more commonly than any other parts of structure, and consequently not producing in these situations any very evident swelling, it has been assumed by Dr. Haygarth†, “that absence of tumour

* Edinburgh Phys. and Lit. Essays, vol. iii. p. 442.

† On the discrimination of chronic rheumatism from gout, acute rheumatism, scrophula, nodosity, white swelling, and other painful diseases of the joints and muscles.—Med. Trans. of the College of Physicians, vol. iv.

constitutes a clear criterion of chronic rheumatism, from acute rheumatism, gout, scrophula, nodosity, and white swelling of the joints."

In a reference to three hundred cases of chronic rheumatism, he remarks, "Out of the whole number, only fourteen patients were noted with any swelling in the seat of the disorder; and it appeared, upon a more careful and deliberate investigation, that all these fourteen cases ought to have been classed under the other genera." I am now confining my notice to gout; and to me, I confess, it appears, that the ground of diagnosis which is here chosen, is neither sufficiently faithful nor comprehensive. The result of my own observations has been, that if the chronic gout affect the foot or hand generally, an œdematous swelling is almost certainly present; but if the chronic rheumatism be so situated, this occurrence is exceedingly rare. If, however, the *bursæ mucosæ* and the *thecæ* of tendons be the seat of complaint; namely, either of chronic rheumatism or chronic gout, the external characters are very similar. Under each disease, the bursal membrane becomes distended to a great degree; and this is often seen in a remarkable manner at the knee joint. The mutual resemblance also in the distension of the tendinous *thecæ*, which, at the wrist particularly, produces considerable fulness of parts, further serves, in my apprehension, to destroy the accuracy of the diagnosis in question.

The textures which have been often affected with gout, become so much debilitated as to be very susceptible of vicissitude of temperature, either in the house or abroad; and in this way, the general disorder may often partake of *rheumatism*. It is only from such combination, that I can attach any propriety to a very common expression, *Rheumatic gout*. I do not conceive that

gouty inflammation and rheumatic inflammation can exist in *the same part*, at the same time; but certainly we find gout and rheumatism occasionally existing in *different parts* of the body at the same time. Of this last fact I have seen many instances. The patient, when he has gout in the regular situations, suffers, in consequence of some partial exposure to cold, a rheumatism in other parts, as in the muscles of the neck, or in the shoulder joint: and a seizure of lumbago at the time of the invasion of the gout is also not uncommon.

The peculiar affection called "Nodosity of the joints*," is distinguished by Dr. Haygarth according to the following characters:—*Almost peculiar to women; without fever; the swollen joints on examination representing an enlargement of the bones themselves; the comparative freedom of the integuments from inflammation; the muscles seemingly not affected†.*

In some examples of this disease which have come under my observation, the bone-like hardness of the affected parts; the general state of freedom from pain, and which at no time was severely felt, together with the absence of sympathetic disturbance of the digestive organs, have sufficiently convinced me of the distinction of the complaint from gout. On the other hand, I have seen a few instances, in which the gout has always been so local, that one or two tendons only on the hand have been occasionally affected with the usual characteristic pains, in sympathy with certain states of the stomach;

* It happens, for the most part, that this complaint rather affects the bursæ and tendons than the immediate structure of the joint; and, on this account, the definition does not appear to me correctly stated.

† Clinical History of the Nodosity of the Joints.

and some cases also, in which, the bursæ belonging to the sheaths of the tendons of the fingers have been so knotted, as truly to be in the state of nodosity; and have now and then given slight evidence of gouty inflammation, no other part having at any time been affected. It usually happens that with this change of organization, the particular phalanx which is affected (usually the last) is so distorted by the action of the muscles, as to present a crooked appearance. I learnt, in the history of some of these cases to which I allude, that the gout was a family complaint. The larger bursæ mucosæ are liable to a peculiar form of disease, distinct both from gout and rheumatism; which, in the chronic form, gives rise to considerable swelling produced by the distension of the bursæ; the membrane secreting a preternatural quantity of its synovial-like fluid. These tumours are free from tenderness, and cause inconvenience chiefly by the muscular weakness which is induced in the affected limb. A swelling of one or more of the bursæ sometimes takes place suddenly in certain constitutions, in connexion with the occurrence of gonorrhœa. I shall have to relate a case of this description.

It is scarcely probable that chronic gout can be confounded with the pains and swellings produced near the joints by secondary syphilis. In such affections, the symptoms have a more increasing progress than those of gout; and are also much less influenced, either as to relief or aggravation, by slight external causes, or by diet, or the state of the mind: and it may be added, that the syphilitic characters presented to the eye, and to manual examination, are in general sufficiently marked to be recognized without danger of error.

OF THE TREATMENT.

FROM the history which has already been delivered of chronic gout, the necessity of an extensive application of the principles of pathology for its elucidation must appear as an immediate deduction. It is only by a theory founded on just physiology, and by a practice resulting from careful discrimination, that our footsteps can be safely directed in the present path. In acute diseases, the bold hand of the Empiric, or even some happy effort of Nature, may sometimes be speedily successful; but when the chronic form of disease is deeply established in the system, no pretended universal, or even expeditious method of cure; or treatment founded on a partial application of principles, has any just claim to our regard.

In the practical arrangement of my subject which I have now to offer (using the privilege of some recapitulation), several distinct modifications of the disease may be presented to our consideration.

I. From that original languor of constitution, or prevalence of the nervous temperament, in which, although the gouty diathesis is strong, the powers of the system have been insufficient to produce the acute form of gout; and pain, swelling, and difficult motion, are the symptoms which chiefly appear. The internal functions are weak and unhealthy, and much nervous sympathy is present.

Here our indications will be chiefly derived from the state of the chylopoetic viscera, and from an attentive observation of the several secretions. It sometimes happens that stimulating medicines, and too cordial a diet, have produced, in this constitution, a slight degree of inflammatory diathesis, joined with plethora; but, even

in this case, general bleeding will seldom be allowable. The employment of purgative diuretic medicine in the saline combination, such as before recommended, with the occasional interposition of a dose of hydr. submur. et pulv. jacobii, and the abstraction of all heating stimuli, will usually constitute a treatment of sufficient activity.

The soothing influence of narcotic medicine should be employed at bed-time, for the relief of pain and nervous irritation. Active doses of opium are however both unnecessary and injurious; and indeed some other narcotic will usually be found sufficiently powerful, and therefore more deserving of choice. I have already, when speaking of narcotics in the treatment of acute gout, entered fully into detail; and shall here only state, that, for the relief of these minor pains of chronic gout, I have found great success in the use of stramonium and lactucarium; and also, of lactucarium with p. ipec. comp.; prescribing the latter in very small doses.

In chronic complaint, when we are not compelled by urgency of symptoms to choose the most active forms of medicine, it is desirable to select those which act in a gradual yet effectual manner. As opium restrains the action of the bowels (a circumstance of only slight objection in the treatment of the acute symptoms, when we are at the same time administering active purgatives), we gain a great advantage in the cure of chronic gout, by the use of the weaker narcotics, which have not this effect; and although the extract of henbane is on this account an excellent medicine, and its soothing powers are not to be disregarded, I am, from comparative experience, quite decided in favour of the united influence of stramonium and lactucarium.

The local treatment is to be conducted exactly on the principles formerly described. Inflammatory tenderness

being removed, friction and bandage will be eminently useful.

The more permanent method of practice comprises the improvement of the internal functions, by a plan of medicine, and of regimen, corresponding to the changing indications.

Tonic medicines are usually injurious, and tend rather to fix than remove the disease. Whatever formula we choose, it must be our care to promote the functions of the liver and the kidneys, to correct the morbid processes of disordered digestion, and excite the action of the bowels, without nauseating and weakening the stomach. The *tinctura benzoes composita*, formed into a draught, and given once or twice a day, in conjunction with *magnesia*, is an useful stimulant to the bowels, and at the same time corrects acidity. The draught mentioned at p. 186, in the saline form, administered in the morning early; and one, in the middle of the day, consisting of *cascarilla* or *gentian* infusion, in equal proportions with *decoct. al. comp. et mist. camph.* to which ammonia or soda may be added, comprise the forms of medicine, which, in a general recommendation, I am most induced to mention.

In a course of treatment, it is desirable that the dose, if of fluid medicine, should not be repeated more than twice in the day; but if the bowels do not become sufficiently excited, the use of some purgative pills at bed-time should be added. I may mention, as useful, a combination of the *pulvis aloes compositus*, *pulvis rhei*, et *sapo durus*, with a small addition of the *pulvis jacobii*. As a part of the plan in question, the alterative use of some mild mercurial oxyd is of especial importance. In the summer season, the use of the warm

sea bath, twice or thrice a week, will usually form a valuable auxiliary to the other remedies.

If, when the secretions are rendered permanently healthy, debility of stomach and general depression should remain, we are consequently led more distinctly to the use of a tonic ; but in this choice we must guard against the effects of heating stimulants.

The regimen, in every particular, must be carefully regulated according to the individual constitution ; but daily exercise in the air of the country, and the avoiding of a diet too full and stimulating, are cardinal points of observance.

2dly, The consequence of acute gout, when its repeated invasions have impaired the energy of the constitution ; and from the weakened circulation which is induced, the chronic diseased action alone takes place. The functions of the internal viscera are more or less deranged ; and the nervous system is much disturbed.

In this example, we commonly see that the primary character of constitution remains to so great a degree, that signs of plethora are often manifested ; and slight local inflammation is readily aggravated by the injudicious use of stimulants. Under these circumstances of vascular susceptibility, in conjunction with languid powers, the treatment which has been stated in the preceding example is generally applicable. The alterative aperient plan will, however, sometimes be required to a greater extent, and for a longer continuance. The state of the secretions will be the true guide to the practice which should be adopted. Material information will be derived from a knowledge of the specific gravity of the urine. I have invariably found, that so long as this has been very high,

the more or less active employment of a diuretic purgative has been attended with the best effects*. The draught, p. 186, regularly repeated twice a day, and the compound calomel pill each other night, usually prove very successful in answering this intention of continued evacuation ; but I have sometimes found the action of the kidney to be so deficient, that it has been necessary to join some other diuretic combination with the draught in question. In a very torpid state of the bowels, I have occasionally added guaiacum to the purgative ingredients with advantage. When the practice now mentioned has been sufficiently pursued, a tonic mode of treatment will follow in the order of succession. But in some instances of prevailing fulness of the habit, local congestion is pointed out, and the consequent propriety of relieving

* It may interest the curiosity of some of my readers that I should mention the following formula of *the Chelsea pensioners'* medicine, which, it will be seen, consists of ingredients that are in common use. This is a heating medicine ; and I should consider it an improper remedy in the paroxysm. I am informed by some individuals, that in the intervals of their fits, they have found it agree very well, as a purgative.

“ Take of flower of sulphur, two ounces.

“ Cream of tartar, one ounce.

“ Rhubarb powder, two drachms.

“ Guaiacum, one drachm.

“ Honey clarified, one pound.

“ One nutmeg, finely powdered. Mix them intimately.

“ Two large tea spoonfuls to be taken night and morning, and to be persevered in till the whole is consumed. For the three first nights, a large tumbler of warm rum and water is to be taken at bed time ; or, if fever is present, white wine, instead of rum.”

the loaded vessels, by cupping, becomes manifest. An obstructed circulation in the liver seems to be the common cause of partial determinations of blood in gouty persons; yet the symptoms are considerably varied. When the right or the left hypochondriac region is affected with pain and tenderness, the blood should be drawn from these situations. If pain, heat, and sensible fulness distress the head, cupping at the neck will be an appropriate remedy; but it should be well remembered, that the symptoms now spoken of are of a secondary character; and the permanent cure will be found in the judicious treatment of the chylopoetic viscera, and in the collateral points of practice.

What has already been said of local management, need not here be repeated.

In the various shades of the chronic gout; between that fulness of habit and remaining degree of action, in which the symptoms approximate to those of the acute form, or even sometimes are actually converted into it; and that universal languor and debility, with which the weakest inflammation alone is found connected; the practice must also be accordingly modified.

In this last case, it may be stated, that although the use of stimulant and tonic medicines is indicated, and demands occasional preference, the purgative and corrective intentions must equally be fulfilled, to a given extent. Subcarbonate, or the neutral carbonate of ammonia, will be found a medicine highly useful when stimulant treatment is indicated, and may be favourably joined with the compound decoction of aloes, infusion of gentian, and peppermint water; the alterative pill also being employed.

For the relief of symptoms of a spasmodic and nervous character, which are connected with debility and

morbid sensibility of the stomach, a stimulant antispasmodic medicine may be given occasionally ; such as a conjunction of the aromatic confection and carbonate of ammonia, or æther, with camphor mixture ; but, to palliative remedies of this kind, the patient should not too familiarly resort.

In certain states of debility and general disorder, into which some gouty invalids decline ; or in anomalous cases of disease, where gout has not yet existed, but is suspected ; and when the symptoms are those of excessive languor, or even of partial privation of function with a threatening of general paralysis, it has been a common practice to invite a fit of the gout by various modes of stimulating treatment.

The propriety of this proceeding is certainly as questionable, as its object is difficult of attainment ; and it does not seem warranted by the amount of benefit, which Nature in her spontaneous production of a paroxysm occasionally confers. It is always hurtful, and sometimes hazardous, to excite a weakened circulation into strong action ; and it cannot be denied that the active means which are thus injudiciously adopted to urge the gout, may produce, instead of it, an apoplectic fit. I apprehend that the true method of treatment, on the occasions to which I have alluded, consists in a regular and persevering attention to the chylopoetic functions, both by means of medicine and regimen, according to the principles of which an account has already been offered.

The learned and experienced Dr. Heberden observes on this subject ; “ In complaints of a chronical nature, whatever suspicion there may be of gout, it would be no bad rule of practice, not to direct the waters of Bath, nor any other remedies which are supposed to give the gout, if they would be improper when the same com-

plaints arise from other causes ; but to content ourselves with putting the general health into the best state, by strengthening the appetite and digestion, and by relieving the urgent symptoms*.”

The Bath waters, judiciously employed, are unquestionably a remedy of great value, in some states of the constitution consequent on gout. Authors seem, however, very well agreed, that they should be forbidden when any inflammatory diathesis is present†.

Dr. Parry informs me, “ that the Bath waters, in no form whatever, are beneficial during the paroxysms of gout, or in any inflammatory disposition which may exist in the interval.”

In that dyspepsia of the gouty, which is joined with a languid circulation and a great deficiency of nervous energy, and where neither organic congestion nor inflammatory tendency exists, these waters appear calculated to be eminently useful ; and the praises of authors and of patients concur in justifying the reputation which they have acquired. The cheerful influence of such a watering place ; the repose of mind which is obtained by those who leave the cares of business behind ; the improved regularity of all the habits ; and the change of air itself, which is usually a remedy of no small power ; and finally,

* Commentaries on the History and Cure of Diseases, p. 45.

† Dr. Gibbes, at the same time that he recommends the powers of the Bath water to relieve “ the debilitated state in which the patients are left after a severe fit of the gout,” observes, that “ when the gout is, as it is expressed, flying about a patient’s constitution, the warm bath is highly dangerous. I have heard of very dismal consequences resulting from an imprudent use of the bath in such a state.”—Treatise on the Bath Water, vol. ii. p. 34.

the confidence of benefit with which the mind is inspired ; concur in procuring the relief, and in effecting those cures which are attributed to the waters of Bath.

I have collected from patients many instructive reports of the effects of the Bath waters ; and shall recite a brief abstract of those which are most interesting.

A gentleman, aged 41, whose constitution was much debilitated by frequent attacks of gout, and who possessed entirely the nervous temperament, suffering in the intervals of the paroxysm chronic pains and much weakness of limbs, pursued a course of the water both by its external and internal use for three months, in the summer season. The *hot bath* occasioned at the stomach both a remarkable sensation of weight and of coldness. The *Cross bath* did not produce any such effect, but its continued use made him feel more weak and languid than usual ; and he quitted Bath without having derived any ease to the limbs, or received any kind of improvement.

A gentleman, aged 28, of robust appearance of constitution, in his second fit was attacked in the great toe of the left foot. Leeches were applied freely, with relief. In three days the right ankle was affected, and leeches were again used. The inflammation was scarcely in any degree relieved, and very quickly the right knee was seized. A large number of leeches was now applied to this knee, and with apparent advantage ; but the left was as immediately attacked, and he describes that all the parts thus treated with leeches, remained extremely weak and subject to chronic pains for nine months, at which period he went to Bath. There he bathed, and received also dry pumping (so called because, in the use of it, the affected part of the body only is wetted, whilst the rest is kept dry) on the weakest parts, for a course of thirteen weeks ; at the same time

regularly drinking the water. He returned completely cured.

A lady, aged 54, of strong constitution, but of the nervous temperament, suffering chronic pains and weakness of limbs, and a spasmodic state of stomach from slight causes, entered on the usual course of the waters. From the unfavourable determination of blood to the head, and consequent confusion which it occasioned, its internal use was soon discontinued. General immersion occasioned languor, and increased the general debility; but dry pumping, both on the stomach and on the weakened limbs, produced the best effects.

A gentleman, aged 33, of the nervous temperament, and not of strong constitution, having often suffered severely both from gout and rheumatism, and being afflicted with the pains of chronic gout, which had been induced by exposure to cold after the use of the hot bath, went to Bath, and entered on the regular course, drinking two rummers a day, and bathing three times a week. He pursued this plan about three months, omitting all medicine, except some ordinary pills to regulate the bowels. Previously to drinking the water, his urine had deposited a pink sediment very copiously; but after a few weeks he seldom noticed this appearance. He found his appetite and spirits much improved. With respect to the local effects of bathing, he described that on those days he had less stiffness and aching than usual; but the amendment was only transient. He used a temperature from 97° to 100°. He quitted Bath without having received the least material improvement.

When this gentleman came under my care, soon after this period, I found very manifest indications of an unhealthy state of liver, and adopted a plan of treatment accordingly. The ankles were thickened at the joint; the

integuments were œdematous; and one knee was much enlarged from bursal distension. All the muscles of the lower extremities were small, relaxed, and very weak. Extreme lameness, with much awkwardness of gait, was the consequence. I recommended the usual plan of salt water sponging to the whole limbs, a stimulating liniment to the weakest parts, and night and morning a systematic method of friction, with speedy good results. The case terminated in the most favourable manner.

A gentleman, aged 55, robust and plethoric, first attacked with gout at the age of 29 (the disposition not hereditary), suffered a severe paroxysm in the beginning of autumn, which was regularly and successfully treated. In September, being quite convalescent, he went to Bath, as it was his occasional plan to do. He favoured me with the following statement: "After the usual preparation by aperient medicine, I commenced the drinking of the water of the Cross bath, with one glass of the middling size before breakfast, and the same quantity before dinner. It agreed with me as usual on former occasions, always giving me an excellent appetite, and an extraordinary flow of spirits. At the end, however, of eight days, I began to feel the approach of gout very sensibly in the feet, and in short was quite lame. My physician considered that the water was too stimulating, and advised its discontinuance. I should remark, that I was not sensible of any fever, and did not notice the usual discolouring of my tongue, nor the appearance of the pink sediment in my urine during this attack. After, however, the swelling of the ankles had subsided, I was still distressed with flying pains about them and my feet. I was next recommended to try the effect of the King's bath, and not to think of the internal use of the water. I bathed in consequence every other day, and finding the

plan agree, and that the pains in my feet sensibly diminished, I continued it regularly, five weeks ; and the result was very satisfactory indeed."

The following concise reports serve to shew the necessity of duly preparing the system, in order to the favourable employment of the Bath water.

C. D. aged 52, of the nervous temperament, first affected with gout a year ago, having been under a short course of aperient medicine, and appearing to be sufficiently prepared, made cautious trial of the water ; but after a few days, he experienced feverish excitement, obstruction of bile, and uncomfortable feelings of the head ; so that it was necessary to return to the former plan. This being pursued a sufficient time, he resumed the use of the water with much advantage.

R. S. aged 66, had been subject to gout thirty-nine years, and lately suffered in a great degree from nervous depression, weakness of stomach, and general languor. He first drank the water, having only premised the use of an aperient draught. It disagreed, causing uneasy feelings in the head. He was then cupped, and the result was so favourable, that he resumed the water, and drank it for three weeks with great benefit.

H. A. aged 50, gouty for ten years past, of bilious diathesis, labouring under weakness of stomach and many nervous feelings, entered upon a course of the water. A very confirmed jaundice soon took place, which was imputed to the injudicious employment of the water.

W. R. aged 45, of the mixed temperament, plethoric, first affected with gout at thirty-five. Without previous preparation, drank the water for ten days ; his sight became dull ; he declares that he felt senseless from the confusion of his head, and also experienced great lassi-

tude. All these symptoms yielded to a course of alterative and purgative treatment.

R. V. aged 51, of the mixed temperament, extremely susceptible to the influence of stimuli; first affected with gout at twenty-six; having a weak state of stomach, and much nervousness, visited Bath and drank a second size tumbler of the water three times a day, for several weeks. He was rendered more indisposed in every respect. He became affected with palpitation of the heart and in the epigastric region, alternately; and also with faint perspirations. He then went to Cheltenham, and took the water No. 4. At first, it caused sickness; but, with perseverance, he found it agree; and gradually all his unpleasant symptoms disappeared.

A gentleman, mostly of the sanguineous temperament, many years subject to gout, found so much excitement from the Bath water, that during the whole course of it, he could not take even half a glass of wine without finding himself inconveniently heated and stimulated. Observing, however, every precaution, he derived benefit from the water.

I am acquainted with many instances in which the chalybeate water of Tunbridge Wells has, after a short course, excited an attack of gout. The patient finds his stomach sensibly strengthened; and, from its early effects, promises himself great advantage; but, ere long, he is overtaken by the sufferings of gout.

A gentleman relates to me, that being in doubt whether on a former occasion his complaint had really been the gout, at a subsequent period, when suffering from dyspepsia, he took this water as an *experimentum crucis* with regard to the gout; and soon after, a decided paroxysm followed.

In most instances, and indeed almost without exception, the waters of Cheltenham prove highly beneficial to gouty persons; and particularly when joined with a medicinal alterative, and with regulated regimen. It not unfrequently happens, that the influence of their stimulating properties predominates, and that they have the effect of exciting a paroxysm; but I do not know of any tedious fit resulting. The water, No. 4, is that which is most suitable to the gouty patient, and especially in the first instance. I mention this by way of adding, that no other kind should be attempted without previous advice.

The waters of Leamington are entitled to similar praise with those of Cheltenham; but I learn from a comparison made by many persons, that their aperient effect is less active. From the waters of each place, it now and then happens, that flatulency and distension ensue in such a degree, as to require that their use should be abandoned; but in other instances, these effects disappear as soon as their operation becomes well established.

In reasoning upon the composition of the sulphureous water of Harrowgate, I have always been induced to consider it as a very fit summer alterative for gouty persons; and for those, who, together with the gout, suffer any inconvenience from cutaneous affection, it particularly recommends itself. I collect a favourable general report of the effects of the water from several gouty patients who have visited Harrowgate; but I must add, that some have found its influence such as to excite an attack; they being, as I imagine, much charged with the gouty diathesis at the time. One gentleman observes, that he took two glasses a day, and used the bath each other

day, remaining in about five minutes. In a fortnight after he was seized with gout, and it proved to be the most severe fit which he had ever experienced.

Another gentleman considered it to be in an equal degree curative. Having recently had a fit, he took three large glasses daily, which acted most favourably. The gout went off gradually, and he regained perfect health.

Some, under my observation, have made use of this water at home, having imported it in bottles of convenient size. One of these patients observes, that two half-pints daily, affected the bowels in some degree, but acted more remarkably in causing perspiration. I may here notice, that I have ascertained by examination, that the Harrowgate water bears removal, and can be kept a considerable time, in sealed bottles, without material deterioration.

Finally—of this water, as of the waters of Cheltenham and Leamington, I have to observe, that in certain constitutions, occasionally it fails to act as an aperient, and causes disagreeable distension and flatulent oppression. The operation of all the waters depends in a great degree upon management; and usually some alterative pills should be joined with their use.

A course of any of the mineral waters which have been now mentioned, with some exception that may be made in regard to Bath, obviously demands a favourable time of the year, in order that their effects may be aided by suitable exercise in the open air. It happens on most occasions, that the gouty patient requires an alterative aperient course of medicine, at a season, when the visit to a watering place is not to be prescribed. It follows then, that we should pursue our plan of treatment at home; and indeed when the gouty diathesis has been evidently strong, I have preferred, under any choice of circum-

stances, the decided activity and efficacy of medicines, to the use of mineral waters. When the patient has been suffering rather from oppression of his powers than loss of strength, in consequence of the influence of a morbid state of the digestive organs (to use a comprehensive expression), I have avoided tonics, and in my usual treatment, have prescribed an alterative pill on alternate nights; the draught, p. 186, in the morning early, every day; and a course of ass's milk, half a pint being taken about half an hour after the draught. If languor prevail, I prescribe in the middle of the day the second draught described at p. 266, using either cascarilla or gentian as circumstances suggest. Regimen is of all possible importance.

The most difficult and untractable cases of chronic gout which I have seen, have been those in which the empirical treatment (whether in the use of the eau medicinale, Wilson's or Reynolds' tinctures, or even the vinum colchici when trusted to as a specific) has been employed; and I have found a disposition to relapse implanted in the constitution in proportion as such treatment has been long continued. It often, therefore, requires much perseverance in the use of regular methods of treatment, before this habit of relapse can be overcome. In the acute paroxysm, the physician sometimes experiences the disadvantage of an impatience in the mind of the patient, from the slower progress which he makes, contrasted with the rapid apparent cure which he has before obtained by one or two doses of his favourite nostrum. But in the chronic gout we have the same evil in another shape. The nostrum has produced the tendency to quick relapse; and has proved how impossible it is to cure, in the true sense of the word, a disorder which has been slowly arising out of the faults of the constitution, by one or two doses of a supposed specific medicine.

I have the satisfaction to add, that I have, in an extensive number of cases, succeeded most satisfactorily, after a time, in gaining the ascendancy over the state of constitution which I have now described; finding that the disposition to gout, by degrees, lessens as much as before it increased. This, indeed, may be expected from the united influence of the corrective and aperient plan of treatment, in conjunction with regimen.

3dly, I have to consider a chronic state of gout also arising out of repeated acute attacks, in which, local changes of structure have been produced in a great degree; the nervous system is highly sensible to the influence of external causes; but the state of the natural functions proceeds for the most part in a healthy course. I have in this description presumed that the energy of the constitution is retained in considerable vigour; and that the injury which the attacks of gout have produced, affects chiefly the limbs. With this form of chronic weakness rheumatism is often blended, and the patient is extremely susceptible to every vicissitude of weather, and especially to wet, and to damp cold air.

Sometimes he rises seemingly strong and comfortable; but after a little exercise, especially if the weather be unfavourable, his joints become painful, affected with a sense of heat, and almost disabled; together with aching of the loins, and general lassitude and weakness. The symptoms are very various in different cases. Some, by the exertion of a walk, dismiss the symptoms with which they rise in the morning. One gentleman informs me, that on some occasions he goes to bed with such threatenings of gout, he expects a fit by the morning; and instead of this, he is so much relieved and alert, that he is able to hunt.

In the pains and frequent threatenings of inflammation, which, under these circumstances, continually occur, the combination of a narcotic and a sudorific appears particularly useful; and for this purpose, the compound powder of ipecacuanha, in small doses, twice or thrice in the twenty-four hours, often proves a valuable medicine, strict attention being at the same time paid to the proper action of the bowels and kidneys. In this form of the complaint, however, I must not lose sight of the recommendation which is due to the combination of stramonium and lactucarium, from which I have repeatedly seen the best effects derived.

The improvement of the diseased and weakened limbs claims our particular attention.

It is in aggravated cases of the kind which I have now in view, that a system of tepid bathing will usually prove most useful; and that *Buxton* promises its greatest benefits.

So far as I can form a judgment on the subject, I should much prefer, as a remedy for certain states of chronic pain and weakness, the trial of Buxton to that of Bath; and this opinion concurs with the result of my whole experience, that a tepid temperature is always useful, and a high temperature is generally injurious to gouty parts. I was consulted by a gentleman crippled with gout and rheumatism, who, after fruitless trials of the hot bath, persevered for several months in the use of the vapour bath; but the limbs became weaker, and in no respect relieved. To him, morning sponging with tepid salt water, and diligent friction with a liniment moderately stimulating, have rendered great service.

From much experience I can confidently recommend this method of treatment, under circumstances of previous relaxing means, both as tonic and preservative in a high

degree. Simple rubbing will usually prove sufficient, except where the energy of parts is very defective; and in that case, stimulating liniments will much assist the effect of friction.

In returning to the subject of Bath and Buxton, I shall offer the following quotation from Dr. Heberden, respecting Bath* :—

“I have not been able to observe any good in arthritic cases from the external use of these waters, either when the distemper was present, or in its absence: on the contrary, it has rather appeared to increase the weakness of the limbs; and sea bathing has contributed far more to recover the strength of gouty persons; many of whom, in the intervals of their fits, have used it with safety and advantage.”

That the employment of sea bathing by a gouty person may sometimes be made with great advantage, is a truth that may be at once admitted; but I would restrain the prescription within very narrow limits. It appears to me that the plethoric state or tendency, which is so common to the gouty, should always be a point of careful consideration, in determining the propriety of using cold bathing. The circulation suddenly checked from the surface, may be unfavourably directed; and in proportion as plethora, or local congestion, or immediate tendency to gout, may be existing, the interference of so strong an agent as cold immersion must abound with hazard. Also, if the energy of the limbs be very deficient, the application of cold is not followed by sufficient re-action on the surface; and pains, as of rheumatism, may be produced.

* Commentaries, p. 51.

A gentleman, aged 36, having with advantage used the warm bath three times a week, for five weeks, to relieve some chronic pains and weakness left in the limbs by a severe paroxysm; in the hope of finding a greater tonic effect from cold bathing, went into the open sea, with the interval of a day, repeatedly. He now experienced occasional pain at the pit of the stomach; tingling sensations in the ribs; some aching in the sciatic nerve of one limb, and, he says, "gouty pains flying all over him." He resumed the warm sea bath, and it agreed perfectly.

It is to be observed, that the temperature of the open hottest bath, at Bath, ranges from 108° to 100° ; of the Cross bath, from 98° to 94° ; and the temperature of the Buxton bath is stated to be 82° . Hence the action of the two waters, as remedies in the circumstances in question, will be widely different. Some illustration of the power of the Buxton bath, in relieving a weakened and painful state of limbs induced by chronic gout, will not be uninteresting.

A gentleman, subject to rheumatism, was seized with gout in the great toe, only, in the first fit. In subsequent attacks, both toes and both ankles were affected. He had also sciatica on one side, and rheumatism in various parts, in connexion with the last paroxysm. All inflammation being removed, but harassed with continual pains, and such weakness in the joints, that he had the apprehension of losing the use of his limbs, he made trial of the common warm bath. It did not afford relief, and served only to increase the debility of the limbs. In this state he went to Buxton, and bathed regularly for seven weeks, with the happy result of a perfect cure.

Another gentleman, much crippled from chronic gout, and also affected with rheumatic pains, received a cure

at Buxton in five weeks. He relates that he was not sensible of any material advantage, until about the fifteenth time of bathing.

A third patient had suffered severe gout in both feet, in the toes, insteps, and ankles. He had applied leeches freely to the inflamed parts, with only slight relief; and considers that they led to the very serious œdematous weakness which he suffered long afterwards. At a distant period from the paroxysm, he was unable to walk for more than a quarter of an hour, without producing swelling of the feet, and much fatigue and excessive aching. In this state he visited Buxton, and by regular bathing, in rather less than a month, his limbs received a cure. I am acquainted with cases in which the external and internal use of this water has, in about a week after its employment, excited a paroxysm; doubtless the gouty diathesis having been in particular force at the time.

In the way of external treatment, Dr. Saunders has stated his opinion, that it is the *temperature* alone which can fairly claim the praise of being a remedy. If this idea be as correct as it seems reasonable to suppose that it is, a domestic plunging bath, affording the convenience of heated water, would be valuable to the gouty patient; and thus the imitation of Bath or Buxton, or the employment of any intermediate temperature according to the circumstances of the case, would be in the possession of those whose affairs cannot permit their absence from home. I am aware that this suggestion may appear more plausible than practicable, from the great expence which is attached to the construction of such a bath, with the suitable arrangements for warming the apartment; but I offer the hint to the affluent.

It must be the patient's care to supply the many collateral advantages of the watering place, by a correct

regulation of all his habits. The superiority which a spacious tepid bath affords over one that is confined, is obviously to be referred to the free motion, which allows the opportunity of using a low temperature during the immersion. It is, in my opinion, much to be regretted, that our sea-side watering places do not possess the convenience of a *plunging* tepid bath. In certain states of chronic rheumatism, and also of chronic gout, the invalid would, I conceive, derive still more benefit from the stimulus of the salt water with a proper temperature, than even from the Buxton bath, and would be spared the trouble of a long journey. Such were the sentiments which I formerly expressed; and they were derived in part from theory, but mostly from the reports of those who had from experience drawn a comparison between the bathing obtained at Buxton and Bath, and other places; at the sea-side especially. But candour and more recent information require me to confess, that, in great probability, I have not done justice to Buxton in the sketch which I have before drawn. It happens more particularly in chronic rheumatism, than in chronic gout, that the bathing at Buxton proves so remarkably useful. According to the report with which Dr. Drever (whose experience on the spot has been extensive) has favoured me, the water of Buxton, used both internally and externally, has a specific power of acting which is not to be referred to mere temperature; and he assures me, that it has, in many remarkable instances, led to a cure, when every other mode of bathing had entirely failed to give relief.

I was consulted by a gentleman, who, together with chronic gout affecting the knee, was much distressed by wandering rheumatic pains. The latter yielded entirely to the Buxton bath; but in regard to the former complaint,

he did not profit by its influence in any degree. Did not this result exemplify, how much less the rheumatism is necessarily connected with the constitution, than the gout?

Under the impracticable circumstances of procuring for my patient the advantages of a tepid plunging bath, I have on some occasions directed a shower bath, consisting of a saturated solution of salt, and raised to a proper temperature by the addition of hot water, to be allowed to fall on the affected limbs; and the result has been always more or less satisfactory. In this plan I have insisted on subsequent friction being very freely employed, as an important adjunct of the treatment. I think it material to suggest, that we should view any mode of bathing which may be adopted, as principally in the light of external treatment, and as a valuable auxiliary to internal means of management. I have seen some striking instances in which a course of bathing has alone been trusted, and the treatment of the internal organs having been neglected, either no material advantage has been procured, or a relapse has shortly followed. These observations are applicable to what I have found to result from the common warm bath, from Bath, and also from Buxton.

A lady entirely of the nervous temperament, after a severe paroxysm, made trial of a warm *sea* bath, with a view to improve the limbs. Her digestive organs were still disordered, for no proper treatment had been pursued. The bath produced so much sense of tightness in the chest, and uneasiness of breathing, that she could not continue its use. It appeared comfortable in its effects to the weakened limbs.

A. S. aged 50, long subject to gout, when suffering from weakness of the limbs, and pains which appeared to be rheumatic, made trial of the hot bath at Brighton.

After bathing a few times, a paroxysm of gout took place, which proved severe.

T. L. aged 52, first gouty at 35, experienced so much sense of debility and perspiration from using the hot bath once only, that he could not venture to repeat it. His muscles were previously relaxed, and his constitution generally was so deficient in tone, that it was an unfit remedy. He adopted it with a view to relieve some chronic pains of his limbs.

Another gentleman, under similar circumstances, found exactly the same effects.

R. S. aged 30; gouty since 24; when suffering from the sequelæ of a paroxysm, and affected with pains of the limbs, which appeared to be rheumatic, after a few times using the warm bath, found himself rendered feverish, the weakness of the limbs increased, and the pains not relieved. These examples are sufficient to shew the importance of considering carefully the state of the constitution previously to the use of so simple and innocent a remedy as the warm bath appears to be.

When proper constitutional treatment has been premised, the use of the warm sea bath often proves a most successful remedy; but a greater perseverance may be necessary than the patient is prepared to expect; and I have known it happen, that although the remedy has agreed, no apparent benefit has begun to appear, until two or three weeks have elapsed; and yet afterward the best effects have been obtained.

Of the good effects of the Leamington warm bath; the following is a pleasing instance:—A gentleman having suffered gout severely, was left very lame, with much stiffness in the joints, and rigidity of the muscles. After bathing two or three times, his improvement was so satisfactory, that he walked almost without difficulty.

The vapour bath is a remedy of considerable power, and should not be inconsiderately employed. I have known some gouty persons derive much benefit from it ; and others have experienced inconvenient languor, and increased susceptibility to the changes of the atmosphere from the great perspiration which it has induced. This result has happened most to corpulent persons. In proportion as the solids are relaxed, and the muscular fibre is weak, I should dissuade the patient from the trial of the vapour bath.

In those cases of chronic gout, in which, from neglect or *mal-treatment*, the flexor muscles of the limbs have so much overpowered the extensors, that a permanent contraction of the muscular fibres, with a rigid and contracted state of tendons, becomes the consequence ; a distinct mode of treatment is rendered necessary.

When I reflect on the examples of miserable lameness, of youth rendered almost helpless, and the middle age of life made decrepid with infirmity, which have repeatedly come under my observation as the consequence of neglected or improper treatment of acute and chronic gout, I feel that I cannot too impressively recommend an attentive professional management of the paroxysm ; or dwell too much on the importance of the method of practice, of which I am now about to speak, when such management has been neglected, and the consequences just described have been produced.

We find in some of these cases that the just antagonist action of the muscles is quite destroyed ; and the patient, if able to walk without the help of crutches, moves at right angles with his arms and legs, awkwardly bending forwards to lessen the superincumbent weight on the knees and ankles. He also walks chiefly on the heels, and with the appearance of fear and uneasiness. The

extensor muscles are found relaxed and small; the flexors also small, but rigid and contracted. The legs, and particularly the feet, are much affected with coldness, and the languid circulation in the extremities is often apparent from the blueish colour of the skin. The bursæ, mucosæ in different situations are distended and tender, and occasionally are much enlarged. This happens chiefly at the ankle joint, around the knee joint, in the ham, and at the tuberosity of the os ischium. The ligaments are thickened and contracted, and very tender; the tendons are rigid and hard; and from these causes, according to the degree in which they prevail, the lameness is general or partial; and according to its duration and other circumstances, is very tractable to attentive management, or admits only of some amelioration.

The usual modes of bathing, either warm or tepid, seldom afford any advantage, and in some instances appear productive of increased infirmity. It is indeed in the rubbing process, which was first instituted by Mr. Grosvenor of Oxford, that the only method of cure for these cases can possibly be found. It is in vain, as I have seen in three examples, that the powers of electricity are employed upon the languid limbs, or blisters, or any stimulants. I have the satisfaction of stating, that in many very urgent cases in which various other active means had been unsuccessfully tried, I have procured the most decisive advantages from instituting this treatment*. It will be readily imagined, that much time and perseverance are required to give due effect to the plan;

* In this metropolis there are persons properly instructed in this process; and who attend on very moderate terms of remuneration.

and I must also add, that most commonly in addition to it, internal means of treatment are indispensably required. The regular plan of friction will be found usefully assisted by morning sponging with tepid salt water; by the use of rollers when œdematous swelling, or a varicose state of the veins are attendant evils; and where a want of energy much prevails, by the use of a liniment at night; or night and morning.

I must in this place observe with regard to rollers, that, if ligamentous rigidity be the chief inconvenience which exists, they prove rather injurious than useful; by restraining, in some degree, the sphere of motion in the joint, which is already too limited.

I was favoured, two years ago, with the following interesting communication by Mr. Davis, who having received the early part of his professional education from Mr. Grosvenor, is highly competent to appreciate the value of the method of practice now under consideration:

“ In May 1815, A. B. came to town in consequence of having had the gout repeatedly during the last year and a half, in both knees and ankles, which had produced an enlargement and distension of the bursæ mucosæ of the knees, a contracted state of the flexor muscles of the limbs, a peculiar rigidity of their tendons, and an enlargement in one knee below the patella on each side, giving a *sensation* to the touch very much resembling that of cartilage. There was a great degree of weakness and stiffness of the ankle of that limb, with a considerable thickening about the tendo-achillis, at the point of its insertion into the heel.

“ The patient, from having totally lost all power of moving the knee joints, was unable, without the utmost difficulty, to move about the room, even with the aid of crutches.

“ Manual friction was applied regularly for one hour and a half in the morning, and again for the same length of time in the afternoon, to the lower extremities, particularly from above the knees downwards. This produced a gradual improvement, and after the continuation of this plan during ten weeks, the weakened limbs acquired their usual tone and powers of motion; and the joints assumed nearly their natural appearance.

“ Previously to the patient's having recourse to friction, he had drunk the Buxton waters, and used the warm sea bath for several weeks, without the smallest benefit.

“ During the time the friction was applied, he had no return of gout; and since that period, July 1815, he has had only very slight attacks.

“ He was first attacked with it at the end of the year 1813; and till that time had enjoyed robust health, and was possessed of great activity.

“ The paroxysm, which chiefly led to the lameness in question, arose from his exposure to cold and moisture for several hours, after having been over-heated by violent exertion.

“ The acute stage of the disease lasted several weeks, during which time he was kept in bed, and subjected to the influence of stimulants, with his joints enveloped in flannel.

“ On the abatement of the gouty inflammation, the joints of his lower extremities remained very weak, particularly the knees; and upon the least exposure to cold, or upon any irregularity in diet, there re-appeared some slight gouty inflammation, which in the end reduced the patient to the situation already described.

“ Here then is a striking instance of the beneficial influence of friction, properly applied to parts of which

the energy and action had been destroyed by the gouty inflammation. I say *properly* applied, because I have had reason to believe, that if the friction be used before the inflammation has subsided, or if it be applied too violently, it brings back the gouty inflammation, instead of producing the salutary effects above mentioned."

In my own experience with the effects of friction, and this has now been very considerable, I have found it essentially necessary to join with the plan, an alterative course of medicine; and, without which, gouty inflammation would be certainly excited. Indeed the local weakness is sometimes so nicely balanced between simple debility, and tendency to inflammatory action, that much caution is required in the use of friction and stimulants.

In many instances it is more prudent to premise the use of medicine for a short time, to the plan of friction.

The patient must be apprised, that for a day or two, the parts which have been rubbed, will have the feeling of being bruised, and probably the general lameness will be increased; but these consequences are only of a transient nature, and are not to be regarded.

In the list of cases of chronic gout which I purpose to offer, I shall relate some instances of the success of friction, in restoring the patient from the most discouraging condition of lameness; the local plan being assisted by corrective medicine. To these cases, therefore, I refer the attention of my reader.

In minor cases of complaint affecting the limbs as the consequence of gout, more partial and also more simple treatment will prove sufficient.

When the bursal distensions are tender and painful, the application of soap plaster, spread on leather, is much to be recommended in addition to moderate bandage. In those gouty enlargements around the joints, which to the eye appear as the growth of bone and cartilage, but are

really produced by the morbid changes of the ligamentous, bursal, and tendinous textures, and a distended cellular membrane, the use of blisters might appear to be indicated; and they have been recommended by several authors*.

In one case, corresponding with the description which I have just stated, I made a full trial of this treatment; blistering the parts several times in succession. The result was much less favourable than the united means of sponging, friction, bandage, &c. employed in other cases; and it must be allowed, that the irritation of a painful remedy (in gout especially) should be avoided, when it cannot be compensated by considerable advantage from its other effects.

In the foregoing arrangement of my present complicated subject, I have entirely excluded those anomalous forms of sympathetic disorder, in which many slow and varying symptoms take place in persons who have never had any evidence of gout; and in whom, from suspicion alone, or loose analogy, and from a desire of removing the appearance of obscurity, by imposing a name, the appellation of gout has been given to the disease. Having before stated my objections on this head, I proceed to the detail of a few cases in further illustration of the theory and treatment of chronic gout.

CASE I.

M. P. aged 60, tall, robust, has a circular chest, is very corpulent, and a perfect example of the plethoric

* Musgrave, &c. Dr. Rush speaks of the employment of blisters both in acute and chronic gout, with much approbation.

habit; of sanguineo-nervous temperament; irritable disposition; accustomed to all the luxuries of a life of indulgence; and with a full diet, uses only passive exercise, although he divides his residence between town and country. Gout unknown in his family. He was first seized at the ball of the great toe, at the age of twenty-four. He has occasionally had feverish attacks, with sore throat; but gout has been almost his only disease. It has scarcely spared any part of the upper or lower limbs; and has produced such a weakness of ligaments, rigidity of tendons, distended bursæ, and thickened aponeurosis, that, in the feet especially, permanent tenderness and lameness are established. To some of his acute attacks, improperly treated, alarming œdema has succeeded; and harassing perspirations have worn the powers of the system, and seriously interrupted his convalescence.

In this gentleman, an acute fit, of late, seldom happens; but chronic symptoms, which now and then have a transient violence, return at no distant interval in the year; and, though inflicting less misery than formerly, considerably embitter the enjoyment of life.

I visited this gentleman in June 1814. The knee was the part most affected. There was much bursal swelling; the skin was tightly stretched, very tender to the touch, inflamed, but not discoloured. He had no ability of locomotion. The feet were a little swollen, and not wholly free from inflammation, and were troubled with shooting pains. He complained of lumbago; and stated that this attack, which had now been of some duration, was excited by exposure to cold. The skin was temperate; but the pulse was 100, with an irritable beat. The tongue was furred, a feverish thirst was present, and

all relish for food was lost. The urine was scanty, high coloured, and depositing much pink sediment ; and the bowels were inert. He had taken occasional small doses of calomel, but without the addition of purgative medicine, so that no effectual benefit had been obtained.

The following draught was directed three times a day :

R Magnes. ℥ss.
 Aquæ menthæ viridis ℥x.
 Aceti colchici,
 Syrupi aurantii āā ℥j.—M. Fiat haustus.

The lotion, p. 254, was applied constantly through the day ; and at night the affected knee was covered with soap plaster. The diet was chiefly diluent for two days ; but afterwards, solids and a little wine were permitted. The bowels became actively excited ; the kidneys soon secreted abundantly ; and, at the end of a week, he was so well recovered, that he undertook a distant journey with but little inconvenience.

With a view to the permanent health of this patient, a more attentive regimen was necessary than he would contemplate with satisfaction, or was willing to obey. He acknowledged his present cure with entire approbation, and promised to observe many good rules ; which I fear, however, were only imperfectly followed. The secreting functions doubtless required watchful attention ; and, as in every other instance of a plethoric habit, all the best means of prophylaxis were imperiously demanded. I am unacquainted with any late particulars of this case ; but I learnt that the gout did not soon return.

CASE II.

February 10th, 1815.—As an example of the chronic symptoms of gout connected with serious visceral derangement, the following detailed particulars appear to me instructive:

A. L. aged 49; a publican; of the middle height; circular chest; very corpulent, with bloated countenance; of sanguineous temperament; extremely plethoric; veins large and full, and in one leg particularly, varicose, with some blotches in the skin; a very free liver both in eating and drinking; his liquors, porter, ale, and gin. Has had gravel occasionally, and is much subject to painful hæmorrhoids. His first attack was at the age of forty-two; and he remained exempted from a return for six years. During this period he did not attend to regimen; but states that he had a regular state of bowels, and very frequent hæmorrhoidal discharge. *Since*, both these circumstances have been much reversed. Gout unknown in his family. The second attack, twelve months ago, affected the same knee as before, and that part only. Six months since, the ball of the great toe in the right foot was severely attacked. In this last fit, first the right foot, then the left, and lastly the right foot again, after appearing to be recovering, have undergone severe gouty inflammation. The fit had been excited by walking, insufficiently protected, on wet ground, and remaining many hours with damp stockings*.

* This patient does not remember to have had warning sensations of a fit; but on the contrary has usually gone to bed on

Both feet are œdematous; the skin is yet slightly red; but this colour vanishes for some little time on pressure; and he is now rather troubled with aching than severe pain. He can with difficulty walk. He is occasionally threatened with a return of active inflammation, having throbbings and prickings in the parts; and cramps in the legs, which harassed him much at the commencement of the acute symptoms, are still troublesome.

The pulse is full, beats strongly, and in frequency 90. The skin is rather hot, and very dry; the tongue is much furred; appetite is lost; the bowels are irregular, but for the most part costive, with foul, blackish, and slimy fæces.

He complains of tenderness, on pressure, in the right hypochondrium, and at the epigastric region, and more especially just at the ensiform cartilage; but I do not discover any visceral enlargement. The urine is scanty, and immensely loaded with pink and mucous sediment. It reddens litmus paper; it contains an excess of urea; is not rendered turbid by nitric acid or heat; its specific gravity 1.028 (see Experiment.iii. p. 120). Four ounces afforded of phosphoric acid 9.2 grains.

In addition to this derangement of the natural functions, and morbid state of secretions, he has a violent cough with oppressed breathing, from which he has suffered many weeks. Expectoration is copious, but does not give relief. The chest is tight and painful, and has the sense of much weight. If he stoops, he is constantly

the night of attack with the ordinary feelings of health. He states, however, that for some time previous to a fit, he has perceived an increase of abdominal corpulency; has had a costive state of bowels, and a deficient secretion of urine.

seized with an alarming spasm across the diaphragm. On the evening which preceded this fit of gout (invading him in the night), in a paroxysm of coughing he fell on the floor, and for several minutes was insensible. He rejected from the stomach much acid and bilious matter, with relief. Now, on coughing, he frequently becomes black in the face. Each night he is distressed with night-mare and frightful dreams; and in the day is sensible of much fulness in the head, with occasional pain. A severe state of hypochondriasis is present.

In this case, every indication presented itself of an overloaded circulation; demanding active depletion. I shall offer a brief recital of the treatment which was adopted. No remedies had hitherto been used.

℥xvi of blood were taken from the arm. Two grains of calomel, the same of antimonial powder, and eight of colocynth extract, were directed each other night, and the draught, p. 186, twice or thrice in the day, accordingly as the bowels should be affected. In the local treatment, I was contented that each morning the skin should be well sponged with tepid salt water. The diet wholly unstimulating, and to consist of milk in any form, vegetables, and puddings.

Immediate relief was felt from the loss of blood, which was sisy and cupped; and evident advantage also followed the purgative and diuretic action of medicine. Three days after, all the signs of fulness being still urgent, the bleeding was repeated to the same extent as before; and, the cough being violent, a large blister was applied to the chest. The medicines before prescribed were continued till the end of ten days.

The gouty pains of inflammation quitted him within a few days; but œdematous weakness, occasional aching, and transient shootings, remained troublesome. The

system having now been considerably relieved, I judged it right, with a view to the strengthening of the limbs, to employ a moderately stimulant liniment, friction and rollers, in addition to the morning custom of sponging.

The kidneys, which at first seemed sufficiently affected by the medicine, now required more excitement by means of stronger diuretics. Ten drops of the tincture of digitalis were added to the draught. In the pill, half a grain of extract of elaterium was substituted for the antimonial powder.

At the expiration of a fortnight much relief was obtained, but many symptoms also remained. The oppression of the chest was sensibly lightened; but the fits of cough, although less frequent, were violent, and the head was full and uneasy. *Nightmare* was abated. $\bar{x}iv$ of blood were removed by cupping at the neck. An opiate squill mixture was prescribed to be taken occasionally. The gums being rather sore, mercurial medicine was suspended. The action of the bowels and kidneys was continued by the effects of the draught twice a day, and by a pill containing two grains of digitalis, a quarter of a grain of elaterium, and half a grain of opium, each night. The appearance of the *fæces* improved, and the urine was now and then light and clear, and of lessened specific gravity; but occasionally also it returned nearly to its former state.

At the beginning of March, the patient's situation was materially different. Appetite and the relish of food were fast returning. Sleep became for the most part tranquil and refreshing. Cough only occasionally urgent, and the breathing easy. The complexion was strikingly changed from a dark yellowish hue, to returning healthy clearness. Much amendment of spirits. The secretions were most favourably changed, but were

not arrived at the just standard. Tenderness was still felt in a slight degree, at the epigastric region towards the right side. In reference to this, 3ss. ungu. hydrarg. fort. was directed to be rubbed in upon the part, each night, until the gums should again become just sensibly affected. An aperient bitter was prescribed twice a day. Light animal food, with table beer, was permitted each other day, in compliance with the urgent entreaty made for restorative diet. The limbs were at this time quite recovered.

In ten days the gums were a little tender, and the mercurial friction was again suspended, as also all other medicine. Air and exercise were advised each fine day. In another week the whole treatment was resumed.

At the middle of April, he was recovered, and had the look, together with the feelings, of cheerfulness and health.

I now made a comparative examination of the urine. It was of a light amber colour, without any other sediment than the usual mucous cloud. Its specific gravity 1.0168. (See Exp. iv. p. 120.) Four ounces afforded of phosphoric acid, 2.97 grains. The quantity of urea and uric acid was also lessened, and apparently in the same relative proportions as the phosphoric acid. The bowels were regular, and the fæces bore a healthy character.

I directed him daily exercise, with a regimen of careful moderation, and that he should strictly abstain from malt or spirituous liquors. He was permitted two or three glasses of sherry daily. The limbs were to be treated with salt water in the usual manner. Costiveness to be carefully obviated by the pill, p. 282; and upon any evidence of vitiated secretion from the bowels or kidney, five grains of compound calomel pill to be taken at bed time.

In this very urgent case, the efforts of Nature in producing a fit of gout had not apparently relieved the system. At one juncture, apoplexy was strongly threatened; and for a considerable time, the evident congestion in the lungs was endangering a rupture of the vessels.

The tenderness at the hypochondriac and epigastric regions; the spasm of the diaphragm on stooping; the continual fulness of the hæmorrhoidal vessels; the dark alimentary secretions; were strong marks of obstructed liver, and of congestion in the circulation of the system of the vena portarum. From these united causes, the head was much affected, and the nervous system was seriously disturbed.

The comparison in the state of the secretions under the disease and at the return of health, and of the remarkable difference in the saline and animal principles of the urine, is worthy of attentive observation.

I saw this patient again, *Feb.* 1816. He reported that he had passed through the summer in perfect health, and entirely free from cough. He became tired after a few months of restraint of diet; and returned to the free indulgence of porter, and probably other liquor. At the return of winter, some cough again occurred. He had been free from all threatening of gout. His erroneous habits were producing plethora. He complained of cough, occasional heat of stomach, and attendant headache. Doubtless without entire reformation of habits, his former symptoms will all return; and that establishment of health, which care and prudence might ensure, will be effectually prevented.

CASE III.

A lady, aged 57, plethoric, but of the nervous temperament, has been subject to gout for eleven years past. About twelve years ago, she suffered continual uneasiness in the left side, which sometimes amounted "to a peculiar burning as if an abscess were forming." From the increased bulk of the part which she describes to have taken place in a short time, and the copious dejection of dark blood from the bowels, which occurred occasionally, my presumption is that the spleen was in a state of congestion and chronic inflammation. The stomach suffered by sympathy, but in addition to ordinary sickness, she sometimes vomited dark blood. She says, "that she received delightful relief from being cupped freely on the side." She has for many years past had hæmorrhoids, attended with frequent and considerable hæmorrhage. Has been much subject to pain and confusion of the head, with excessive sense of weight. Often dyspeptic, and subject to depression of spirits. Has suffered gout in almost every part; and many of the attacks have been induced by very slight causes.

At my first visit, I found this patient under a severe attack of gout affecting both feet and both hands. It had been induced by accidental exposure on a foggy day in November, with the wind in the east. The affected parts were vividly red and much swollen. She complained of violent throbbing, and severe cutting and pricking sensations, and frequent cramps. Her looks were expressive of great anxiety and suffering. The complexion was jaundiced. The left side was tender to pressure, and there was evident visceral fulness. A troublesome cough was present, and the respiration was

uneasy. Appetite was lost, and the stomach was much affected with flatus and with occasional sickness. The matter rejected, she described as either yellow or green; and that as one or the other prevailed, it was more sensibly bitter or acid. It was excessively slimy. I saw a portion which had so much the appearance of pus, that I was curious to give it attentive examination; and found it to be very concentrated mucus*. The urine most

* I used the ingenious method recommended by Dr. Young (Introduction to Medical Literature, p. 546), for the distinction of pus and mucus, of viewing the suspected fluid through pieces of plate glass before a candle; concluding it to be pus if a circle of coloured rays should appear, and vice versa. In this instance I found the test satisfactory, for no coloured light was produced; but I must observe, that our inferences must be carefully drawn, when we do find that the secretion from any surface, internal or external, exhibits the coloured rays. It is familiarly known, that a mucous membrane under disease, produces puriform secretion, when there is no breach of surface. We see this in gonorrhœa, and also in the expectorated secretion which attends many kinds of cough either recent or chronic; and which, in its appearance, often alarms both the patient and the practitioner. I have repeatedly found the expectoration attendant upon an alarming cough so entirely puriform in every character, that the lungs had been considered in a dangerous condition of disease; but perfect recovery took place. Hence we are taught to derive our judgment from the associated constitutional symptoms, and not from the threatening characters of what is expectorated. It is not generally known, how readily a mucous membrane under irritation takes on the morbid action of which I am treating. I have examined the secretion of the conjunctive membrane of the eye, when under very slight inflammatory irritation; of the Schneiderian membrane under catarrh; and of the urethra when affected by gleet; and have discovered by the optical test,

commonly deposited pink sediment, and the state of the bowels was such as also declared a morbid action of the liver.

In addition to the particulars already related, she mentioned that in the winter season she usually felt "as if she had rheumatic pains all over; and that the cramps and other uneasy feelings of her stomach and side were most troublesome when the limbs were the most comfortable; and the reverse, when gout was in the limbs." I learnt that to a degree of extraordinary idiosyncrasy, she would be affected to salivation with a small portion of mercurial medicine, if repeated thrice or even twice *at short intervals*. She was equally sensible to the power of opium on her nerves, and declared that she would rather endure any pain than take composing medicine.

I shall avoid the details of treatment in this case, which would be tedious; and confine myself to a general account. From the draught, p. 186, in regular use, an occasional pill of hydrarg. submur. et pulv. antimon., together with the free use of the lotion, the inflammatory symptoms were soon and most favourably removed. Pain and spasm were also controuled by pills of the stramonium extract and lactucarium, which did not prove in the least degree inconvenient to the nervous system.

a slight degree of the coloured rays. The circle of rays appears more distinct, and the colours more vivid, in proportion as the substance is of the true nature of pus; or, to give the philosophical explanation, accordingly as it consists of globular particles, by which the light is refracted. A chemical test for the distinction of pus and mucus, is yet a desideratum. At present, we do not possess any which is useful, and until such a discovery is made, the optical method claims our preference.

The side was relieved by blistering; and unless this had succeeded, the aid of cupping would have been added. After a period, with a view to strengthen the stomach, the draught, p. 266, in effervescence with lemon juice, and the pills there mentioned, were prescribed; but so repugnant was the system to tonics, that it could not be taken without some consequent heat and irritation more than once a day; and it was therefore continued only at noon; the pills or the colchicum draught being taken at night, and the mercurial pill about every fifth night. If repeated with more frequency, I had the proof that mercurial fever and irritation would ensue. The weakened limbs, when no longer affected with inflammatory tenderness, were treated with the morning sponging, diligent friction, and rollers, and a liniment occasionally, with the best effects. Ass's milk formed a part of the regimen. I have the pleasure to add, that, at the end of two months, the patient was very favourably convalescent from all the symptoms; and acknowledged a state of radical improvement, and of acquired spirits, strength, and comfortable feelings, which she had not permitted herself to expect.

In the following cases I shall endeavour strictly to confine myself to the most interesting details:

CASE IV.

J. P. aged 44; corpulent and robust; plethoric; of the mixed temperament; first affected with *acquired* gout at 36; has suffered most severe fits, never less than once, and sometimes twice in a year; was subject also to painful attacks of gravel until he discontinued the use of ale; laboured under chronic gout, and gave me the

following account of his case when he consulted me in March. In the preceding October, received a fall from a horse, by which his right arm was so much strained and bruised, that it became immediately discoloured throughout, and extremely tender. The gout had been hanging about him for a week. He was bled; took purgative medicines; and twelve leeches were applied to the arm. In two days after the accident, gout attacked the right foot very severely; and, in succession, the left foot, the right and left knee; lasting three weeks. In the injured arm no gout took place; and the painful sensations which had before troubled him in this part, now ceased to engage his attention, so severe was the suffering from the gout in the lower limbs. Through November and December he continued very lame; his feet and ankles swelling, and being very painful after slight exertion; often losing his sleep at night from actual pain of some part affected with sudden gouty inflammation; or from distressing starting of the limbs, or agonizing cramps. In January he obtained a temporary amendment by the use of Reynolds' specific; but he relapsed in February; and, in addition to the gouty state of the feet, he was tormented by rheumatic pains in the loins and arms; was occasionally affected with a spasm of the diaphragm, which he described as the cramp of the stomach; he was hypochondriacal in a high degree. I first saw him under these circumstances, and found the tongue furred; the appetite capricious; the bowels for the most part confined; the fæces clay coloured; the urine scanty and depositing pink sediment. I directed the feet to be sponged very thoroughly four or five times a day with the lotion, till inflammatory tenderness was removed; small doses of hydr. submur. pulv. jacob. et extr. colocynth. comp. each other night; the saline form of the

draught, p. 186, twice a day ; pills of pulv. ipec. compos. with lactucarium twice or thrice a day as pain might prevail ; and a moderately supporting diet. From the moment that this treatment was commenced, his amendment was rapid. The active alterative just mentioned, was, after a short time, exchanged for the pilula. hydrarg. ; the draught was administered in the first of the morning only, and followed by some ass's milk ; salt water sponging, and friction in the morning, and the liniment, p. 271, at night, were substituted for the lotion. In one month, the patient was, in every respect, perfectly and satisfactorily recovered ; no occasion having arisen for the use of tonic medicines.

CASE. V.

V. R. aged 51 ; slight ; of the mixed temperament ; highly irritable in constitution ; first affected with *hereditary* gout at 26, and has suffered of late years a severe paroxysm both in spring and autumn. The pain so continual and violent, that he has been deprived of sleep for eleven days and nights. The premonitory symptoms, cramps in the calves of the legs ; great irritability in the general state of the constitution ; restlessness ; head-ache ; furred tongue ; loss of appetite ; confined state of bowels ; and deficient secretion of urine. When this gentleman consulted me, he was affected with gout in the hand and elbow, and the chief attendant symptoms were such as I have now described. He informed me, that, in the course of the preceding twelve months, he had been six times very painfully attacked with gout, which he had unsuccessfully treated with Wilson's tincture ; and, latterly, thought his stomach and

nervous system much injured by that medicine. I never witnessed a more irritable habit. Palpitation of the heart affected him on slight occasions during the state of gouty diathesis. He remarked, that sometimes the skin has been so morbidly sensible, that upon touching any very cold substance, little bladders have immediately arisen. From agitation or slight bodily exertion, he would have faint perspirations. The tongue was furred and very white. The urine deposited a sediment consisting of the phosphates. The fæces were knotty and clay coloured. Antimony, in the smallest doses, had been found always to disagree; and few purgative medicines had suited his stomach. He was delighted with the excellent effect of the saline form of the draught, p. 186; and a pill containing hydr. submur. half a grain, and pilul. hydr. gr. iv. given each other night, agreed perfectly. For the relief of pain and restlessness, he took with success lactucarium, together with a dose not exceeding two grains of pulv. ipec. comp. which was occasionally repeated in two or three hours after being in bed; but more commonly a single dose proved sufficiently sedative. A poultice, with the lotion, and its use also by means of sponging, proved very successful. In this case, I preferred such modes of applying it, to the plan by linen compresses. Ass's milk was an useful auxiliary, causing the draught to act more pleasantly both on the bowels and kidneys. When the inflammatory symptoms were entirely overcome, I resorted to such treatment as was most calculated to improve the strength of the constitution, and restore the weakened limbs. The lower extremities were quite disabled from debility, induced by the previous neglect of all proper management. Friction for an hour daily, with the other means so often mentioned, produced the best effects. The only form of tonic medicine which could be

taken without stimulating and heating effects, was the infusion of columbo, with the carbonate of soda, and the addition of lemon juice, at the time of taking it, to constitute a saline draught in effervescence. To conclude, this gentleman recovered and retained a state of constitutional health and spirits, and strength and ease of limbs, to which he had, for the previous two years, been a total stranger.

CASE VI.

The subject of the following interesting case, was a medical gentleman, to whom I had the satisfaction of advising a successful mode of treatment. I shall offer an abridged statement of his own observations.

J. W. aged 45; short; of middling bulk; of the nervous temperament; first affected with *acquired* gout at 21, in the instep, and which was treated by mistake as a sprain. Till within the last two years has been attacked at short intervals; and sometimes one severe fit has followed another, with the respite only of a month. The hands, feet, knees, elbows, and hip joints have all been affected. The usual first notice of an attack was a pain on leaving the bed in the morning, which quitted him as the day advanced, and by means of taking his usual exercise; but returned about seven or eight in the evening, and continued till nearly the same hour in the morning. Inflammation immediately succeeded. To describe, he says, the violence of the pain, is impossible. It was attended with a sense of weight, wrenching, gnawing, burning, and as if some one were beating the parts with a large hammer; and, in short, no torture can be imagined more severe than that which I suffered. The pulse was quick, the tongue furred, the bowels torpid, the kidneys

at first active, with a copious discharge of clear urine for the first day or two; then scanty, high coloured, and depositing abundantly the pink sediment; the spirits much depressed, and with so much nervous restlessness, that I could not bear the approach of any one near me. With the continuance of the fit, the stomach became extremely irritable, and often affected with violent retchings. Cramps were severe, particularly at night, together with watchfulness and delirium. The duration of the fit was usually three weeks without any intermission of suffering, either day or night. As the attack sometimes commenced without notice, so it quitted as suddenly; leaving much general exhaustion, and great stiffness and tenderness of the affected joints. As soon as the pain subsided, the state of the secretions improved, but did not become healthy. The appetite was voracious, and much sleepiness prevailed. It commonly happened that a relapse followed in a fortnight, and which sometimes gave rise to symptoms equally, or, if possible, more severe than in the previous attack; but their duration was shorter, and the convalescence more rapid." In regard to the treatment, he had made trial of calomel to the extent of salivation, and ordinary purgative medicines; with the local use of leeches, fomentations and embrocations, without relief to his sufferings. From Wilson's tincture he procured present ease; but the gout so quickly relapsed, that he was dissatisfied with its effects; and two years ago, under a protracted state of sufferings, he resorted to the combined method of treatment which I advised. In reporting to me lately the effects which it produced, he says, "I am happy to inform you, that the plan succeeded beyond my most sanguine expectations; and, from that period to the present time, I have never lost two nights' sleep from the gout." He men-

tions that he has made a comparative trial of the draught as described at p. 186, with and without the acet. colch.; but that the omission of this ingredient most materially lessened the good effects of the medicine.

CASE VII.

C. L. aged 53; slight; of the nervous temperament; first affected with *acquired* gout in the ball of the great toe at 50; and since in both feet on three occasions. I found him suffering from a chronic state of gout in the feet; and alarmingly ill with an urgent cough, copious puriform expectoration, a quick pulse, some degree of hectic fever, and a countenance which bore a consumptive stamp. On the day of my first visit, some urgent business had called him from home, and he was accidentally exposed to wet. He was subject to a severe cramp at the pit of the stomach from slight causes; and on the night following to this exposure, he was seized with this distressing symptom about two in the morning. It continued a considerable time. He described the sensations which followed the first sudden seizure of pain, as the feeling of weight and coldness like that which might be produced by a piece of freezing ice; or at another time, when the weight was more oppressive, as if cold lead were forced into the stomach; and the aching was severe. Repeated draughts of warm brandy and water afforded relief; but, in case of its return, I directed a mixture of camphor, æther, and opium. On the succeeding night he was attacked in a similar manner. The first dose of the mixture contained fifty drops of the tincture of opium; and two succeeding doses, which were taken with intervals of half an hour, fifteen. This quantity completely re-

moved the pain. It returned slightly on the next night, but yielded to a dose of the mixture containing thirty drops of the tincture, and never after returned. The cough and attendant constitutional disturbance now engaged my attention. The alvine secretions were such as to mark a very unhealthy condition of the liver. The saline form of the draught, p. 186, pills of pulv. ipec. comp. et lactucar. and a mild dose of calomel each other night, answered my utmost wishes. For ten days I avoided any treatment to the feet, which continued to be affected with chronic gout, and which had not alternated with the condition of the stomach already described. I feared that any application, however cautiously made, might incur the imputation of causing some internal inconvenience. At this period, however, the ankles being œdematous, tender and aching on the least exertion, I employed, in succession, the lotion by means of frequent sponging, the tepid salt water, and the stimulating liniment. The best effects resulted, and not the least disadvantage to the constitution arose. This gentleman completely lost all his symptoms, with the exception of a slight degree of habitual asthma, in the space of six weeks. I never witnessed such threatening symptoms of phthisis in connexion with gout, as in this instance, yielding so perfectly to treatment. An interval of a year has now elapsed, and the patient has remained well.

CASE VIII.

L. M. aged 50; corpulent; plethoric; of the mixed temperament; first affected with *acquired* gout four years ago, at the ball of the great toe. I found him suffering from chronic gout, and that the symptoms had continued

in each foot, alternately or together for two months. He had refrained from any treatment, from an idea that the gout was a disease which would cure itself; and from some apprehension connected with the recollection of an irregular action of the heart, which occurred a few years before; an account of which I shall state in his own words. "So well as I can remember, the irregular action of my heart, and consequently irregular pulse, came on in the year 1811. I cannot say that it was really palpitation, nor was it in the least painful; but from a peculiar sensation which I felt in my left breast (always there and no where else), I could tell every beat, or rather the want of beat in my pulse, as plainly as possible. The sensation was like that produced by a small electric spark. This symptom, without any variation in the sensations, continued till September 1814; when, for the first time, I had a week's fit of the gout. I had tried various medicines for the relief of the heart, without the least success. I took particular notice, that on the very evening when my gout first came on, the feeling which I had in my left breast (and from which I could plainly tell the missings of my pulse without examining it) entirely left me; and my pulse became quite regular, and has continued so to the present hour, September 1818." This gentleman attributed his singular indisposition to the suspension of his former active exercise on foot and on horseback; so that he became more corpulent, and full in habit. In regard to his chronic state of gout, I found that the symptoms had been supported by congestion in the abdominal viscera, and an unhealthy condition of the secretions. The aperient and alterative treatment, with regulated regimen, perfectly succeeded; and the limbs were restored by the usual plan of proceeding, which I have detailed in other cases.

CASE IX.

About a year ago, I was consulted by a gentleman, aged 46, who, together with occasional painful fits of gout, the first of which occurred at the age of 40, was subject to very painful attacks of head-ach; and for four years past had been sensible of an irregularity of pulse, which prevailed, more or less, constantly. The symptoms premonitory of gout, were, cramps in the legs, acidity of the stomach, confinement of the bowels, a scanty secretion of urine, and a deposition of pink sediment. The stomach was exquisitely sensible to the influence of acid, so that he never could eat even a ripe orange with impunity. The feet were habitually very cold, unless when taking strong exercise. He mentioned, in regard to the head-ach, that on the day before, he felt a depression of spirits, and a desire to take more wine than usual; he being in his general habits very temperate. On the morning of being affected with the head-ach, he could not bear the use of cold water; although at all other times he derived much comfort from free ablution with it. Together with pain he had many nervous sensations in the head, and a troublesome pulsation. I have stated these details, having the satisfaction to add, that the course of alterative aperient treatment, and regulated regimen, which proved of the utmost service to him in removing the disposition to gout, was equally useful in counteracting the attacks of head-ach. The necessity for this method of treatment was manifested by the faulty state of the alvine secretions, and by the frequent deposition of pink sediment in the urine. Observation alone could lead to these conclusions; for the appetite was regular, and the bowels seldom failed to act daily. In

this example we see a proof afforded, how much the cause of gout is generated below the stomach; how insidiously the disorder forms by slow degrees, when the patient, possessing appetite and a seemingly healthy digestion, and regular action of the bowels, thinks that all the functions go on well. The truth is, that venous congestion is produced in the vessels of the respective viscera of the abdomen, which part of the body increases in bulk; and at length sensations of oppression are felt. Nature, in her attempt to cure this state of evil, excites the secreting vessels of the liver and other parts, the kidneys especially, to increased action; and the excreted fluids are more stimulating in their nature. Although, therefore, disorder arises in the alimentary canal, either spontaneously or excited by medicine, the *vis medicatrix naturæ* is at work; and I should apply the theory which I have offered at p. 139, in regard to the urine, in explanation of the condition of bowels which takes place. It is the object of medical interference to remove this obstruction of the vessels, assisting Nature's intention; and, afterwards, to bring about a more healthy œconomy by means of corrective medicine and regimen, concluding, when necessary, with the judicious use of tonics. It too often happens, however, that medicines of the latter class are given in the first instance; and no proper foundation having been laid for a superstructure, the visceral obstruction becomes only fixed and confirmed.

In the present case, the patient derived material advantage from the daily plan of sponging the feet with tepid salt water; and washing the head freely by means of a coarse towel dipped in cold water; using afterwards sufficient friction to dry the skin.

CASE X.

In the following case, the prevalence of the nervous temperament, in concurrence with the gonty diathesis, produced a combination of symptoms highly distressing and remarkable.

A gentleman, aged 40, rather plethoric and inclining to corpulency, was first affected with hereditary gout at 32, in the great toe of one foot only, severely, for ten days; since which the attacks have not been acute, nor attended with much pain. He described that three years ago he became ill with the following symptoms; and of which he was complaining when he consulted me. The gout, he said, seemed to be hanging about him. He experienced great languor; occasional palpitation of the heart; uneasiness and confusion of the head; exceeding depression of spirits; broken rest, and uneasy dreams; a sense of weariness in the arms, so that it was a painful exertion to hold them up; a dull aching pain between the shoulders; a remarkable coldness of the lower extremities, and general cramps. On two occasions blisters had been applied to the legs with a view to excite gout in the feet; and on one occasion success was procured in this object, but he did not obtain the desired relief to the constitution. A year ago, after exposure to wet and cold, he was alarmingly affected with torpor in the limb, and a threatening of general paralysis. The stomach was affected with sensations of spasm, and some sickness. The symptoms increased at night; quitted for two or three days; and then returned with the utmost aggravation of all the general nervous symptoms. I found abundant evidence of an obstructed state of the liver, and an unhealthy state of the alimentary canal. The appetite itself was seldom deficient.

The feces were sometimes black in appearance ; at others, clay-like. The pink sediment almost constantly appeared in the urine. I instituted a regular course of alterative treatment, by corrective and aperient medicines, in conjunction with ass's milk and regulated regimen ; allowing him daily three or four glasses of Madeira. His tranquillity and rest at night were much assisted by the use of lactucarium. I directed that the limbs should be sponged every morning with tepid salt water, and freely rubbed at night with the liniment, p. 271. The head was treated with the ablution before mentioned. The happiest results succeeded to these measures steadily pursued for two months. Previously, he had taken purgative medicine only occasionally ; and had not been put upon the necessary course of remedy to eradicate the evil. Finally, I prescribed a draught of *sp. ammon. compos. decoct. al. comp. infus. cascarill. mist. camph.* to be taken, at first twice, and afterwards once a day ; and with such good effect, that the disease, at length, yielded quite to my satisfaction.

CASE XI.

The case I have now to describe, was one of chronic gout, very obscurely marked in its origin and progress ; but which became remarkably tractable under a change of treatment.

C. R. aged 54, tall, rather corpulent in the abdomen, plethoric, of the mixed temperament, first at 43, complained of some swelling of one ankle with stiffness and weakness ; which occurred, repeatedly, with the interval of a few months. Gout was unknown in his family. In 1812, when fishing, he strained one knee. On the fol-

lowing morning he complained of stiffness, which he expected would be carried off by exercise, and he walked accordingly, but the uneasiness increased; and on the third day active inflammation took place, with swelling and pain. This was not considered to be gout. It was treated with poppy fomentation. Much weakness was left in the part, for which camphorated spirit was applied without advantage. A stimulating lotion was next used, and also a tight bandage. This treatment produced inflammation and pain. The Buxton bath was tried for one month without benefit. A course of friction was next resorted to for seven weeks; at first for twenty minutes, and latterly for an hour, twice a day. The knee was restored by this plan; but at the end of the seven weeks, the other knee and both ankles became suddenly affected with pain, and with real symptoms of gout. There was not much redness, but the throbbing was considerable. The symptoms continued three weeks. In the succeeding four years, he was troubled with several attacks of short duration. In February 1817, the knee originally affected was again attacked with swelling, stiffness, and much pain. He took a purgative medicine. On the following night the other knee became affected. Took half a bottle of Wilson's tincture. Procured sleep, and next morning both knees were so much improved, that he could walk down stairs without much trouble; and in three days after he went into the country. A relapse took place almost immediately, the ankles becoming chiefly affected. Took the same dose as before, of the tincture. No apparent effect from it until twenty-nine hours, when it acted upon the bowels violently for two hours; no sickness. It gave relief. In ten days suffered another relapse, and he repeated the tincture, but without any relief. It caused sickness without affecting the

bowels. After this, he took the *can medicinale* and Wilson's tincture alternately, till the month of June; with no other distinguishable difference in the effect, than that he found a larger dose of the tincture necessary to produce an equivalent influence over the symptoms. The patient continued to have relapses, parrying off severity of pain by one or other of these medicines, in frequent small doses; but he was rendered very sensibly weak and languid, and altogether nervous by the treatment; and in the hot weather of June, a more decided attack coming on, he determined to discard the empirical remedies, which he felt to be injurious to his constitution, and at best only an uncertain palliation. He was then put on the use of a draught composed of sulphate of magnesia, peppermint water, tincture of hop, and some cloves. His stomach became improved, and he thought himself recovering; but in the beginning of July, he had a slight attack in one knee. He now discontinued the use of wine and all fermented liquors, and lived in the most simple manner. At the end of the month, one knee and the whole of both feet became more affected with decided gout than at any former period, and he suffered severely. This attack left him very lame for a long time. When I first saw this gentleman, in the middle of September, he was using crutches. The bursæ at the knees and ankles were much distended and tender; and in his constitution he was much enervated. Upon investigating the case, I found very strong evidence of unhealthy secretions; the abdomen was out of proportion large; the muscles of the lower extremities were small and relaxed. I prescribed my usual aperient and alterative treatment; ass's milk and regulated regimen; and, to relieve the languor of the system, I advised two or three glasses of wine daily after dinner. *Lactucarium* was

taken occasionally at night, with an useful effect in producing tranquillity and sleep. The affected parts were at first repeatedly sponged every day with the lotion, p. 254, and when inflammatory tenderness was quite removed, the united means of salt water sponging, the liniment, p. 271, moderate bandage and friction, were employed with unremitting attention. Each morning, ablution with cold water was used to the head with much comfort and advantage. The crutches were quickly laid aside; and not only have never been resumed up to the present period, February 1819, but during the whole period he has scarcely had the least return of gout, and has never been confined to the house a single day. On feeling any threatening of indisposition, he resorts to the draught, p. 186, and occasionally takes an alterative pill. Formerly he suffered at times from gravel, and often passed sharp crystallized portions. Then he took malt liquor without particular restriction. Since the present plan of treatment, he has not once been indisposed with this complaint. No disadvantage has arisen from the daily indulgence of three or four glasses of wine; but, on the contrary, the constitution has appeared benefited by it, quite in an equal degree to the comfort which it has afforded him.

CASE XII.

A gentleman, aged 59; corpulent and robust until reduced by gout; first attacked with acquired gout at 43, having induced the disease by freedom of living, suffered severe and frequent paroxysms, which for many years he left chiefly to their natural course; but for a few years past had taken the eau medicinale and Reynolds'

specific, procuring from each medicine a palliation of symptoms ; yet followed by so much frequency of relapse, that at length he gave up this treatment. From the can medicinale more particularly, his limbs became remarkably weakened ; and his head disordered by nervous sensations in an alarming degree. He next prepared colchicum in rum and water by infusion, and took small doses daily for a long time ; deriving occasional ease, but having an attack of gout either in the upper or lower limbs every two or three weeks. At my first visit I found him labouring under a distressing complication of gout, rheumatism, and gravel. From the last complaint he suffered to such a degree, that the urine was often bloody. The change of organization in the limbs was remarkable. The fingers were for the most part knotty and contracted. The left knee was enlarged with bursal distension, and at that time affected with gouty inflammation ; the ligaments were in a tender state ; the right knee was almost in a state of perfect ankylosis ; the bursæ were distended ; the ham-strings contracted, and as hard as wires. The foot of the same limb was almost ankylosed, and its form distorted ; the other foot nearly in a similar condition, with the integuments œdematous, and the skin of the leg partly in the state of ichthyosis. The ligaments had evidently undergone such thickening and shortening, as to forbid much expectation of useful treatment. The internal functions were highly disordered. The appetite was deficient ; flatus of the stomach and bowels was infinitely distressing ; the bowels were quite torpid, and the discharges were either black or clay-coloured. I could scarcely enumerate the train of urgent symptoms which afflicted this patient ; but I must add, that rheumatic pains affected the arms, the muscles of the ribs, and the loins.

By day, he was wholly confined to the chair; at night, he scarcely procured sleep. When, by means of the alterative and aperient treatment, the use of the black drop in small doses twice or thrice in the twenty-four hours, the lotion and poultice externally, I had succeeded in removing the gouty inflammation; I adopted the plan of friction twice a day for an hour, salt water sponging, a liniment, and the application of rollers to the ankles, in conjunction with the continuance of the medicines just mentioned, and the addition of a tonic. Ass's milk, and a strictly regulated regimen, were added to the medicinal agents. At the expiration of three months, the tendency to gouty relapse seemed obliterated; the rheumatism was cured; and even in a short time from the commencement of the treatment, the complaint of gravel was overcome. The improvement of the limbs exceeded my utmost expectation. He acquired the power to walk with a pair of sticks, a cork-heel being added to the shoe, that he might be enabled to bring the foot of the shortened limb in contact with the ground. His health and spirits were delightfully improved; and he removed into the country for the benefit of change of air; with every prospect of gaining as complete a recovery as the circumstances of a case, previously so much neglected and ill-treated, would possibly allow.

In a case of a similar nature to this which I have related, the gentleman having been confined wholly to his chair for several months, in consequence of the stiffness of the knees and ankle joints (almost amounting to complete ankylosis), occasioned by rigidity of tendons, thickening and shortening of ligaments, and distension of the bursæ, with frequent occurrence of chronic gout; I had the satisfaction, by perseverance with the plan of friction and other external treatment, in conjunction with

the use of internal medicines, to effect such a change that the patient was enabled to walk with the help of short sticks, and in fact to undertake a long tour upon the continent. In this instance the use of friction was for a considerable time interrupted by the occurrence of a small vesicular eruption, spreading generally on the legs, and which had been produced, before I saw the patient, by the irritation of the skin, occasioned first by the bites of leeches, and afterwards by a blister; these remedies having been used with the hope of subduing the chronic inflammatory action; but the above mischief took place, and not the smallest benefit was afforded by this treatment.

CASE XIV.

A gentleman, aged 46, slight in form, and of the nervous temperament, was first affected with *acquired* gout at 26, in the ball of the great toe. Subsequently, with the exception of one paroxysm, his attacks have been of a very anomalous character. In September, 1816, after exposure to wet and cold, he was seized with general symptoms of fever, painful irritation of the bladder, and a mucous discharge from the urethra, with much ardor urinæ; no cause having been applied to produce such a form of complaint. In a few days, a violent inflammation of the left knee took place, with excruciating pain; a numbness and soreness of the right hip joint; a feeling of coldness at the pit of the stomach, with sickness so as to reject every thing almost as soon as taken; and a torpid state of the bowels. Soon afterwards, the shoulders, elbows, wrists, thumbs, and some of the vertebræ, were also attacked; and, very shortly, the ankle, metatarsal bones, and great toe of one foot. The pain of some of

the parts, particularly of the affected knee, was so violent that he could not move in the smallest degree without assistance, and it was necessary to change his situation from one part of the bed to the other by means of a folded sheet. During twelve months, a series of remedies, too numerous to be mentioned, were employed without success. Mercurial salivation was produced on two or three occasions, but always with the effect of causing high constitutional irritation, and an aggravation of local sufferings. Anodines gave imperfect relief and disordered the head. Balsam of copaiva, the decoction of sarsaparilla, liquor potassæ, the alterative use of the blue pill, and other medicines, were tried in succession. At one period the nitro-muriatic acid bath was used, with no other effect than that of producing troublesome pimples on the skin. The hot bath, at a high temperature, was employed with diligent perseverance. He was put on a milk and vegetable diet. To the diseased knee, acute inflammatory action having subsided, a liniment of tartarised antimony was applied with friction, which occasioned excessive inflammation, swelling, sloughing, and a foul ichorous discharge from the ulcerated parts, accompanied with urticaria over the whole body, producing intolerable irritation and itching, and much sympathetic fever. Finally, when the parts were healed, the enlargement of the knee was reduced, and the pain mitigated, but the urticaria continued very distressing; and the general health was not at all improved by the influence of the medicines which were joined with the local treatment. I visited this gentleman first in December, and found him entirely disabled in his limbs, so that he had not the power to move about the room even with the aid of crutches. At the hips, knees, and ankles there was swelling and much tenderness. The muscles were exceedingly relaxed; the

countenance wore every mark of languor and debility. From the free use which had been made of the hot bath, and an excessive quantity of flannel clothing, the skin was in a state of unhealthy perspiration. The secretions were much vitiated; the urine depositing a copious sediment, which consisted, at different times, either of the phosphates or urates, or the two deposits mixed; and the alvine discharges were either deficient in bile, or contained that secretion of vitiated quality, as was denoted by a greenish colour. The tongue was coated with a dirty white fur. The nervous system was in a state of high irritation. He was much affected with cramps; the pulse was quick; the appetite deficient; the spirits depressed. I adopted the following treatment. The lotion p. 254, was used lukewarm, by means of a sponge, to all the affected parts, frequently in the day; and was continued so long as inflammatory tenderness remained; after which, salt water sponging was applied night and morning; and regular friction, by a skilful rubber, was employed, at first for an hour in the morning, and afterwards morning and evening. A chamber horse was used as a mode of exercise. In the two preceding cases also I recommended this simple expedient as an auxiliary to the general plan. It affords considerable exertion to the muscles, and particularly to those of the abdomen; and serves therefore to quicken circulation and action where most required. The internal remedies consisted in the alterative use of a mild preparation of mercury, so cautiously administered as not to cause any feverish excitement; and in the use of the draught, p. 186, for a considerable period twice, and afterwards once a day; ass's milk being taken in the morning early. The plan of diet was changed, and gradually made quite of a restorative kind. A little wine of the best quality constituted a part of this

regimen. The patient improved slowly, but without interruption; and when the month of May had arrived, he could accomplish walking exercise to the extent of eight or ten miles in the day, without the least inconvenience. The puriform discharge of the urethra had been in a great measure dependant on some local inflammation; but, in spite of every treatment, it continued, more or less, until the constitutional health was restored. This case, therefore, exhibits in the most favourable light, the advantages of the alterative and aperient treatment steadily followed up, the tonic part of the plan being made to consist in good diet. When the patient made trial of tonic medicines, under my direction, he did not derive from them the least advantage; but found either their stimulating or astringent influence to be unfavourable. Friction and the other treatment for the limbs produced the best effects. By recent intelligence from this gentleman, I learn that he has continued perfectly well.

CASE XV.

The following is a more unequivocal instance of a puriform discharge from the urethra being entirely connected with a gouty state of the constitution; and continuing until the general health became restored.

A gentleman, aged 45, robust, inclining to corpulency, plethoric, of the mixed temperament, was first affected with *hereditary* gout at 23, in the great toe only, but since in both feet, the knees, hands, wrists, and elbows. The paroxysm has sometimes been so severe for two months, that he has required to be fed the greater part of the time. Has taken about thirty bottles of the eau medicinale, with the usual palliative relief in the early use

of the medicine ; but latterly it began to fail in giving present ease ; and his stomach became so weak, as to render him extremely dyspeptic ; and, to use his own words, “ he was often gouty in the stomach.” The head became so alarmingly affected with giddiness, that at length he abandoned the medicine with dread. Has taken *vinum colchici* with less injurious effects, but without the least permanent benefit ; relapses of gout occurring very frequently. I was consulted by this patient first in June. He was then suffering from chronic gout, and was troubled with a puriform discharge from the urethra, and ardor urinæ in a great degree. He states, that these symptoms had continued from the month of February, with the exception of a fortnight, when acute gout affected the feet, and completely suspended them ; but they returned with all their previous virulence at the cessation of the paroxysm. The alterative and aperient plan of treatment, with the use of ass’s milk, and regulated regimen, proved completely successful. At bed time he took pills of *lactucarium* and *pulv. ipec. comp.* with much advantage. At the end of two months, every feeling of gout was removed. After a few days use of the medicines, the symptoms affecting the urethra very much abated, and had ceased at the end of a week ; but then they recurred again for a short time ; finally, however, yielding completely to the constitutional means of treatment.

CASE XVI.

The following case may be offered as the most common example of the lameness and chronic condition of gout, which are left by the acute paroxysm, when injudiciously treated or wholly neglected. It very favourably

exhibits the advantages of friction, and of constitutional treatment.

A. B. aged 40, of middle height and bulk; robust, of bilious diathesis, and of the sanguineo-nervous temperament; was first affected with *acquired* gout at the age of 27; and, since that period, has not had a longer interval from gout than nine months; the fit sometimes returning four or five times in the year. He has been attacked severely in the feet, knees, hands, wrists, and occasionally in the shoulders. In his last paroxysm, he suffered intense pain in the groins. This gentleman possesses a strong constitution. Till within the last two years he took the eau medicinale freely; and usually succeeded in removing the immediate symptoms, but the fit returned with four-fold frequency, and without any abatement in severity; the stomach became weakened; the loins remarkably affected with aching pain, stiffness and debility; and the ankles swelling upon every slight exertion. At a subsequent period he made free use of Wilson's tincture, and found it a less active medicine, both as to benefit and injury, than the eau medicinale. It proved only a temporary palliative, his disorder increasing in the constitution in the same proportion that he attempted this mode of cure. I found that he had scarcely been free from gout during a year past; and was then so lame as to require the use of crutches. On first leaving his bed in the morning, he was almost unable to bear in the least degree on the feet, such was the weakness and tenderness of the ankles and knees. With some patients I have met with this state of infirmity in an extraordinary degree, although, when the day has advanced, they have walked tolerably. To proceed with my account, he complained of all the symptoms just now stated; was distressed with cramps, and at night with

starting of the limbs; had flying pains in the shoulders; aching at the right shoulder blade: the appetite was very good; but from any incaution in the quantity or quality of his meal, he suffered oppression at the stomach, and heart burn. The complexion was yellow. The urine deposited much pink sediment, and the alvine secretions were quite disordered, the fæces variously appearing mucous, very dark, or clay coloured. I shall conclude with briefly stating, that friction was used for an hour daily, its use being begun as soon as the tenderness of the parts was sufficiently removed; and such was the happy result of the combined plan of external and internal treatment, which was attentively pursued, that at the expiration of three weeks, this gentleman was enabled to walk without lameness; and, in six weeks, his general health was quite restored.

In the two following examples, the immediate connexion between an unhealthy state of the digestive organs, and the occurrence of gout, as cause and effect, is strikingly shewn.

CASE XVII.

C. D. aged 52, tall and of middling bulk, of the nervous temperament, was first attacked with gout a few months ago. Neither of his parents had this disorder, but on the maternal side he referred it to uncle and aunt, grandfather and grandmother. His ankle and the internal edge of the foot were the parts affected. The skin was red and shining. Intense pain came on about two in the morning and continued till five. With the first invasion of the symptoms, he was seized with cold per-

spirations, and fainting; to which he has for many years been occasionally subject. The acute symptoms were tedious. He had taken only some ordinary purgative occasionally. Much weakness was left in the foot; and it was œdematous after slight exertion. The constitution was more seriously affected. This patient was a clergyman, and described, that, since the gout, his nerves had been so weak that the exertion of preaching had the effect of rendering him extremely faint and exhausted. He often complained of a sense of tightness in the head, noises in the ears, and a frequent vision of black spots, or of flies.

He also felt, when in cold air, as if the wind were penetrating into the head. The appetite was far beyond the powers of digestion; as he constantly suffered from heart-burn and flatus. Frequent support to the stomach was necessary, without which he experienced a sense of sinking at the pit of the stomach, and general languor. The secretions were quite unhealthy. The tongue was mostly white, and occasionally furred; the saliva viscid, and often of a saltish taste. The urine deposited a mixed sediment; the bowels were torpid; and the fæces were often knotty, and either very dark, or of the clay-like appearance. Latterly he had taken some stimulant bitters, upon the theory that debility alone prevented the healthy functions of the alimentary canal. This treatment aggravated all the symptoms; and with it he experienced occasional sensations of gout in the great toe. It was evident that purgatives and alteratives, with regulated regimen, would constitute the only safe and useful plan of cure in this case; and which was immediately adopted. Each other night he took pills containing hydr. submur. gr. ss. pilul. hydr. extr. colocynth comp. extr. hyoseyami. āā gr. iv. and the draught, p. 186, once or

twice daily for a fortnight; and afterwards only in the morning early; and in the middle of the day, one composed of equal parts of decoct. al. compos. infus. cascarrill. mist. camph. with sp. ammon. compos. ʒss. the alterative pill being reduced simply to pilul. hydr. gr. iv. For the fourth week, the last mentioned draught was taken twice a day, and the former omitted entirely. He derived much comfort from the occasional use of lactucarium at night; and of a cordial mixture in the day, when feelings of languor returned; which, however, he seldom experienced. It was matter of no small surprise to him, that the early part of this treatment which he apprehended would prove of a lowering nature, seemed to improve his strength and lighten his spirits, when the previous plan of tonics had really oppressed him and increased his feelings of debility. Ass's milk, regulated regimen, the morning ablution of the head, and sponging of the feet, were means which entered into the general treatment of the case; and the termination of it was most successful.

CASE XVIII.

G. B. aged 48, tall and slight, of bilious diathesis, and sanguineo-nervous temperament, first affected with acquired gout at 38. He found himself becoming more corpulent when first visited by gout; and he thinks that some fulness of habit has been an admonishing circumstance in most of his subsequent attacks. He has invariably found himself highly bilious when affected by gout. His habits are remarkably active, and for a long time past he has been very temperate in diet. One of his attacks appeared to be consequent to the disuse of regular horse exercise. This gentleman was labouring

under chronic gout, chiefly affecting one foot, when I was first consulted. The integuments were œdematous; the great toe much enlarged, very tender, and occasionally painful. I adopted my usual plan of treatment, and with the usual success; but found more than ordinary occasion to follow up the use of moderate doses of alterative aperient pills, in conjunction with the draught p. 186. The discharges were usually of a greenish colour, often exhibiting an oily appearance on the surface, and peculiarly foetid; sometimes they were knotty. When describing, in a written report, this state of the bowels, he says, very emphatically, "My toe is to bile, what a barometer is to the weather. In the same proportion that the bowels were freed from their improper contents, and from the green bile especially, the tension and pain about my toe sensibly and indeed rapidly diminished. I feel convinced, that had I abstained from calomel another day, I should have had a return of gout; perhaps of a severe nature."

When all the symptoms of gout were removed, he resumed his regular horse exercise with very great advantage; and I must here observe that this important remedy should always be recommended in the general directions of the physician, when the convenience of the patient permits. Horse exercise, and the air of the country, are powerful aids towards a perfect recovery, when the stage of convalescence is advanced.

I have introduced these last two cases, for the purpose of shewing the immediate connexion of gouty action in the limbs with the vitiated state of the internal secretions; and this at a time when the appetite is good and regular, and the tongue is bearing a healthy appearance; so entirely is the fault of the digestive process, under the

present circumstances, below the stomach. Nay, it sometimes happens that the urine will not, for the most part, appear unhealthy either in quality or quantity; and the whole error consists in the obstructed state of the liver, in the vitiated secretion of that important organ, and in the secretions of the intestines themselves. We are ignorant of the proper functions of the spleen and of the pancreas; but it may be entertained as a probable opinion, that the healthy œconomy of these parts is also interrupted. In those instances of remarkable corpulence about the abdomen, with indications of fulness of the habit, which have been gradually taking place, before the gout has declared itself in a fit, I consider that the vessels, and the veins especially, of the whole contents of the abdominal cavity, have been in a state of *engorgement*; to use an expressive phrase. In most persons, and particularly in those of plethoric habit, we see a very distended state of the veins in the extremities, either just at the time, or rather before the invasion of the fit. How well then does the treatment by alteratives, and aperients, and regimen, unite with this pathological doctrine. The practitioner must not too hastily satisfy his judgment that he has rectified the secretions, or done all that is required toward the lasting improvement of the constitution. One day may afford the appearances of healthy secretion, while, on the following day, the very contrary evidence takes place. Also it must be his care that he reduces the size of the abdomen within proper limits; and he must instruct the patient to keep watch against its insidious increase. At the same time every care is to be employed to improve the bulk and firmness, and muscularity of the limbs, by suitable local treatment and exercise. Contrast the recovery of the gouty invalid, conducted upon these principles, with the results of

the empirical modes of treatment. Witness, from the latter, the state of the patient, with the stomach weakened, and the general secreting functions rendered torpid, the abdomen large, the muscles and the limbs small and weak, with the integuments œdematous; and, lastly, the nervous system, sooner or later, shattered to the foundation. A few constitutions of extraordinary vigour may afford exceptions to this strong picture; but I sincerely affirm, from my general observation, that the colouring is faithful, and not too deep in a single tint. I have still a few words to add.

In the cases which I have here offered, I may appear to have drawn too favourable a sketch of the treatment of chronic gout. I have not exaggerated a single fact; but at the same time, I do not wish to keep out of sight that in some cases the cure is with difficulty effected. Yet I must contend, that however tedious the recovery may in some instances prove, the principles of the practice do not on that account become invalidated. Patience must be exercised; for the alterative method is of necessity slow. Yet, though slow in progress, its effects are sure.

A protracted state of chronic gout is almost invariably maintained by an unhealthy condition of the functions of the liver; and when this is the fact, the patient should be rather taught that he is undergoing a probationary course for the cure of this organ, than for the cure of gout alone. With such a truth impressed upon his mind, he will the more readily accommodate his expectations to the periods of weeks or months, which may elapse in obtaining his cure. I am clearly convinced, that no treatment but that which is truly constitutional can procure solid and lasting benefit. If, also, in the progress of the cure, a relapse of gout do take place, it does not contradict the propriety and value of the plan of treatment; but rather proves the deep seat

of complaint which has been fixed in the constitution; and our conclusions should animate us to persevere in our means of gradually eradicating the evil. I have before remarked, that the most difficult cases of gout are those in which some quackery has been previously employed; and from which, a disposition to frequent relapse has been acquired; but I have also observed, that due perseverance in regular principles of proceeding, will conquer even this difficulty. I have dwelt, tediously I fear, on these points; but I have done so, because experience has taught me their importance.

At p. 342, I have adverted to the blended symptoms of gout and rheumatism which occasionally takes place. Cases of this description are usually met with in those who have injudiciously nursed themselves with flannel, a great quantity of cloathing in bed, and a very hot apartment, during the period of a paroxysm; neglecting also, in all other respects, the proper methods of treatment. The liability to relapse, upon every exposure to a humid atmosphere, or a cold wind, proves a distressing circumstance. The physician, when consulted in such a case, must give the patient an impressive warning on the subject of exposure; for otherwise his best means of practice will fail, and incur undeserved reproach. The following instance will exemplify my present statement.

J. R. aged 60, of slight form; of the nervous temperament; was first attacked with *hereditary* gout at 30. For three months in the autumn of 1817, he suffered a series of relapses of gouty pain and slight inflammation, affecting the feet and knees. Each attack confined him to the bed or the sofa. He used such an excess of flannel covering to the parts, that he produced constant perspiration, locally and generally; and infinitely increased the debility which the disorder itself occasions. For several nights he took a drachm of *vinum colchici* with

some magnesia. For the first two nights he experienced a little relief; but afterwards he states, it had not any useful effect. In the short intervals of imperfect recovery, he yielded to the calls of his affairs, and either changed his apartment or went abroad; and on one occasion, being accidentally exposed to wet, he brought on gout in one knee severely; and also was tormented with flying rheumatic pains in the arms, and in the sciatic nerve of each limb together or alternately. The gout on this occasion took possession of its former situations in the feet, and in the other knee; with but little appearance of inflammation, and chiefly affected the bursæ and the tendons. I visited this gentleman under this complicated state of suffering; and knew from experience that his recovery would necessarily be very tedious. He complained of clammy perspirations, attended with coldness. He was languid and debilitated. His appetite was not deficient; but the bowels were inert, and the action of the liver was torpid, as was indicated by the whitish appearance of the fæces, which were passed chiefly in pellets. The urine deposited pink sediment; although that which was secreted in the night was most commonly transparent. I viewed this circumstance as the index of a quick digestion in a nervous constitution, the digestive organs being at the same time unhealthy. I shall avoid the tedious details of treatment, and briefly mention, that each other night an alterative dose of hydr. submur. was given in conjunction with extract. colocynth. comp. and extract. hyoscyam.; the former being changed in about a fortnight for pilul. hydrarg. The most satisfactory relief of the rheumatic pains was derived from the use of gutt. nigr. administered at three intervals during the day, and partly in admixture with the draught, p. 186. The limbs were treated on the principles which

I have so often described. When inflammation, requiring the use of the lotion, was removed, the joints were surrounded with empl. saponis, spread on leather; which appeared to afford the double advantage of causing a favourable moisture on the surface, and of protection from the influence of variable temperature. In the next stage of the treatment, salt water sponging; friction with the liniment containing tinctura lyttæ; and flannel bandage, were the means employed; and, finally, friction by a skilful rubber completed the cure. The restoration of the constitution was assisted by change of air to the sea-side; and by blending the use of tonic medicine with judicious regimen. I must add, however, that a period of three or four months elapsed before the recovery was perfectly accomplished.

The two following cases I shall relate very concisely, for the sake of *diagnosis* alone.

CASE XX.

A gentleman, aged 45, of middle height and bulk, well formed, his chest circular, of the nervous temperament, when at the age of 24, suffered much from chronic rheumatism; and at 43 was first attacked with gout in the great toe, inheriting the disposition from his father. He has since had repeated fits, affecting both feet. His habits of living had constantly been very intemperate, even in the use of spirits; and his exercise in hunting violent. About two years and a half ago, he was repeatedly exposed to wet and cold, accustoming himself to travel on the outside of a stage in wet weather, insufficiently protected by clothing. In the origin of the unfortunate disease which forms the subject of my

present communication, and which commenced after this exposure, he complained of pain between the scapulæ passing through to the chest, as if of rheumatism. No relief was afforded by medicine, and the symptoms gradually increased. The pain was sometimes so acute, as to convey to his mind the idea "of hot needles being suddenly stuck into the flesh." At other times he felt a severe deep-seated gnawing pain, attended with the sensation of having been bruised; and not unfrequently as if a heavy weight were laid upon the chest. Very aching sensations affected the arms. On pressure in the epigastric region, he not only complained of much tenderness at that part, but also suffered increased uneasiness between the scapulæ. He declared that he had never felt the least respite from his sufferings either day or night, but on two occasions, when labouring under severe symptoms of acute gout. It is particularly worthy of observation, that so long as the urgency of the paroxysm continued, it served to suspend the pain which affected the chest; but which returned again as the symptoms of gout abated. Medical treatment was wholly unavailing, except that sedatives and rest now and then seemed to have a slight palliative effect. Some months after this period, a prominence on the left side of the sternum was discovered, attended with a strong pulsation. A pulsating tumour by degrees became very evident; and the melancholy case shewed itself to be unequivocally an aneurism of the aorta.

CASE XXI.

A. B. aged 41, robust and plethoric, but of the nervous temperament, suspected himself to have been

affected with gout, by right of inheritance from his father, under the following circumstances.

A year ago, when having gonorrhœa in its advanced stage, he was suddenly attacked with a swelling of the little finger, and, in a few days after, in the following parts, in quick succession:—the right knee, right instep, left knee, and the left instep. The parts were swollen, free from discolouration, and not painful; but he was rendered completely disabled from walking for three weeks. The discharge from the urethra ceased immediately upon the above attack becoming general, and never afterwards returned. He mentions that he was singularly languid and weak, and imputed the attack itself to the active effects which the use of Cheltenham water had produced upon his bowels.

On a second occasion, gonorrhœa having first existed, he suffered a similar attack to the preceding, on the morning after having used a hot bath. He became suddenly so much disabled, that he could not move either hand or foot. On this occasion, the discharge from the urethra was not suspended by the affection of the limbs.

A third time, after having laboured under gonorrhœal infection about a fortnight, and being under the active influence of aperient medicine, he was suddenly seized with a swelling of the knee, which did not cause pain, nor tenderness even to pressure; but the stiffness was extreme, and he could scarcely bear upon the limb. The discharge from the urethra continued. I saw this gentleman under these circumstances; and from the characters of disease which I found, and from his accurate description of his former attacks, I was convinced that he had never been affected by gout; which was the name always before given to the disorder. The swelling in this instance, and which he said resembled the former

appearances, was entirely bursal. The distension was very great. It seems, therefore, that in certain constitutions, a diseased condition of the bursal membrane, in one or more parts of the body, ensues as a sequel to gonorrhœa; giving rise very suddenly to an increased secretion of the synovial-like fluid of the bursa, and to much consequent distension, and lameness. In regard to the treatment in this case, I shall briefly mention that a mixture, consisting of balsam. copaiv. mist. amygd. and vin. colch. appeared useful, and certainly influenced the state of the urethra very favourably. A blister was applied to the knee with advantage; and when the skin was healed, friction with a stimulating liniment, and the use of a bandage, seemed to assist in completing the cure. As on the former occasions, a remarkable degree of languor and debility were experienced for some time; but he derived sensible benefit from a mixture of decoction of cinchona with an aromatic tincture, and the carbonate of soda; lemon juice being added at the time, so as to produce effervescence.

I shall conclude the general subject of Chronic Gout, by a brief discussion of the treatment

OF THE GOUTY CONCRETIONS.

THESE concretions, of which I have already made mention at p. 47, 115, 116, 145, and in cases i, ii, vi, xiii, xiv, xv, were described by the ancients as constituting the tophaceous* kind of gout. Sydenham, after all the pre-

* A general term to express concretion, derived from the Hebrew.

ceeding humoral pathologists, believed them to consist of indigested gouty matter thrown upon the joints, and changed into their peculiar state of hardness by the heat and pain of the joint. Van Swieten, on the same subject, speaks of this chalky matter as being formerly in a condition to circulate through the vessels, and views it as a deposit from the circulation. Believing the concretion to be of a real chalky nature, he mentions that it proves soluble in acids, and advises the muriatic in conjunction with the oil of turpentine as a solvent. It is truly curious, that, from asserted experiment, he should recommend, as a remedy, that acid in which the uric compound would be the least soluble. He afterwards however remarks, that alkaline applications had been more generally employed; and he says also by himself, with much success. Such were the unsettled conclusions into which this eminent man was led by the ignorance of chemistry, which was universal at that day.

In the first deposit of the uric compound which constitutes these concretions, it may be much, if not altogether, in the power of remedy to obviate the inconveniences which neglect would certainly produce.

From the easy solubility of the uric acid in pure potash liquor, I was led to the employment of this medicine as an external application; and in three instances of recent deposit, it was so successful, that the concretion which had been visible under the skin, became gradually removed. I have directed it in dilution with an equal quantity of recently prepared almond milk, to be applied by means of friction, two or three times in the day. Should this degree of strength irritate the skin, it may be more diluted; but I have usually found that it is borne in these proportions without inconvenience.

When the concretions have been of long standing,

and are much indurated, their absorption is with difficulty excited, and perhaps cannot be accomplished.

Even this case however must not be abandoned. A patient who came under my care, had long suffered pain and occasional inflammation from enlarged bursæ mucosæ in each hand. Their extreme distension and hardness gave the appearance of large bony tumours. On examination, it was evident that the bursæ were filled with uric matter. The use of the alkaline liniment, in a short time, was so far efficacious in lessening the size of the tumours, that the tightened skin became relaxed, and the use of the fingers was much improved.

In three cases, in which the concretion had partially forced itself from the cellular membrane and cutis through the cuticle, the application succeeded in causing the removal of the remaining deposit.

As foreign bodies, these concretions occasionally produce ulcerative irritation, and consequent sores; requiring surgical treatment*. Mr. Hunter remarks of them, "they leave the parts not easily excited to inflammation; the chalk shall remain for years without producing inflammation, and seldom produce it at all, but from quantity†."

A gentleman, aged 63, who has suffered from acquired gout, for the last thirty-three years, has several ganglions in the fingers; and in two of the bursæ there is an evident deposit of chalk-stone, or, to speak cor-

* See a Paper on this subject, by Mr. Moore.—Medical and Chirurgical Transactions, vol. i. p. 112.

† Mr. Brodie informs me, that he has succeeded in conquering this morbid action of the vessels, by the occasional application of the argentum nitratum to the diseased open surface.

rectly, of the uric compound. He has two ulcerations in each heel, and a constant secretion of this kind is taking place. The parts ache very much in cold and in changeable weather; and give him a strong warning, if gout is at hand. When, from much walking, increased tenderness has been produced, a simple bread-poultice produces relief; and, at all other times, the emplastrum saponis, spread on leather, affords a comfortable defence. Doubtless the true method of cure in such a case, is to cut away the thickened cuticle; and, arriving at or near the diseased surface, to make a judicious application of argent. nitrat.

In cases xiii, xiv, and xv, p. 325, 328, and 329, I have related how favourably an uric abscess allows of the lancet; and that it does so, even when the bursa mucosa suppurates. This treatment comes within the province of the surgeon; but the brief general history which I have given, may be useful.

In regard to the constitutional treatment in this peculiar disposition of the exhalant vessels, it seems to me very doubtful whether any medicines will be found to have a specific operation, as chemical agents; although, I confess, that, in the practice which I am about to suggest, I hold this principle partly in view. The digestive functions of those persons, in whom this morbid process is going on, are usually weak and irregular; and they are much disposed to acidity of stomach. I believe, in every instance in which I have seen these concretions, that the liver has been more or less diseased; and a corresponding treatment should therefore be pursued, when such indications are presented. In reference to the *ricarious* secretion of uric acid now under consideration (if I may adopt this hypothesis), the powers of magnesia, in

conjunction with liquor potassæ, may with propriety receive a trial. The following formula may be offered :

R Magnes. gr. x. ad ʒi.

Mist. amygd. ʒxiv.

Liquor potass. ℥ xx. ad ʒi.

Syr. tolutani ʒi.—M.

Fiat haustus, bis quotidie sumendus.

An intelligent gouty gentleman informs me, that many years ago, he was troubled with chalk-stones in several fingers ; and that from one finger there was an occasional oozing of chalk-like matter. He adds, “that by means of a course of magnesia, taken in regular daily doses, all the chalk-stones gradually disappeared.” Upon examination of the fingers, I find, at present, only a very slight trace of concretion in one of the bursæ. Patients are so often deceived in the belief of having *chalk-stones*, in consequence of the knotted state of their tendons, or the induration of the small bursæ mucosæ, that I cannot allow myself to receive the account of this cure with implicit confidence.

In cases of this description, a steady perseverance in any means which are adopted is quite essential. In several instances of concretions which were existing in connexion with very unhealthy functions of the liver, the mild alterative course of treatment which was adopted, had a decided influence in promoting the absorption of the uric matter ; and from such share of experience as I have had in this peculiar deviation from the ordinary course of gout, I can venture to affirm, that much good may be afforded by an attentive plan of medicine, and of regimen.

Sydenham, when treating on this part of the subject, makes the following observation, which, although not founded on correct pathology, deserves considerate attention: "I have experienced, in my own particular case, that not only the generation of these concretions may be prevented by daily and long-continued exercise, which duly distributes the gouty humours through the whole body, that otherwise attack a particular part; but it also dissolves old and indurated concretions, provided they do not come to such a degree, as to change the external skin into their substance."

RETROCEDENT GOUT.



WHEN, during the existence of gouty inflammation, either in its acute or chronic form, a sudden cessation of the external action takes place, it sometimes happens that an internal organ becomes immediately and violently affected. When this event occurs in the height of a paroxysm, the symptoms are acute and run a rapid course; but when it arises in chronic gout, they are sometimes of slower progress. In either case, therefore, the symptoms bear a relation to the previous state of the system.

The retrocession of gout in the paroxysm, constituting the case of danger of which I am now speaking, is of rare occurrence, and probably never happens except from the patient's want of care, or from some injudicious management.

The transference is most disposed to affect the stomach or intestines; or both in succession.

The symptoms which attack the stomach are exquisite pain and spasm; and sickness is an usual attendant. If the intestines be more distinctly affected, enteritis in its worst form is produced; and vomiting, which most commonly occurs, is more or less urgent, accordingly as the seat of disease is near or distant from the stomach. In either case the danger is pressing; and, unless relief be speedily rendered, *death* soon closes the scene.

If the transference take place to the brain, in its worst form, apoplexy is produced ; and, in all probability, proves of fatal termination. The head is sometimes affected with pain and distressing sensations in connexion with the gouty diathesis, even before the first occurrence of a paroxysm, as is shewn in the following case.

A gentleman, of the nervous temperament, previously to his first attack of gout, which occurred at the age of 27, had been for a few weeks complaining, as he expressed it, of oppression of the brain, remarkably affecting his powers of mind and spirits ; and occasionally, also, he suffered severe head-ach. He took purgative medicines, and was cupped two or three times with advantage ; but was not cured. One evening, his head became suddenly relieved, and the great toe of one foot as suddenly affected with symptoms of gout. The exchange was almost instantaneous.

During the gout, the head and stomach become peculiarly affected in certain constitutions.

A lady, entirely of the nervous temperament, relates to me that she has suffered much from the head and stomach at the period of having the gout. She experienced lancinating pains of the head, and a remarkable sense of depression at the pit of the stomach. Gout taking place in the limbs, relieved all those symptoms ; but if the gout ceased, they returned again immediately.

A gentleman describes, that when his feet were inflamed with gout, his head was suddenly seized with intense pain and great confusion ; the gout, however, not quitting the feet. He procured relief by fomenting the head with hot brandy and water, and afterwards applying flannel, so as to produce perspiration.

Dr. Cullen observes, under the division of Retrocedent Gout, that “ sometimes the internal part is the

heart, which gives occasion to a syncope; sometimes it is the lungs, which are affected with asthma."

There is an apparent cause, why these organs should be less liable to be affected than the brain; namely, the greater sympathy subsisting between it and gouty parts; and more particularly from the great tendency of a determination of blood to the head, in those who have long been subject to gout; and that the alimentary canal should most commonly become the seat of the retrocedent action, might be expected from the active sympathetic connexion so often subsisting between it and the extremities, during the phenomena of gouty inflammation.

With respect to the retrocession in question to the heart or lungs, I am not acquainted with any clear facts of its occurrence. In persons who are subject both to gout and asthma, the existence of either one or the other taking place with some degree of alternation, must not be considered as a fair example of retrocession; according to the definition which I have given. In case of the lungs becoming the part affected in repelled gout, I should expect that inflammation and not asthma (unless in an asthmatic person) would be the actual form of complaint.

A question of theory has sometimes been agitated, as to the propriety of the definition, *retrocession*. Mr. Hunter observes, "I should be inclined to suppose, that its effects on the brain or stomach are not similar to those on the extremity, or probably it does not advance so far in its effects to them; or it would certainly kill."

It appears to me an useless discussion in a practical point of view, to argue upon the identity of the phenomena, in these particular situations. We see that the certain event follows the certain antecedent; and hence

I conceive that the established opinion of an occasional *transference* of inflammatory and spasmodic action from external to certain internal parts, in gout, is clearly enough made out; and that it is important in practice, as well as admissible in doctrine.

Dr. Cullen states two other affections, the one of the neck of the bladder, producing pain, strangury, and a catarrhus vesicæ; the other of the rectum, sometimes causing pain alone in that part, and sometimes by hæmorrhoidal swellings there. He adds, "In gouty persons I have known such affections alternate with inflammatory affections of the joints: but whether to refer those affections to the retrocedent, or to the misplaced gout, I will not presume to determine."

In two patients, I have seen that chronic inflammation of the prostate gland has been so curiously modified by the gouty diathesis, that the surgeon has pronounced the gland to be affected with gout.

That gouty persons are remarkably subject to hæmorrhoidal affections, and an irritable state of the bladder and urethra, I have already admitted and explained; but in connexion with the paroxysm, I have only observed an occasional increase of sympathy in a high degree; in which case, I conceive that the terms misplaced or retrocedent gout, as applied to such affections, express much more than is warranted by the phenomena.

In two instances I have known a painful affection of the testes and scrotum assume a very doubtful character, so as to perplex the judgment of the surgeon for many weeks in succession; and at last entirely give way to an unexpected occurrence of acute gout.

One gentleman relates to me, that for the last three years he has been distressed with an irritable state of bladder, sometimes giving rise to urgent inconvenience;

but that the symptoms were remarkably suspended by his last fit of gout. Now, he rises five or six times in the night; but at the time of the paroxysm, not more than twice. I strongly suspect that in this instance there is a calculous concretion in the bladder.

I have observed that those gouty persons who are most disposed to be painfully affected with gravel, are also the most liable to spasmodic attacks of the diaphragm or abdominal muscles; and which the patient always describes as gout in the stomach. I have seen this occurrence chiefly happen in the absence of the paroxysm, and therefore should call it a spasmodic affection, or spasm mixed with inflammatory action, in the gouty constitution, but not retrocedent gout.

Upon these and other anomalous forms of complaint, which are partly inflammatory, and partly spasmodic in their nature, it seems both just and sufficient to consider, that a modification in the symptoms is produced by the influence of the gouty habit. Of this point, and of the nervous character of the gouty constitution or temperament, I have already treated at some length.

The subject of conversions of diseases, or the superseding of one disease by another, is matter of curious and important inquiry. In the cases which I have recently related, I have shewn the extraordinary power of gout in suspending or removing long established morbid actions. I may add a few further examples.

A gentleman of highly irritable constitution, after an exposure of some hours to wet and cold, was attacked with erysipelas in the face. The symptoms were severe for a few days, but yielded quickly to the gout, which took place in the feet.

A gentleman, for many years subject to gout, was exposed to cold on the 4th of the month. In the evening

and upon the following day he suffered from severe pains in the chest, with difficulty of breathing. On the 6th, gout began in one foot, which increased on the following day in a great degree, and his chest became entirely relieved.

J. B. has long been subject to a purulent secretion from the ear. An attack of gout, which he lately had for the first time, wholly suspended the discharge; but it returned as soon as the gout quitted.

A gentleman, who has suffered from gout for many years, informs me, that in the early part of his life he had occasional attacks of erysipelas; but that since the gout, he has been free from that complaint.

In some instances, from a change in the state of the constitution, gout fails to return; or is suspended by another disease, to which the individual has acquired a stronger tendency.

A gentleman was subject to gout during several years of his life; but, for the last seven, he has been affected only with asthma, which succeeded to the gout.

I proceed now to consider the remote causes in relation to the retrocession.

CAUSES.

THE most frequent cause which produces retrocedent gout in its most genuine form, is sudden vicissitude of temperature applied to the body generally; or cold, more or less continued, offered to the affected parts.

Professor Home, of Edinburgh, in his lectures, relates the case of a gentleman, who exposed himself to the influence of wet and cold, when the gout was slightly pre-

sent in the feet ; and on the same afternoon, enteritis followed, which in twelve hours proved fatal.

Dr. Parry informed me some time ago, "that in the same winter he had seen two instances of extravasation in the brain, from the removing of gout in the extremities by immersing the feet affected in cold water."

A gentleman having a relapse of gout in the foot, in consequence of a strain, applied a cold lotion very freely, thinking that the inflammation being thus produced might be of the common kind ; but the pain was so much aggravated, and his general feelings were so unusual, that he was afraid of internal gout, and soon desisted from its use.

Another gentleman applied to the inflamed parts a cold lotion, consisting of equal parts of alcohol and liquor ammon. acet. with seeming relief to the inflammation ; but at length he felt so alarming a spasm at the stomach, that he laid aside the remedy in fear.

An elderly gentleman, severely gouty, of the nervous temperament, on one occasion, when both feet were much inflamed, applied laudanum, hartshorn, and spirit of turpentine. He says, "that he repelled the gouty action from the feet, and that his head became immediately affected, feeling as if a wooden wedge had been driven in between the bones, and was forcing off the crown of the head. In the course of a few hours the gout returned to the feet, and the head was at once relieved."

It is with much satisfaction I state, that, in a subsequent severe paroxysm, this patient made free use of the evaporating lotion, p. 254, with perfectly good effect.

Lately, I saw a gentleman, who, when slightly affected with gouty inflammation in his feet, walked on cold damp ground ; and his stomach quickly became so severely

pained, that, but for timely remedy, the event would have been uncertain.

When cold is the hurtful agent, the internal symptoms which are produced are probably, for the most part, of an inflammatory nature. I have formed this opinion from such cases as have come under my own observation; and from the general information which I have collected.

The blood being checked from the surface, while the gouty diathesis is present, a preternatural determination to some internal organ succeeds; and inflammatory action arises, which is marked by the utmost intensity of symptoms, and a rapidity of course that is almost peculiar. If, however, the retrocession take place after the free employment of all the means which reduce inflammatory action, notwithstanding that cold may have been the exciting cause, the symptoms which arise will probably prove of a spasmodic nature. This appears to be illustrated in case x, at p. 313.

I must also add, that in those constitutions in which the nervous temperament strongly prevails, symptoms of pure spasm will often arise, although the exciting cause shall have been an exposure to cold; or the influence of cold in any way applied.

Phenomena, in part similar, appear to be produced by the occasional agency of certain stimuli, in suspending the external gouty action, as in the operation of hellebore; or as seen in the effects sometimes produced by the *eau medicinale*. In cases of this description, however, the noxious cause being applied internally, there is some obvious difference in the beginning and progress of the symptoms. The increased determination of blood to the internal organ, is here the *first* event in the series of the diseased actions.

The influence of indigestible food produces one form of the disease; but in this example I have seen that the pain, which is intensely severe, is dependant rather upon spasm, than upon active inflammatory action. Sickness is a certain attendant; and I believe that the stomach, under the present cause, is always, in the first instance, the entire seat of the disease.

I must repeat my observation, that the character of the attack, as to inflammation or spasm, materially depends on the temperament of the individual; and an investigation regarding this should not be neglected. I shall presently relate some cases which exemplify this point.

Violent passions of the mind may possibly induce retrocedent gout; but I am not acquainted with any example of the fact.

DIAGNOSIS.

To attempt a practical distinction between the internal disease of which I am now treating, and similar forms of disease where gout has not been present, or in persons not gouty, is probably both useless and improper; as it involves theoretical considerations of too delicate a nature to admit of satisfactory reasoning; while the treatment must be prompt and vigorous, and strictly adapted to the actual symptoms.

In order to possess a power of correct discrimination upon the question, whether the case shall be considered simply as one of spasm, or of spasm and inflammation mixed, or of pure inflammation; our minds should be thoroughly stored with sound principles of pathology. Useful conclusions may be derived from a reference to the exciting cause, on the principles which I have stated.

In an attack purely spasmodic, the rigidly contracted state of the abdominal muscles, and the relief which is afforded by strong pressure, are very distinctive. When it is purely inflammatory, the tender state of parts to the slightest weight or pressure; the more regular diffusion, yet greater fixedness of the pain; the sympathetic fever which is instantly produced; and indeed the very physiognomy of the patient in the comparative situations of attack, will, to the experienced practitioner, be a description of the nature of the disease. The state of the pulse, as whether small and indistinct, or full and oppressed, or in vigorous action in any way, will materially direct the judgment; and the state of the skin and features, whether cold and collapsed, or in contrary states, is a guidance of importance. Dr. Cullen and authors in general appear to have considered it as a settled axiom of practical doctrine, that debility and spasm, and not inflammatory action, seize the internal organ in the case of retrocedent gout.

A perfect conviction prevails in my mind, that, in a genuine example of retrocedent gout to an internal organ, inflammatory action is the more common occurrence, and that spasm alone is comparatively rare. The mixed action of spasm and inflammation may, however, be expected to happen still the most. In reference to inflammatory attacks, we see that many of the cases related by authors have evidently terminated in gangrene*.

* Morgagni, Ep. 57.—Rush's Inquiries and Observations, vol. v. p. 153.—According to these authorities, the *black vomit* occurred before death. In the Hunterian Museum there is a preparation (No. 52) of a gouty stomach, thus described: "A specimen of a portion of the œsophagus and stomach of a person

OF THE TREATMENT.

IN conformity to the state of the fact which I have been now discussing, the means of practice are to be determined.

The life of our patient hangs on the discrimination which we exert. In every case of retrocedent gout, Dr. Cullen† has directed a treatment entirely stimulating; and probably the same doctrine is prevalent in the schools of medicine, and consequently in general practice.

It is to be admitted, that in some particular constitutions, or in certain nervous states of the constitution of any individual, simple *spasm*, either affecting the stomach or the diaphragm, of the most intense kind, is the instantaneous or speedy effect of some offending agent, accompanied with an abatement or cessation of the gouty inflammation in the extremities. In this case the patient feels some benefit from strong pressure applied to the stomach and abdomen; and the pulse is contracted. He almost instinctively flies to brandy, or hot brandy and water, for relief; and, in such an example as this, very probably with every advantage. In an urgent attack of this kind, the state of the patient appears indeed truly

who died suddenly of the gout in his stomach. There was considerable inflammation, even, in some places, to the extravasation of blood."

† Having spoken of strong wines joined with aromatics, and to be given warm, when the stomach and intestines are the seat of disease, he adds: "If these should not prove powerful enough, ardent spirits must be employed, and are to be given in a larger dose."

alarming; and the most powerful remedies at hand are felt to be the only expedients to save life. I am acquainted with some striking instances of this kind, in which the stimulant practice, carried to a free extent, succeeded; but this description of pure spasm, in my opinion, forms the only exception to a very different general rule of practice, which I would propose in opposition to that laid down by Dr. Cullen.

If retrocession have been excited by indigestible food, the sickness which is present, and the appearance of the rejected matter, point out that the vomiting should be promoted. An emetic of ipecacuanha is well adapted for this purpose; and its operation is to be assisted by draughts of warm water in the usual manner. If the pain be thus relieved, the bowels should next be acted upon, and five or ten grains of calomel should be given as an immediate dose. As soon as the stomach can retain a purgative medicine, the draught, p. 186, may be given every three hours, until a full operation is produced; or the combination of sulphate of magnesia and infusion of senna with an aromatic tincture, may by some be preferred. Whatever medicine of this description may be chosen, the dose should be repeated at short intervals, until a full effect is produced. The aid of an injection will occasionally be required. If, however, violent pain should still continue, after the stomach has been cleared of its contents, tincture of opium, in a dose from forty to one hundred drops, may be given without hesitation; it must be repeated also in free doses, at an interval of half an hour, or an hour, until pain and spasm cease, or satisfactorily abate; and at the same time, purgatives which will have their effect delayed but not prevented by the opiate, must, on no account, be omitted. Pills of calomel, colocynth, and soap, constitute the form of active

remedy which is most to be recommended; and the fluid purgative should follow their administration.

When the pain has ceased, and the circulating and nervous powers have so far recovered, that the *re-action* of the system produces its effects, we should be upon our guard, lest inflammation take the place of spasm, and insidiously prevail, unseen and unrelieved.

When exposure to cold, or the influence of cold in any way, or violent stimuli, have been the exciting causes of injury, in a patient whose circulation is strong; and in proportion as his habit is plethoric, we have to expect that the disease will be inflammatory; and, accordingly, as the symptoms are marked and violent, sixteen, twenty, or thirty ounces of blood should be instantly taken from the arm; the quantity and repetition being adapted to the several indications and circumstances. The inflamed bowels must be treated, as in ordinary enteritis, with equal promptness and decision. The rules of practice, in relation to this point, are too well known to require being detailed. With regard to collateral treatment, however, something further may be offered. The attempt to solicit back the gout to the extremities (or, probably, in more correct language, to divert the diseased action from the vital organ by remote excitement) will be made with great propriety by sinapisms; or by topical warmth to the limbs; as by a stimulating fomentation; a pediluvium of hot water simply, or containing mustard and salt; by warm covering, and such means: but to the bowels themselves, as in the treatment of common enteritis, I am induced to prefer the use of a *tepid* application to one of a higher temperature. With this view, upon the principle of evaporation, the lotion of camphor and alcohol (p. 254) applied just tepid, or at that temperature which is felt to be the most comfortable, con-

stantly and universally to the abdomen by means of folded cloths, constitute a remedy which may be adopted with advantage.

In some cases of severe enteritis, I have had decisive proof of the superiority of its effects over the ordinary method of hot fomentations.

In those instances in which the general powers of the circulation are weak, in connexion with visceral inflammation, local cupping may be preferred to the more general detraction of blood; or when this valuable remedy cannot be conveniently put in practice, the free use of leeches must be the substitute.

In reference to the fact, that gouty persons, in the intervals of their fits, are sometimes seized with violent pain at the stomach, and spasm that seems to threaten life, the most powerful stimuli alone giving relief, I must remark, that such occurrences are not examples of *retrocession*.

A lady, possessing the true nervous temperament, just as she was convalescent from the paroxysm, imprudently exposed herself to the evening air, walking on damp ground. Soon after entering the house, she was seized with sudden and violent spasm of the diaphragm. She was free from sickness, but breathed with exceeding difficulty. She thought "that she should have expired with the pain." She took hot brandy and water, and applied warmth to the feet, with speedy relief.

If apoplexy be the disease which succeeds to the retrocession of gout, copious bleeding, to the extent that the pulse permits, is the only remedy which can save the patient; and if no rupture of vessel have taken place, it will most probably be effectual. Cold water, or, with still more effect, a cold evaporating lotion, freely and constantly applied to the head by means of folded cloths; and warmth and stimulus to the feet, according to some

of the methods already stated; form important aids of treatment. To speak of purgative means, and the general management, would be only to repeat the well-known rules of general practice, and to fatigue the attention of the reader.

Should the viscera of the thorax be affected, the usual principles of practice are to be followed.

The following cases will serve to shew the distinction between the symptoms of spasm and inflammation, and the relative treatment which is to be employed.

CASE I.

S. R. aged 47, corpulent, of bilious diathesis, and of the nervous temperament in a most exquisite degree. Was first attacked with *acquired* gout in the external part of the foot at 22. Seven years ago took the *eau medicinale*, in the dose of half a bottle, two nights in succession. It did not materially relieve the pain, and induced a feebleness of limbs almost amounting to paralysis. He never afterwards recovered the comfortable use of them, which on all former occasions he had done. He has suffered from gout to a degree of martyrdom; and when under the influence of the paroxysm, his nervous system is affected in the most distressing degree; having, together with intense pain, unaccountable horrors and apprehensions. Seven months ago he was advised to discontinue the use of wine, which he had been in the habit of taking regularly, and often convivially. He felt the loss of this cordial stimulus in a very sensible degree. He said, expressively, that his nervousness was so much increased, that he was afraid to meet a friend. He would prefer to turn away, in order to avoid conversation. At this period, therefore, seven months having elapsed, he

was exposed to wet and cold, and the feet were slightly wet. On that day he did not suffer inconvenience; but on the following night, having only a slight degree of gout in the feet, he was suddenly seized with excruciating pain in the stomach, and such an oppression from flatus that he thought the stomach would actually burst. He was relieved from these alarming symptoms by the use of hot brandy and water; æther and opium; and fomentations of hot water applied to the stomach. A severe paroxysm of gout in the feet and knees soon followed. I saw this gentleman, on a subsequent occasion, under the influence of spasmodic symptoms, which were induced by cold. An extraordinary torpor appeared to have come over the whole frame. The pulse was very slow and languid; the skin was cold; the features were collapsed; the expression of the countenance was that of terror mixed with a degree of insensibility; the breathing was laboured; the muscles about the throat and trachea were so much affected, that the act of swallowing and speaking was rendered very difficult. Stimulants, as on the former occasion, were here employed with success. Towards the relief of a severely painful state of gout in the limbs, for which, soon after, I had occasion to prescribe, the application of the poultice made with the evaporating lotion agreed perfectly, and was very beneficial; and also the occasional sponging of the part with the warm lotion had a good effect: but with a patient of such peculiar tendency to retrocession, I would not adopt my usual method of more active evaporation. This was a case of exception to the usual management of the paroxysm in regard to diet; as I found it expedient to allow the use of Madeira daily, to the extent of two or three glasses, even during the height of the painful symptoms.

CASE II.

R. C. æt. 52, of the nervous temperament, was for the first time affected with gout in the foot at 31. When the symptoms were yet urgent, after eating rather heartily and drinking cold Madeira and water, he was for a short time exposed to cold air. He was suddenly seized with cold perspirations in the limbs, attended with a sense of heat arising up from the bowels, and an alarming oppression affecting the breath, but unattended with any pain. The foot became immediately easy. He experienced a slight degree of sickness; but he was most alarmed with sensations of sinking, to the degree even of apprehending a fatal fainting. The palpitation of the heart was excessive. He drank warm water as being the readiest expedient, and produced vomiting. Much acid matter was rejected from his stomach, and with immediate relief. The attack in question ended with a return of active gout to the foot. On a subsequent occasion he suffered in a similar manner in the paroxysm. About seven in the evening the pain in the foot was very severe. At twelve, having suffered much from pain and restlessness, his stomach was suddenly seized in the same manner as before, and with the former attendant nervous sensations. His foot was as immediately relieved. Without delay he took an emetic; and, discharging from his stomach a large quantity of acid matter and mucus, of green appearance, he became relieved. In the morning the gout again took possession of the foot.

In this case, the timely employment of purgative treatment would, in all probability, have prevented these attacks; which, if not attended with real danger, were

highly alarming to the patient. He had not interfered with the natural course of the gout.

CASE III.

C. A. aged 64, of full habit and inclined to corpulence, of bilious diathesis, and of the nervous temperament, was affected with hereditary gout first at 63, and has suffered no less than six attacks within twelve months. Much subject to cramps in the legs. Formerly had jaundice; with such urgent symptoms, that obstruction of the duct by a gall stone was apprehended. After having pursued a course of medicines at home, he went to Cheltenham, and derived much advantage from the water in regard to the bilious obstruction; but injudiciously continuing to take it without intermission for five weeks, he brought back his *dyspepsia*, which proved very troublesome. Previously to his first attack of gout, he laboured under a very troublesome state of dyspepsia, a cutaneous complaint, and cough. He hailed the arrival of gout, therefore, as an expected remedy for such complicated indisposition; and would not, of course, interfere with its progress. Observe, however, the failure of Nature to relieve the constitution, even by six paroxysms of gout in one year! and, in the last, the stomach became affected as follows.

The usual symptoms of gout in the feet had continued about a week. They were attended with dyspepsia and some vertigo. He had taken a purgative medicine; had fatigued himself by over-exertion; and at his dinner had eaten more of solid food than was judicious. He was awoke about three in the morning with a sense of cramp at the stomach, and a hiccup which continued for several

hours. He became relieved on the following day, but felt much languor and depression. In the evening he complained of general coldness, which was soon followed by an urgent sense of oppression at the stomach, feelings of cramp across the diaphragm, hiccup, and a degree of burning in the chest. The complexion was rather jaundiced. His immediate symptoms were relieved by a mixture consisting of conf. aromat. mist. camph. sp. ætheris sulph. comp. et tinct. sennæ; but the most sensible advantage was obtained from the active operation of a calomel purgative. The attack returned for two or three nights in an abated degree. Each time it was unattended with sickness. It left a considerable soreness about the stomach. The appearance of the excretions fully pointed out the necessity of purgative treatment; and in this, the radical cure consisted. The gout in the feet gradually declined; and I should observe, that no local treatment had been used. The active use of remedies was followed by the adoption of an alterative course of medicine; and at the end of a month this gentleman was perfectly recovered.

CASE IV.

A gentleman, aged 50, of the nervous temperament, first affected with *hereditary* gout in the great toe at 30; and every year since, has had one or two severe fits. Towards the close of the last attack, he imprudently applied a cold lotion to the foot; and almost immediately experienced a severe pain in the chest and in the shoulder. He drank hot brandy and water with speedy relief. Two days after, most inadvertently, he again used a lotion cold, and with the same consequences. He then re-

sorted to a powerful dose of essence of ginger, which caused an expulsion of flatus from the stomach, and afforded relief to the pains. He obtained, however, his more decided amendment, and final cure, from the steady employment of aperient and alterative medicines, used in conjunction with regulated regimen.

CASE V.

L. S. aged 52, rather plethoric, but of the nervous temperament, first affected with hereditary gout at 29; subject to cramps in the muscles of the legs. On one occasion he applied twelve leeches to the feet, and, during the promoting of the bleeding in the usual manner, he was suddenly seized with an excruciating pain at the pit of the stomach. He instantly, and as if by instinct, drank half a pint of brandy, and with immediate relief. He next encouraged the gouty inflammation in the foot, on the principle of keeping the disease in that part. Lately he exposed himself to a cold east wind, unprotected by a great coat. Rheumatic pain immediately affected the shoulder and scapula, and gout followed in a few days. He states that toothach, without any previous exposure to cold, has, more than once, preceded a paroxysm.

He relates, that on another occasion, after a tedious fit, he reduced his regimen to a great degree of abstinence, and suffered a severe spasmodic attack of the stomach, which yielded to a draught of brandy. This gentleman has been much subject to dyspepsia, and to bilious obstruction.

CASE VI.

R. G. aged 65, corpulent, plethoric, and of the mixed temperament, first affected with acquired gout at 23; and since has been a severe martyr to the disease. At the close of a long paroxysm he exposed himself to cold, and soon experienced an intense pain of the head, with great confusion and occasional delirium. In addition to the use of purgative medicine, he was bled largely from the arm. The head became relieved; but next the abdomen was affected, the pain being severe, and the tenderness even to slight pressure was so great as to point out the presence of inflammation. During these symptoms gout suddenly and entirely quitted the feet, and did not afterwards return. The bleeding was freely repeated, and he was also cupped upon the abdomen. A very large blister was next applied. These means, with the continuance of purgative medicine, soon effected a removal of every alarming symptom.

CASE VII.

R. B. aged 50, corpulent, plethoric, and of the mixed temperament, was first affected with *hereditary* gout at 22, in the ball of the great toe. Has since been a martyr to the disease. Two years ago first resorted to Reynolds' tincture, which, for a few trials, relieved him considerably; but his relapses became so frequent, that he lost his confidence in the medicine. Since the use of it he has found his stomach remarkably weak, and subject to the sensations which he calls spasm. Four months ago, when a paroxysm of gout in the feet had first ceased, and when

weakness only seemed to affect the limbs, he remained for a considerable time in a cold apartment, and was suddenly seized with a pain at the pit of the stomach, which gave him the sensation of being tied at that part with a strong cord. He drew himself in, and pressed his stomach with both his hands in the most forcible manner; using such and other muscular actions as this kind of spasmodic pain makes instinctive. He drank, in quick succession, eight tumblers of hot brandy and water, rendered still more stimulating by the addition of essence of ginger. He very soon obtained relief by this treatment.

A month since, I was first consulted by this gentleman, and found him lying in very severe pain, which affected the greater part of the abdomen. Gouty inflammation had existed in the feet and knees about a fortnight. As on the former occasion he had been in a cold apartment. The gout had suddenly, in a considerable degree, quitted the extremities. The pulse was hard, full, and strong. Now he took pains to reverse all the former actions of his muscles; and he lay upon his back with his knees a little bent. The abdomen was very tender to pressure. Bleeding from the arm, cupping upon the abdomen, and afterwards a large blister, with the use of pills of hydr. submur. pulv. jacob. extr. colocynth, compos. and the draught, p. 186, so freely administered that the bowels were kept in free action, were the remedies employed. I must add, however, that, for the purpose of removing the pain and soreness which remained, he took, with great advantage, the anodyne pills, p. 224, at stated intervals. An increase of gout at first affected the limbs, but this yielded to the influence of alterative aperients; and the patient recovered in the most favourable manner.

CASE VIII.

T. N. aged 55, disposed to occasional plethora, and of the mixed temperament, when affected by acute gout in one foot, took an active dose of calomel and James's powder. It appeared from what occurred, and from reference to a former occasion, that James's powder disagreed with this patient in a peculiar manner, producing excessive perspiration and faintness. Being accidentally out of bed, and when perspiring profusely, he fainted away, and was thus exposed to cold. In the course of the day a pain came on at the pit of the stomach, passing through to the back. He ate some light animal food at dinner; soon after, he was seized with vomiting, but this did not give relief. Mustard cataplasms were applied to the feet, although the gout had not receded from the affected part. The stimulating treatment was adopted very actively, and, in quick succession, hot brandy and water, antispasmodic medicines, hot fomentations, and the hot bath, were administered. The pain, however, continued, and the soreness of the abdomen very much increased. In this extremity, he was bled very largely from the arm, and with such immediate relief that it almost proved a cure.

From a review of these cases, it appears that, even when inflammatory action affects the abdominal viscera in the gouty patient, the symptoms have a different disposition from that which belongs to the same form of disease in one not gouty; and it is in our power to administer opium after depletion by bleeding, and the use of purgative medicine, when in other cases such practice would be inadmissible. The present examples point out how much the character of the retrocedent action is in-

fluenced by the temperament of the patient. The cases which I have offered would seem to show that spasmodic action is of more frequent occurrence than the inflammatory, in contradiction to my first position ; but I repeat, that in the most genuine form of retrocedent gout, namely, when the inflammation suddenly quits the extremities in consequence of the influence of cold, in a patient not previously reduced by treatment or by the continuance of disease ; and, more especially, if the temperament be more of the sanguineous than the nervous, the internal diseased action will be that of inflammation distinctly, or inflammation mixed with spasm. It appears to me of great importance that the rule of Dr. Cullen, to treat retrocedent gout with powerful stimulants, should be protested against, as absolute ; and that the danger which may appertain to such practice should be pointed out.

I ought, in justice to my own method of treating the paroxysm of gout, to add, that, in the cases of retrocession which I have here introduced, most of the patients have come under my care when affected with the regular acute symptoms, and have made use of the lotion and the other several means which I have detailed, without the least disadvantage, or any threatening of retrocession. I have been led, therefore, to conclude, that the chief cause of the retrocedent action has consisted in the neglect of proper purgative and alterative treatment ; in conjunction also with that injudicious use of stimulating diet, which places the patient in a state of morbid excitement, and creates the tendency to transference of disease from parts without, to that organ within, whose functions are in the greatest degree disturbed from the healthy course.

As an example of visceral inflammation supervening on the symptoms of chronic gout, and of the treatment which was successfully employed, I shall relate the following case :

CASE IX.

J. G. aged 58, tall, robust, and corpulent; has a circular chest; is very plethoric; of a sanguineo-nervous temperament; of bilious complexion; of free habits both in eating and drinking; formerly when he lived in families as butler, indulging in wine; but of late years in porter and spirits; not subject to other diseases; gout unknown in his family, although a numerous one. Has for many years suffered much from occasional cramps, both in the legs and the abdominal muscles; and sometimes, also, with spasm of the diaphragm. He became gouty at thirty-five. First attacked in the great toe of one foot only; but since, in knees, hands, and elbows; and adds, that in the paroxysm his head has sometimes been painful, even to delirium. Some fits warn him of their approach by unusual depression of spirits, and by uneasiness and increased weakness of the joints. Others make their invasion after midnight, when he has gone to bed in seeming health. Of late years has had a fit, both in autumn, and in January or February. States cold to be the usual exciting cause. The disease has become more and more severe in its progress.

In January last, was attacked in the feet successively, and in the hands afterwards. No crisis formed in this paroxysm. The symptoms, during the month preceding the attack now to be described, had been entirely chronic; the inflammation and pain also changing place frequently,

Describes that there was much œdema in the feet, and above the ankles, and also in the hands; much numbness of parts, and coldness and heat sometimes alternating; spirits extremely depressed; appetite good; bowels irregular; the faces foul and slimy. Under these circumstances, his limbs being more free from complaint than usual, he exposed himself, unprotected, to a cold east wind and rain. I found him at night, on the 20th of March, 1815, complaining of severe pain at the stomach, with considerable difficulty of breathing; the right hypochondrium and epigastric region tender to pressure. He related, that, for three or four previous days, he had been troubled with colic pains and a disordered state of bowels. On this day he had been costive. Urine scanty and high-coloured, depositing much pink sediment. The tongue much furred. Had been vomiting green acid matter, and was still sick. There was much abdominal fulness. The pulse 84, very full and strong; the veins generally much distended; the skin hot and dry. The complexion was considerably jaundiced, of a blackish yellow hue. He was immediately bled to \bar{xvi} . He took first an emetic of ipecacuanha; afterwards five grains of calomel; and a purgative draught of sulphate of magnesia, infusion and tincture of senna, every four hours. Thin gruel was ordered freely; and tea only in addition, with a little bread.

On the following day, 21st, much relief had been obtained. The blood had a thick buffy coat, with strongly contracted edges*. Much slimy matter had been dis-

* A comparative examination of the properties of the blood, in the circumstances of health and disease, would be a task of exceeding difficulty upon a sufficient scale for useful conclusions. Its relative specific gravity would be an interesting fact to be as-

charged from the stomach; and the fæces were dark and offensive. The skin remained hot, and the pulse was not abated in fulness or frequency. The bleeding was repeated to the same extent. The draught at p. 186, was directed every four or six hours; and the pills, according to the formula, p. 276, at bed-time.

On the 22d, reported that some pain had returned at the stomach on the preceding night, but less urgent. The draught continued.

On the 23d I found that he had passed a severe night. The pain, preceded by excessive coldness of the extremities, returned early in the evening, and got to its height about 2 A. M. No sickness, and the bowels freely open; now some discharge of fresh-secreted bile; but previously the appearance had been muddy, or sometimes as of yeast and water. A deep inspiration was painful to the right hypochondrium; some cough was present; a severe sense of heat was felt at the stomach.

certained, and this investigation is not of difficult accomplishment. I shall take a convenient opportunity to direct my attention to this point of inquiry. The appearance of the buffy coat (the fibrin) was referred by the ingenious Mr. Hewson to the slow coagulation of the blood; whence, the red particles having the greatest specific gravity separate from the fibrin, and fall to the bottom. In healthy blood, which coagulates quickly, they remain entangled. Whether or not, inflammatory blood (as it is called for the sake of distinction) contains more of fibrin than healthy blood possesses, is a question well deserving experimental inquiry. The firmness of the blood, and its contracted edges, when taken from a vessel by means of a lancet, under inflammatory disease, is still more instructive to the practitioner than the buffy coat simply. It indicates that the vessels are in strong action; and the blood sometimes appears as much drawn in, as if it were itself possessing a contractile living power.

The pulse was yet full, but abated in its action. The bleeding was repeated; the medicines were continued; and to the pills a grain and a half of crude opium was added. A large blister was applied to the parts affected with pain.

On the evening of the 24th he was considerably mended. Had procured much sleep in the preceding night, and had been free from pain through the day. The pulse 80, calm and soft. The limbs warm, and the whole skin of comfortable temperature. The blister had acted with sensible advantage. The blood appeared firmly coagulated, and was contracted; but less sizzly than before. The secretions had been of variable appearance, but for the most part as already described. The draught was continued, and the pills, with the intermission of a night, were repeated. Some broth was added to the diet. Two days were passed in great amendment; and the slight pain which returned was at night, at the same time that he was formerly accustomed to feel the accession of gouty pain in the limbs. At this period the hands and feet were almost free from inflammation, but were swollen, tender, and weak.

On the 27th I found that he had experienced a severe relapse, at the usual time of the night (about 1, A.M.), without any apparent cause. Said, that the bowels had been "drawn into balls," with excessive pain. The fæces continued foul, and the urine now, still more copiously than before, deposited pink and mucous sediment. He bore pressure on the abdomen much better than formerly, but it still produced some pain. The day was passed in ease. The pulse being yet full, the bleeding was repeated to xxxiv . The medicines were continued.

Now, as on the former occasions, he bore the bleeding well, and received sensible relief. The blood was still

firm in its crassamentum; but less so than before, and was free from the buffy coat.

On the 29th his amendment was very satisfactory. No pain had returned. Pulse calm and soft, at 76; skin cool; tongue becoming clean; urine lighter in colour, and with little sediment; the *fæces* of improved appearance, but still unhealthy; he bore abdominal pressure without complaint. The complexion yet retained some of its jaundiced hue.

A draught of *magnes. sulph. infus. rosæ et tinct. calumb.* was directed twice a day; and the compound calomel pill (five grains) every other night. Each night *Θij* unguent. hydrarg. fort. to be rubbed upon the right side. The diet to be gradually and cautiously made more nutritive.

From this date he became steadily convalescent; and the secretions by degrees returned to a healthy state. After the first week from its being adopted, the mercurial friction was employed only on alternate nights for about a fortnight, and then relinquished. The mercurial pill was taken for a fortnight, as above mentioned. Then increased attention was given to the weakened limbs; and a strict future regimen in every particular was enjoined. The bowels to be regulated by pills consisting of gum gambog. *pilul. hydrarg. pulv. alöes comp. et sapo. durus.* I shall subjoin a statement of the comparative properties of the urine at different periods.

March 27th.—It copiously deposited pink and mucous sediment. Its specific gravity (see Exp. xix. p. 122), 1.0207. Four ounces afforded of phosphoric acid 4.1 grains.

In the middle of April the urine was light, transparent, and free from all sediment, except the mucous cloud of health; its specific gravity 1.0087.

In February, 1816, the health appearing almost established, but the digestive functions still occasionally disturbed, I made the comparative examination of the urine (see Exp. xx). It was of an orange colour, with much mucons cloud, and an abundant deposition of uric crystals; of specific gravity 1.014. Four ounces afforded of phosphoric acid 2.1 grains.

In a review of this case, we observe a plethoric state of the system existing, to which a long course of gout had brought no effectual relief; a vitiated condition of the secretions become exasperated and confirmed; and then, under exposure to cold, visceral inflammation, chiefly affecting the liver, was produced. From the pain occasionally extending itself to the stomach and intestines, we may infer that the inflammation was chiefly peritoneal. The intervals of ease occurring almost regularly in the day; and the exacerbations at the usual time of night, when the pain of gout always most affected him; form an example of the modifying influence of the gouty diathesis over internal chronic inflammation—producing a case, which may, with apparent propriety, be denominated chronic retrocedent gout.

I did not see this patient in the interval of April 1815 and February 1816. At this last date his countenance had a clearer and more healthy appearance than before. He reported that he had for the most part enjoyed comfortable health during the above period; and had been free from gout.

I may add the following short case, which instructively represents the nature of a spasmodic affection of the stomach, in immediate connexion with obscure gout in the feet; and so alternating, that it seems to come under the head of chronic retrocedent gout.

CASE X.

A gentleman, aged 54; slight; with small vessels; of the nervous temperament; was first attacked with acquired gout at 29; has suffered many bilious attacks, and from his complexion bears the stamp of having an unhealthy state of liver. He is sensible of pain when pressure is applied to the right hypochondrium. Exposure to a cold wind, or the act of standing on a damp pavement, very quickly produces pain of the stomach. Has been subject to feel pains in the stomach in alternation with pain in the ankles and feet. For the last fortnight he has been much indisposed in this manner. He goes to bed at nine, and falls into a comfortable sleep. About twelve or one, has for the last fortnight very regularly awoke with violent pain in the epigastric region, shooting up the sternum, accompanied by spasms, urgent flatulence and heartburn. No difficulty of breathing. The pain relieved by pressure. It remains four hours; and, being then at its acme, gradually declines; and towards morning, ease and sleep return. One night he discharged much bilious matter from his stomach. Lately has had slight gnawing pain in the ankle and tarsal bones of one foot, and two nights ago, "much twinging pain in the ball of the great toe." Yesterday he took an active purgative, and to-day makes the following report:

"I went to bed at my usual time, and awoke about twelve with some pain in the right ankle, and a sense of heat, throbbing and stiffness, about the ball of the great toe, but without any affection of the stomach, except a little heartburn. The symptoms continued during two hours. I then fell asleep, and rested well. This morning I do not find either pain, discolouration, or weakness in the parts so recently affected."

The treatment of this case evidently consists in the employment of means to restore the health of the digestive organs, by suitable medicine and regimen.

Of cases of apoplexy occurring in the paroxysm, and successfully treated, I shall state concisely two examples :

CASE I.

J. M. aged 60, robust, of sanguineous temperament, corpulent, and very plethoric; has suffered severely, at short intervals, for ten years past. Had been under the paroxysm about a week, one foot being much inflamed and painful, and the other about to be affected. He had kept the parts wrapped in flannel, and had not restrained himself in diet; from an idea that the gout should be *encouraged* in the feet. He had also allowed the bowels to be costive.

In the morning, while in the act of stooping to adjust his dress, he fell upon the floor in a fit of apoplexy of alarming violence. Instant relief was at hand. Twenty ounces of blood were taken from the arm, by a large orifice, and active purgative treatment was adopted and continued. Every bad symptom was quickly removed, and no paralysis followed. The gout continued in a mild form only. The ordinary treatment of the constitution, as in apoplexy unconnected with gout, was steadily continued with complete success.

CASE II.

F. L. aged 63, of very similar make and constitution to the foregoing. Gouty during the last twenty years.

Had been gradually increasing in corpulence; and used of late only passive exercise. The bowels were costive; and the secretion of urine, though irregular, was for the most part very deficient. The gout was severely affecting one foot. It had been nursed in flannel. As in the former case, the diet was indulgent. Soon after dinner he was seized with apoplexy. Blood was freely taken from the arm; a stimulant injection was administered; a large dose of calomel (10 grains) was given as soon as it could be forced down, followed by sulphate of magnesia with infusion of senna, &c. in repeated doses. In four hours, although the bowels had been freely emptied, and depletion had been abundant, the pulse was strong and throbbing, and the head was painful and confused. Twelve ounces of blood were then taken from the jugular vein; a cooling lotion was freely and constantly applied to the head; the purgative treatment was pursued; and the diet was restrained so as to afford but little nourishment. Gout confined itself to the foot, and continued with moderate symptoms, for a short time. The patient recovered without any unfavourable consequence. In these cases, the disease of the brain could not be considered an act of gouty retrocession; for the inflammation in the extremities was not suspended at the moment of attack, and became abated only from the active depletion which was adopted.

It should rather be stated, that the mismanagement of the patient had induced a plethoric state of vessels, too predominant for an effort of Nature to afford relief by the gouty action; and aggravated also in the determination of blood to the brain, by error of diet; by the influence of heat in clothing; and by neglect of the alimentary canal.

I have met with instances of gouty persons, between

the fiftieth and sixty-fifth year more particularly, who have ceased to have their usual attacks of gout, and have been irregularly affected with nervous symptoms, in so remarkable a manner that they have entertained the idea that gonty matter has been wandering over the frame, without fixing upon the individual part on which it should exclusively exert its action. In some such examples the head is affected with pain and confusion; the eyes with occasional sense of mist or other disturbance of vision; the ears with unnatural sensations of noise, as the hissing of a tea kettle, or the sound of a bell, or a slighter but more continued ringing. When the head rests on the pillow a distressing pulsation is felt. In more urgent cases, this beating affects the head also during the day. I have known the patient to express the curious idea of the hair starting up to a point, and causing a feeling of soreness at the part. Upon any agitation of mind, or over-exertion of the bodily strength, the heart will partake in this nervous disturbance, and be affected with palpitation.

I saw a gentleman labouring under most of these symptoms in a very urgent degree. He was aged 60, and had been subject to gout since 25, till about five years ago; when, for the relief of symptoms affecting the brain, which were supposed to threaten apoplexy, he was bled, within a short time, to the extent of 150 ounces. This was certainly an injudicious use of a remedy, only proper within certain limits. No gout has ever taken place since this large depletion, but he has suffered extremely in his general feelings. When I saw this patient, affected as I have described, his lower extremities being also remarkably cold, I prescribed alterative and aperient medicines, a pediluvium of hot water containing mustard and salt to be used daily, and a stimulating embrocation

to be diligently rubbed in afterwards; a supporting diet; and the avoiding of all anxious and fatiguing exertion. By these means he was speedily relieved; and by a removal into the country, and still following up the use of medicine and careful regimen, he was happily restored.

A patient of slight form and nervous temperament, reports to me the following remarkable form of complaint which he experienced.

After exposure to wet and cold, he suffered from symptoms of inflammation of the lungs. He was bled largely in the morning, with immediate relief to the chest. In the afternoon gout seized the knee. At night the pain suddenly quitted the knee, and his head became affected with extreme confusion, followed by delirium, and entire loss of vision. These alarming symptoms continued many days, and were attended with an anasarcaous swelling of the limbs. The head was but little affected with pain. The active use of purgative medicine removed the symptoms. On first recovering his sight, he found, on attempting to read, that all the letters were reversed.

It occurs to some gouty persons of the nervous temperament, and not in particular to those in years, to be affected with sudden colic upon accidental occasions of exposure to wet and cold, or of disagreement from acid fruit, or some article of diet. It almost invariably happens, that, in these attacks, the action is spasmodic and not inflammatory; and it is a frequent result that some strong stimulant, as hot brandy and water, or compound spirit of ammonia in hot water, gives relief. But an evacuant should be used afterwards. If the stomach itself be disordered, as would be shewn by much sickness being present, an emetic of ipecacuanha should be first administered; and next a dose of calomel with an aromatic purgative; or, if the emetic be not required, the latter remedy may be

used alone. The bowels can scarcely be acted upon with too much freedom. I will not omit to mention, that, in conjunction with the purgative treatment in these cases, if the symptoms should proceed, anodyne medicine is materially useful for the relief of pain, irritation, and restlessness. I have found very satisfactory effects from the pill described at p. 224. It is a safe, and often an important practice, to abstract some blood from the tender part of the abdomen by cupping. Although the attack be not in its original nature inflammatory, yet the continuance of pain causes more or less of congestion of blood in the affected part. The application of a large blister as a subsequent remedy to the cupping will sometimes be demanded. The intensity of pain and spasm has yielded to the immediate remedies employed at the early moment of alarm; but it may happen that the symptoms assume a continued form, although so much subdued in violence. We must therefore keep a watchful eye upon the case.

The patient very naturally conceives that his colic is a certain attack of gout in the stomach; or that gout is affecting the head, if that be the disordered part. He prays for the return of the gout in the extremities; and feelingly declares, that he would gladly endure any torture in the foot, to be relieved either from the distressing sensations which affect the internal parts in a chronic form; or, the alarming acute attack of which I have been treating. Dr. Cullen has distinguished these anomalous forms of complaint by the term misplaced gout; but I would rather define them to be disordered functions of internal organs in a gouty constitution, and modified therefore in their character. In the case which I have described at p. 460, the highly stimulant applications used to the feet, although they acted strongly as rubefacients, did not give

rise to any degree of gouty action. Also in the colic of the gouty patient, when I have used a similar treatment with a view to favour the production of gout, I have not succeeded in this object. But I find that when gout has been retrocedent from the extremities, and such treatment has been employed, it has almost with certainty returned to the extremities. It does so even spontaneously on many occasions, as soon as the internal disease subsides, although no stimulating applications have been used to the external part. I consider therefore that the results in question are to be received as strong presumptive proofs of the nature of the internal disease.

One gentleman informs me that at a time when he was entirely free from every sensation of gout, he walked about on a wet day, and upon his return home was seized in a moment with a pain at the pit of the stomach, so excruciating, that he could suppose the stomach itself to have been tied up in a knot with a strong cord. He instantly drank brandy, and in his alarm took it so freely that he became intoxicated. Yet this proved the cure.

Another gentleman, when free from gout, produced a similar attack from incautiously eating some Orleans plums. He described his pain as if some one were drawing a sword across the stomach.—Such are the intense sensations of gouty persons. He was restored by the use of cordials and aperients.

I advise patients of this description in the first place to pursue an exact regimen; secondly, to avoid wet and cold with extreme circumspection; and, lastly, to provide themselves with an anti-spasmodic medicine as a *vadē mecum*. This may consist of a combination of ammon. carbon. conf. aromat. mist. camphor, sulph. æther, tinctura opii. A full dose of calomel and a purgative, made moderately aromatic, should follow the use of the anti-spasmodic.

The application of heat to the abdomen is an important auxiliary to the other treatment, and may be made conveniently by means of a large bladder filled with hot water. I am now describing the action which belongs to spasm, and not to inflammation. The pediluvium with mustard and salt should be added to the list of general remedies; but I must again observe, that the radical treatment consists in the free use of purgative medicine; the soothing use of opium being conjoined, as I have already pointed out; and the remedies used according to the continuance of the case, and as the judgment of the practitioner will direct.

IN bringing the present treatise to a conclusion, I deem it necessary, *briefly*, to make a more particular reference than I have yet done, to the principal constitutional diseases, to which, gout and the habits of the gouty may be considered as pre-disposing.

A few persons, as I have before stated, pass through a long life, suffering periodical visitations of the gout, and never incurring any other disease; until at length the natural infirmities of years, aggravated by the effects of gout, emaciate and wear out the frame.

I knew well a gentleman, advanced to a great age, who, during forty years of his life, was not on any one occasion absent from his office of business, which was situated five miles distant from his residence and daily required his attention, upon any other account than the gout, to which he was subject at regular periods.

Another gentleman, now aged 86, states that he has not been subject to any other disorder than the gout.

These examples of the gouty life protracted to a great age are comparatively rare; and may be said to appear only in the chosen few, who, in strength of constitution, were formed for remarkable longevity.

Those *who neglect the proper management of gout*, are more liable to die prematurely*; and others, who reach a considerable age, having left the disease to its own course, are afflicted with perpetual discomfort and frequent misery.

The occasional occurrence of *apoplexy*† among gouty patients, is a familiar fact; and to which either immediate death, or paralysis, for the most part succeeds. It is also a truth which deserves to be stated, that some gouty persons, advancing in years, lose the disposition to their former attacks of gout, apparently in the same proportion that they acquire a tendency to apoplexy. The individuals whom I have seen to exemplify this statement, have possessed more of the sanguineous than the mixed temperament, with great plethora of vessels; and have been robust and corpulent. Their habits growing more and more sedentary, *and their indulgences in living not being diminished*, the liver becomes obstructed; the action of the bowels and of the kidneys is not adequate to the excess of ingesta, and of the chyle that is assimilated; whence, a redundancy of blood in the system, and an interruption of the healthy balance of circulation, follow as the conse-

* Dr. Sutton, in commenting upon the prevalent notion that gout rather tends to lengthen than abridge the duration of life, observes, that those, whose pecuniary interests are concerned in this particular, make a different estimate, and by which, "those subject to the gout are placed under the predicament of paying a greater premium for the insurance of life, than those who have never suffered from the disease."

† Dr. Heberden forcibly remarks, "It can hardly be reckoned one of the disadvantages of the gout, that after destroying all the comforts of living, by this weight of misery, or by bringing on a palsy or apoplexy, it immaturesly extinguishes the powers of life."

quence. It appears to me as a further reason why the gout does not occur as formerly, that these elderly persons are seldom exposed to the most active of all the exciting causes, wet and cold. In the temperament and structure just described, the apoplexy which occurs, is, for the most part, and especially unless prompt assistance be rendered, immediately fatal in its event. In the mixed temperament with less fulness of habit, the attack is probably more commonly followed by paralysis; or, sometimes, by the agreeable consequences of perfectly successful treatment.

Obstruction of the liver; jaundice; cholera morbus; dyspepsia; hæmorrhoids; constitute the chief assemblage of the morbid states of the digestive organs, to which gouty persons have the strongest tendency.

Asthma occasionally supervenes on gout; and the invalid has only short intervals from one or the other disease; a respite which he owes principally to the less variable weather of the summer season. It sometimes happens that an asthma, which has chiefly been depending on a faulty state of the chylopoetic viscera, becomes suspended for a long time; or, in more favourable cases, is even removed by the occurrence of gout taking place as a new disease in the constitution.

Hydrothorax is an occasional result of that complicated state of visceral disease to which the gouty constitution predisposes. A few, whose lungs are unhealthy, and whose excesses in living are great, die before middle life with all the symptoms of consumption.

Ascites, arising out of the errors of diseased liver or spleen, or of both viscera in conjunction, sometimes happens, as a remote sequel to gout.

Gravel occasionally takes place in gouty persons, but for the most part it is the disease of their early life before the invasion of gout; and *stone in the bladder*, although (as in Sydenham) it now and then unites its tortures with the gout, is, according to my observation, as I have stated at p. 43, of rare occurrence.

Erysipelas attacks some gouty persons; and sometimes seems to represent, or come in the stead of, the expected fit. It has appeared to me that gouty women are more disposed than gouty men to this additional disease.

Erythema and *Urticaria* may be mentioned among the incidental complaints of gouty persons. I have seen an instance of each of these complaints, shortly preceding the paroxysm; and, doubtless, arising from a faulty state of the stomach and bowels.

The occasional union of *Rheumatism* with gout, has already engaged our attention.

Cramps, although a symptom only of a morbid condition of the nervous system, and usually depending on an unhealthy state of the chylopoetic viscera, is with some gouty persons so urgent a suffering, as to form the leading feature of disease which most engages their attention.

The most important of the diseases, which I have here enumerated, must receive their pathological explanation from the two-fold cause of an over-loaded circulation with partial debility of vessels. Accordingly, as one viscus is more weak than another, partly from original structure, and partly from the frequent interruption of its healthy functions, so will become the consequent seat and nature

of the supervening disease. Whether it happen, that some one particular organ acquires, from obstruction and weakness, a state of congestion which renders it incapable of performing its former share in the general circulation; or whether, by the peculiar action of stimuli, an accidental increased action of vessels take place in a particular part; the healthy balance of circulation is interrupted, and, from either cause, similar effects may be produced. Thus, apoplexy, in a plethoric person, may arise out of the effects of a continued obstruction of the circulation of the liver; or it may be produced in a more sudden manner by the excessive stimulus of intoxicating liquors acting on the brain. When the picture which is here drawn (and it is one that is rather softened, than too closely copied from life), added to that of the sufferings which gout directly inflicts, is well considered, it seems surprising that the idea which many gouty persons fondly entertain, that a paroxysm is an indication of health and strength of constitution, rather than an actual disease, should ever have been cherished.

Let them, ere it be too late, change their destructive habits of indulgence; and instead of reposing a mistaken confidence in the palliative powers of the *eau medicinale*, and other nostrums; or even trusting to the supposed curative influence of a fit of gout; let them adopt, with virtuous resolution, a true *prophylactic regimen*.

In connexion with the present general view of my subject, I shall offer a slight discussion on the practical principles which are applicable to the *acute* diseases under which gouty persons, in common with others, may labour; and lastly, the treatment of apoplexy incidentally occurring to the gouty, when wholly free from gout, is to be considered.

The prejudice which has for so many years existed against the propriety and even safety of general bleeding in gouty cases, whatever the incidental disease might be, although for the last very few years it has certainly been on the decline, appears to me to be yet in considerable force; operating on the public at large, as well as influencing in some degree the medical profession.

Dr. Cullen, however, who does not make the least mention of blood-letting as a remedy in any of the forms of retrocedent gout, which he has stated, allows this practice in the phlegmasiæ, which, occurring in a gouty person, he calls misplaced gout; and thus expresses himself: "In this case, the disease is to be treated by blood-letting, and by such other remedies as would be proper in an idiopathic inflammation of the same parts *."

Dr. Heberden † speaks of general bleeding with caution and some distrust, in the accidental diseases of a gouty person, lest the gout should be at hand; but he grants this exception, "that it will be far more hazardous to neglect bleeding in an inflammatory distemper, than to take away blood in the gout."

In the fourth volume of the Medical Transactions of the College of Physicians, Dr. Haygarth states an interesting case, which he considered to be *carditis arthritica*, and relates that in consultation he proposed this question: "Is it necessary and safe to take some blood from a vein, even though the gout be the cause of this inflammation? If there were no such suspicion, the violence of the disorder would require large and repeated venesections."

I offer this quotation, as a strong proof of the sentiment which prevails against the use of the lancet in the inflammatory diseases of the gouty.

* Par. 583.

† Commentaries, p. 45.

The recommendation expressed by Dr. Cullen, and just now quoted, may be held as the true axiom of practice; and the facts indeed in support of it appear to my observation so familiar and evident, that many details upon the subject need not be added.

I visited an elderly gouty gentleman suffering from an inflammation of the kidneys. Bleeding from the arm to a free extent was one of the means of treatment, and was attended with the best success.

A gouty man, of plethoric habit and sanguineous temperament, was seized with inflammation of the lungs, which had been induced by exposure to wet and cold, when on horseback. Repeated venesection, in concurrence with the usual general treatment, was employed with the same favourable results as would happen in a similar disease of one not gouty.

A gentleman, more than seventy years of age, who had been afflicted with gout through the greater part of his life, was labouring under severe cough, much pain of the head, excess of vascular action, and evident plethora. No symptoms of gout were even threatening. He had an insurmountable objection to the remedy of bleeding. In a short time, hæmorrhage from the nose took place, which was almost alarmingly profuse; but the system became effectually relieved, though with too much expense of the powers of the constitution.

In this case, a definite quantity of blood abstracted at a timely period by art, and in co-operation with the use of medicine, might have proved equally successful in its effects, and would have been much less debilitating than the spontaneous hæmorrhagic action of the vessels. Na-

ture seldom or never errs in her intentions ; but the measure of her performance does not seem always just. So in spontaneous hæmorrhage. The process in its beginning may have been necessary, and proves salutary ; but its uncontrolled continuance might be destructive.

When I considered the subject of general bleeding under the head of remedies in the gouty paroxysm, I stated, that if the inflammatory diathesis were strong and permanent (a full strong pulse with a continued hot skin), and no contra-indication presenting, the remedy of venesection should not be delayed ; but otherwise, a sufficient dependance might be placed on the derivation produced from the circulation by purgatives and diuretics.

When a local inflammation of any kind is set up in an external part, the use of general bleeding is more optional, than where an inflammatory action of the heart and arteries prevails alone ; or, than in the case of an internal inflammation ; because we know that the determination of blood to the inflamed external part, is to a certain extent a relief to the general circulation, and does not endanger the destruction of the whole machine ; but if the inflammatory diathesis of the system be violent ; or inflammation exist in an internal organ ; venesection should be practised without restraint from those fears which belong only to false pathology. Whether the patient be gouty or not, the same principles of treatment must be exercised ; or life itself may fall the sacrifice.

The importance of accommodating our practice to existing circumstances, and to real indications, is still more manifest in the case of a constitutional tendency to apoplexy occurring in the gouty, unconnected with the paroxysm. This subject, then, is now to be discussed.

A gouty person should take early notice of sudden

increase of corpulency, whether it be general, or more partially confined to the abdomen; and, in proportion as his structure favours plethora and congestion in the vessels of the brain, should this attention be paid. Also a consequent regulation of regimen, both as to diet and exercise, and the hours of rest, should be adopted. A daily action of the bowels is a point of attention of the greatest magnitude; and a due secretion of the kidneys is also never to be overlooked. I shall not repeat the arguments which I have already advanced in this volume, upon the high importance of this function to the health of the system.

Among other symptoms of serious warning, may be mentioned, pain of the head with a sense of fulness, confused vision and giddiness; excessive drowsiness in the day, and dulness of the faculties with severe hypochondriasm; and in the night, profound sleep with deep and slow inspirations in the breathing, frightful dreams, and night-mare. The dyspepsia, which is attendant upon these symptoms, should be treated by purgatives and correctives; and every species of tonic should be delayed till a correct balance of the circulation, and a healthy condition of the several secretions are restored.

In proportion as the signs of congestion in the vessels of the head are manifested under the general circumstances which have been stated, the propriety of cupping on the neck will be pointed out; or, if an inflammatory diathesis of the system be also present, general bleeding will deserve the preference; but in cases of this description, *of the fullest amount of symptoms*, undoubtedly both local and general abstraction of blood will be required.

I shall relate a few examples, which have come immediately under my own observation, illustrative of these different positions:

A gentleman, aged 45, of plethoric habit and sanguineo-nervous temperament, who had suffered repeatedly from gout during seven years; at the accustomed season of the year, and at the usual time of the invasion of the fit, was awoke about two in the morning with dreadful night-mare, and all the horrors of suffocation; and these symptoms recurred two or three times. He was cured by the free employment of purgatives and diuretics; by restraint of diet, and by general regimen: and no attack of gout occurred.

A tall robust man, aged 64, of full habit and sanguineous temperament, who had been gouty since the age of 30, but for a year past had been free from attack, had become very plethoric from indulgence. After dinner, when sitting in his chair, he suddenly felt an extraordinary sense of coldness and numbness, from the head downward on one side, and had not the power to rise. He was immediately cupped to a free extent; and the bowels being also sufficiently acted upon, every unpleasant symptom vanished in a few hours.

A gouty man, aged 71, robust, corpulent, and of the mixed temperament, free from gout for three or four years past, was lately seized, under great indulgences in living, with a slight apoplectic fit, which was followed by a considerable degree of hemiplegia. The faculties of the mind were much weakened. In this case, there was an evident obstruction of the liver, and the several secretions were extremely vitiated. The bowels were obstinately torpid. So much inflammatory diathesis was present, that general bleeding was at first employed. The excitement and correction of the functions of the liver, alimentary canal, and kidneys, constituted the regular treatment; but in addition to these means, although the diet

was abstemious, occasional cupping was required. The patient recovered, and no paralysis remained.

A corpulent man, aged 53, plethoric, and rather of the sanguineous than the mixed temperament, had been subject to gout several years, and had suffered a severe paroxysm about two months since. It had been left entirely to its own course, no treatment having been adopted. He had never intermitted his full habits of living. After having complained, during the day, of much pain in the head; and a very slight distortion appearing in the muscles of the face (yet, not seeking medical advice); as he was stooping to undress himself in preparing for bed, he fell down in a fit, and expired.

The last case, to which I shall refer, was a seizure of apoplexy in a gentleman between sixty and seventy years of age. For many years he had been a severe martyr to gout; but latterly it occurred only in the chronic form. Under the united influence of a diet too full with relation to other circumstances, of a torpid state of bowels, and of habits entirely sedentary, this attack of apoplexy suddenly took place, and with a violence which threatened to be instantly fatal. Blood was copiously taken from a large orifice; and the usual general treatment was pursued with perfect success.

In these cases, the prognostic of permanent recovery is favourable, in proportion as paralytic symptoms disappear; as tranquillity and comfortable feelings return; as the sleep is less profound and more refreshing; as the faculties become clear; and the pulse recovers its regularity. So long as the pulse remains variable and intermitting, we may be assured that the functions of the brain are yet unsettled. Our judgment is further governed by the state of the secretions, and by the obedient action of the bowels. If assistance be fortunately at hand, and blood

be freely drawn, while no rupture of vessel has yet taken place, success is very probable. The half-erect posture, while in bed, instead of one that is quite recumbent, should be observed; and the free application of an evaporating lotion to the head forms a valuable auxiliary to the general means of treatment. A cool state of the apartment is an additional point of attention of much importance.

A prophylactic regimen for the remainder of life, and a strict regulation of the bowels, must be insisted upon as indispensable to justify the expectation of future safety. Tight clothing should be avoided, and especially a tight neckcloth. Such persons as are corpulent and plethoric, cannot too much observe a rule to avoid stooping; and they should be careful not to turn the head suddenly, without turning the body at the same time. The washing of the head every morning with cold water, by means of a coarse towel, is a practice of very great utility; and it may be repeated at any time, with equal advantage, when the head is painful, in these cases of local congestion of the vessels. Hot rooms must be shunned, as being highly injurious; and calmness of spirits, and serenity of temper, are comprised amongst the most important sources of security. By the united care of mind and body, according to the rules of the physician, any threatening of apoplexy may be turned into a lesson of safety: but those who relapse into error, must experience those melancholy consequences, which are too well known to require a further description.

A TABLE of the *Analytical* Method of investigating a Case of Gout, and the general History of the Disease, adopted by the Author in forming the present Treatise.

1. The age; sex, &c.?	2. General structure; bulk; skin; complexion; family resemblance?	3. Habit; temperament; general points of constitution; idiosyncrasy?	4. Place of abode; climate; to what other diseases subject?	5. Station of life; employment?	6. Habits of living?	7. Gout in family, and in what degree of relationship?	8. At what age first attacked, and in what part?
9. In what parts subsequently, and in what order? Different parts together, or in succession?	10. At what part of the twenty-four hours, in general?	11. Time of year, and periodical?	12. General causes, predisposing and exciting?	13. What the premonitory symptoms?	14. What relation has the violence of the attack to the particular predisposing and exciting cause?	15. Which the most painful part?	16. Most pain day or night, and at what periods?
17. Local sensations in the height of the paroxysm.	18. Local appearances, and characters; temperature of the inflamed part contrasted with other parts. Description as the inflammation goes off?	19. General symptoms, as to pulse, skin, tongue, action and state of bowels, kidneys, kind of secretions, &c.	20. State of the mind? what nervous symptoms?	21. Irritable urethra? and cramps; and in what muscles?	22. What the usual treatment; and with what results?	23. Gout ever retrocedent; to and from what part; and what the exciting cause?	24. What the longest and shortest duration of a paroxysm?
25. Longest and shortest interval between the paroxysms?	26. What change of organization produced in the parts affected in the paroxysms?	27. Does the gout increase or lessen in progress, as to the severity or duration of the paroxysm?	28. What tendency has a violent paroxysm been found to have in prolonging the subsequent interval?	29. On what disease has gout supervened? and the contrary?	30. Has the patient considered his constitution benefited or injured by gout; and how does this fact appear?	31. To what other diseases has gout been disposed?	32. Particular practical opinions of various authors noticed and examined.

APPENDIX,

Comprising a Series of Experiments made upon the Dog with several Gout Nostrums of the present day, and the different Preparations of Colchicum; with a View to demonstrate the Agency of these Substances on the Animal Œconomy; and to furnish some useful Conclusions, in Reference to the Employment of Colchicum as a Medicine in Gout.*



SOME of my Readers will probably condemn these Experiments upon Animals, as not creditable to humanity; but I beg it may be considered, that this investigation was undertaken solely for the purpose of obtaining information on some important points relative to the health of Man.

It may probably be asked, are these experiments really deserving confidence when their professed object is to compare the effects of medicines upon man and the dog, which are in their œconomy so differently constituted?

* I must not omit to acknowledge my obligations for the very material assistance afforded me in these Experiments, by Mr. Probart of Great Portland Street; and his brother, Mr. Frederick Probart, a zealous medical student.

I answer, that in the structure and functions of the stomach and alimentary canal, there is a sufficient general similarity for the purpose of these experiments. I offer them only in illustration of the results which I have found on the human subject; and as confirming, but not instituting, my rules of practice. They certainly authorise some positive conclusions as to the properties and strength of the different preparations of which I have treated, being fairly comparative; they set forth their mode of action upon the animal system; and shew the nature of the symptoms by which death is produced, when the medicines are administered in the fatal dose.

EXPERIMENT 1.—*Eau Medicinale*.—One hundred and sixty drops, measuring 135 minims, of eau medicinale, were injected into the jugular vein of a full-sized very strong dog, at a quarter before one *p. m.*

He immediately ran about, apparently not affected. In ten minutes the pulse* was 100, and irregular; in a quarter of an hour he sat on his hind legs, licking his wound and looking sleepy and slightly languid; he soon after walked steadily. In three-quarters of an hour he ran about without particular inconvenience, still however

* In the early part of these experiments, I did not make a previous examination of the pulse, considering that its average frequency in dogs was 140 in a minute. Afterwards, however, directing my attention to this point, I discovered a great variation both as to the frequency and regularity of the pulse of the dog in his healthy state. This variation in different dogs was from 120 to 180; and if the animal became at all alarmed, the pulse would not only become very frequent, but in several instances intermittent. As far as I have been able to ascertain, the average number of inspirations in the healthy dog may be stated at 36 in the minute.

having the sleepy look. The pulse intermitted after every five beats, and was 90 in the minute.

In an hour and three-quarters, the dog was evidently much disordered; had no disposition to move; and the drowsiness was increasing. He had vomited twice, bringing up a glutinous matter of a greenish yellow colour, and had two dark alvine discharges. In four hours, a discharge of yellow-coloured viscid mucus flowed from the mouth. He exhibited great lassitude; the eyes were dull; and, when left alone, he seemed heavy and lethargic; the pulse was 94; and the inspirations were natural.

In five hours and a half, he was sitting up, but looking exceedingly dejected; his pulse was 69, and irregular; the inspirations were 24.

In six hours and a half, the inspirations were not apparently difficult, and 20 in the minute; his eyes closed; he appeared sound asleep. In seven hours and a half, he had changed his situation, and was lying with the head erect, endeavouring to swallow some drink placed near him. In a few minutes he vomited a clear liquid, stood erect, appeared relieved, and drank freely of some water: the pulse was 90, and intermitted at every three or four beats.

In eleven hours, the inspirations were only 18; and the dog was manifestly worse in all respects.

The following morning, at eight o'clock, he was found with his head reclining on the ground. On moving and trying to place him upon his legs, he was wholly unable to support himself, and instantly sunk down on the straw. The inspirations were quick, irregular, and difficult.

Nine o'clock.—The breathing was harder; and every 15 or 20 seconds the whole body was violently convulsed.

Half-past nine.—Continually convulsed, and still more violently; and he was much distressed.

At ten o'clock the convulsions had ceased; and the animal lay extended, and moaning.

Eleven o'clock.—Life appeared to be almost extinct: a large quantity of thick, bloody mucus was observed near him; his inspirations were 14 in a minute; a quantity of saliva flowed from his mouth. In a quarter of an hour, his inspirations were but 12. In another quarter of an hour they were only 10. He was now and then convulsed.

Three o'clock.—Throughout the morning he was in a state of complete insensibility; his eyes were now fixed, and the eyelids shut; the teeth were firmly closed, and power of voluntary motion was lost. The inspirations were increased to 26 in the minute.

At four o'clock he had expired.

Appearances on Dissection.—On the following morning the body was opened. The stomach exhibited slight appearances of incipient gangrene; the duodenum and jejunum were highly inflamed; the ilium, colon, and cæcum, less; and a small spot of ulceration was seen at the commencement of the rectum. The internal coat of this and of the other intestines, through their whole course, was here and there marked with deep rose-coloured spots, about the size of a pea. The auricles and ventricles of the heart, and the vena cava, were much distended with black blood, partially coagulated; the gall-bladder was filled with bile.

EXP. 2.—*Wilson's Tincture.*—One hundred and sixty drops of this medicine were injected into the jugular vein of a mongrel dog, the size of a large terrier. He became languid in a few minutes, and a slight convulsion

of the muscles of the thighs was observed. In half an hour he voided some urine. In little more than an hour, the symptoms were more favourable. In an hour and a half, he seemed so much recovered as to be capable of running many yards, and up a pair of stairs. In three hours, he was discovered to be in a sound sleep. Some greenish yellow mucus appeared around the mouth. In five hours, he ran about from place to place as if restless, and appeared debilitated. In seven hours and a quarter, we were much surprised to find that he had expired. This dog lost six ounces of blood in the operation. There was a small sediment in the bottle of the tincture, which was shaken up, and injected with the fluid.

Dissection.--The œsophagus appeared in a natural state. The whole surface of the stomach exhibited marks of inflammation, and the mucous membrane, towards the pylorus, was almost gangrenous; the stomach contained bile and mucus; the duodenum, near the stomach, was much inflamed; the other portion but slightly. The jejunum, at its commencement, was highly inflamed, and covered with an abundance of coagulated albumen. The inflammation extended through the whole course of the jejunum and ilium, but gradually diminished. In the colon the appearances of inflammation increased; the cæcum was slightly inflamed; and the rugæ of the rectum exhibited numerous red points. The lungs bore slight appearances of inflammation; the liver was partially inflamed; the urinary bladder was much contracted; the brain was in a natural state.

EXP. 3.—*Reynolds' Specific.*—At twenty minutes past one *p. m.* a hundred and eighty drops of this tincture, measuring two drachms, were injected into the jugular vein of a young mongrel dog. He passed at the moment a natural alvine discharge. At three o'clock he had

vomited a large quantity of mucus, tinged with bile; the inspirations were hurried and performed with difficulty, and were 62 in the minute; he hung down his head, and there was universal tremor.

At four o'clock he was making ineffectual attempts to vomit; looked miserably dejected, and stood with difficulty.

Six o'clock.—He was sitting up, but much enfeebled; the inspirations were 28 in the minute.

Seven o'clock.—It did not appear that he was suffering any pain; he was lethargic, and indifferent at having his head raised by the ear, or being in any way disturbed: his inspirations were now apparently free from distress, and 36 in the minute.

At nine o'clock he was found dead; a white jelly-like substance appeared to have passed from the rectum.

Dissection.—The stomach was highly inflamed, and completely distended with grumous blood; the inflammation had extended with great violence through the intestines, as far as the rectum, which was slightly inflamed and abundantly marked with deep rose-coloured spots of the size of a pea. An ounce of colourless limpid fluid was found in the chest; the gall-bladder was much distended; the blood in the auricles, ventricles, and veins, was black and coagulated.

EXP. 4.—*Hellebore and Laudanum.*—One hundred and sixty drops of this mixture, in the proportions recommended by Mr. Moore*, were injected into the jugular vein of a large dog.

He was instantly motionless and insensible, and to all appearance dead. In six minutes respiration was perceptible, but extremely quick and difficult; he attempted

* See p. 196.

ineffectually to vomit. In ten minutes the pulse was 90. In half an hour, he passed an involuntary loose discharge of mucus, tinged with blood, and some blood was observed to come from the mouth; the breathing was convulsive and extremely difficult. In three-quarters of an hour his inspirations were only six in a minute, and very deep; the whole body was drawn up, and exhibited emprostotonos. In two hours the symptoms were abated; the respiration was much easier, and there was no longer any convulsion. In two hours and a half, he was still better and capable of sitting up. In four hours and a half, there was a further improvement in his appearance; and he walked many yards with facility. In six hours, he was found upon his legs, much recovered; his eyes were bright, and no other symptom appeared than debility. In eight hours, the ligature had given way, and the wound was bleeding freely. On the following morning he was running about, and had a good appetite. There had been several alvine discharges during the night and throughout this day. The next and third day, he was quite well.

EXP. 5.—*Hellebore and Laudanum*.—The last experiment was repeated; and one hundred and sixty drops were injected into the jugular vein of a good-sized strong female dog. The operation was instantly followed by emprostotonos, and a spontaneous liquid alvine discharge; the limbs were convulsed; the pulse was 116, and irregular.

In six minutes the convulsions were general; the animal made ineffectual attempts to vomit; the inspirations were laboured and very deep, and only eight in the minute.

In sixteen minutes they were so rapid, that they could not be counted, with occasional intermissions. In eighteen

minutes she was found moaning. The inspirations, in three minutes after, were only 3; the pulse was 100 in the minute. In an hour, she passed another very copious discharge; the pulse was regular; there was abundant salivation, and she appeared much better. She occasionally uttered cries; now rose upon her legs, staggered a few paces, and fell. In two hours, the tongue was protruded, and saliva flowed from her mouth; the body was slightly convulsed, and the respiration was laborious. In half an hour, the convulsions became frequent; the respiration as before; there was occasional whining. In three hours and a half she was sensible, and appearing to recover; but, as yet, incapable of supporting herself. The inspirations were irregular and difficult; the tongue still protruded; the eyes were open and bright; but she started and seemed terrified at the light of a lamp. In four hours and a half, she was sitting on her haunches, but was incapable of keeping this posture for any length of time. She was breathing slow but freely; she no longer shrunk from the light, and appeared to be recovering. In five hours and a half, her strength was returning, and she was in every respect better. In six hours and a half, her appearance exhibited complete paralysis of the hind extremities. She seemed quite sensible, and looked very animated. On the following morning she appeared well, and ran about without any inconvenience whatever.

Dissection.—In the course of the day, this animal was destroyed for the purpose of ascertaining what internal lesion of structure had resulted from the experiment.

The whole viscera were found healthy except the lungs, which here and there presented livid spots. The auricles of the heart, and the vena cava thoracica, were filled with very black and firmly coagulated blood.

Exp. 6.—*Elaterium and Laudanum.*—Two grains of

elaterium and sixty drops of landanum (forty minims*) with about a drachm of water, were injected into the jugular vein of a middle-sized terrier.

He instantly lost all power of motion, and was quite insensible and lethargic. In twenty minutes he had a natural alvine discharge; he was much convulsed; the pulse could not be counted. In rather more than an hour, he was sensible to pain on being pinched; was still much convulsed, and continued lethargic when left to himself. In three hours, there was no farther remarkable change. In five hours and a half, there was excessive languor, and he was still insensible. In seven hours, he seemed to suffer great pain, moaning continually, but insensible to any object near him; his eyes were closed and fixed, and the teeth were closed; there was a convulsive motion of the hind leg; the inspirations were sixty in a minute. In little more than eight hours he was visited again, but there did not appear any remarkable change. He died during the night.

Dissection.—The left lung presented the appearance of inflammation; the right was also inflamed, but in a much less degree; there was a very partial inflammation of the left lobe of the liver. The whole surface of the stomach was slightly inflamed; the inflammation extending with increased violence through the duodenum and jejunum, and into the ilium, which was much contracted; and here also the inflammation was still more strongly marked. The brain had a healthy appearance.

EXP. 7.—*Sir Everard Home's Vinous Infusion of Colchicum.*—One hundred and sixty drops of the vinous in-

* See p. 196,

fusion of colchicum, prepared according to the directions of Sir Everard Home* (procured from Fisher's), measuring two drachms, were injected into the jugular vein of a strong dog. For the first fifteen minutes he did not seem to suffer the least inconvenience. In an hour and a quarter, he was sitting on his hind legs; the eyes were bright, and there did not appear any remarkable change. In five hours he was still capable of sitting up, but appeared much dejected; and there was some difficulty of breathing, attended with an occasional husky cough. His pulse was 113; the inspirations were 56. In five hours and a quarter, he vomited some bloody mucus, and expired.

Dissection.—The stomach was in a state of gangrene; the duodenum, jejunum, and ilium were in a high state of inflammation, approaching to gangrene. About two ounces of highly offensive grumous blood were found in the stomach; there was also blood in the duodenum, but not offensive. The colon, cæcum, and rectum were much inflamed; with, here and there, deep rose-coloured spots of the size of a pea. One ounce and a half of fluid was taken out of the cavity of the chest. The arteries contained very little blood. The veins, auricles, and ventricles were much distended with blood of a purplish black colour, not coagulated. The bladder was nearly full of urine of a deep saffron colour; and the gall bladder was much distended with bile. There was an effusion of bile upon the liver.

EXP. 8.—*Tincture of Colchicum.*—At one p. m. one hundred and sixty drops of this tincture were injected into the jugular vein of a mixed terrier of middle size.

* See p. 195.

On being immediately loosed, he staggered considerably, and fell down a pair of stairs.

Forty minutes past one.—He foamed at the mouth.

Two o'clock he was languid and lethargic.

Four o'clock.—Extremely lethargic; and a tremulous motion pervading the whole body.

Six o'clock.—He remained much in the same state, except that his inspirations were deeper, and only twelve in the minute.

Seven o'clock.—The inspirations were only ten in a minute; the eyes were fixed; the teeth firmly closed; and he was to all appearance dying.

At nine o'clock he had expired.

Dissection.—The heart presented the most extreme appearances of inflammation, looking even black and gangrenous. There were about two drachms of water in the pericardium; the stomach was slightly inflamed; the inflammation increasing in the duodenum, and extending in a slighter degree as far as the rectum; the bladder was distended with urine.

EXP. 9.—*Expressed Juice of the fresh roots of Colchicum.*—At 40 minutes past one o'clock, *p. m.* a hundred and twenty drops of the expressed juice of the fresh roots of colchicum were injected into the jugular vein of a young terrier. Immediate signs of great debility were produced; he passed a natural alvine discharge.

In two minutes he vomited about half an ounce of frothy bile of a gamboge colour. Rising upon his legs, he staggered as if intoxicated, and immediately fell. In five minutes the respirations were 52. In seven, the eyes were fixed; the pupils contracted; the hind extremities were stiff and drawn up. In ten minutes he was upon his legs, passed a scanty bilious discharge, walked about, and

soon again made great efforts, without effect, to relieve himself; uttering at the time a cry of distress. He ran into a dark part of the room. On lying down, he had some rigors, and there was a profuse salivation.

Three o'clock.—He was on his legs, with his back raised, and shewing signs of distress in the bowels. The inspirations were 42; he was capable of walking, but seemed very languid.

At four o'clock the lassitude had increased, and he supported himself with difficulty on his legs; a copious discharge of mucus or saliva was proceeding from the mouth.

A little before five, he appeared to be suffering much; was moaning continually, and took no notice on being disturbed.

At six o'clock, he was still much distressed in his inspirations, and these were only twelve in the minute; the eyes were fixed, the teeth firmly closed, and he appeared to be dying.

At seven o'clock there was no perceptible change, except that his inspirations had increased to 14 in the minute.

Nine o'clock.—He respired only 8 times in the minute, with much difficulty. His moaning was much fainter.

Ten o'clock.—He was found dead, and was quite cold.

Dissection.—The stomach was highly inflamed, containing about an ounce of dense mucus, mixed with grumous blood; the duodenum, jejunum, and ilium were highly inflamed; the inflammation decreasing in the colon, cæcum, and rectum. The whole intestines were lined with coagulated albumen. The blood found in the auricles and ventricles of the heart was fluid, and very black; the gall-bladder was much distended with bile.

EXP. 10.—*Acetum Colchici*.—Three fluid drachms of the acetie preparation of colchicum, neutralised by carbonate of ammonia, were injected into the jugular vein of a terrier dog of middle size.

In three quarters of an hour he seemed languid, and trembled a good deal, but was capable of running about. In three hours he exhibited great lassitude; the pulse was 108, and irregular. In five hours and a quarter, on our trying to ascertain his pulse and other symptoms, he started off, as if quite well. In another hour, he was seemingly free from pain. Soon after, the eyes were bright and open. In ten hours he appeared tolerably well.

On the following morning, at eight o'clock, he appeared quite recovered; looked lively; and was sitting up. There was no disposition, however, to move; he looked frightened. At six in the evening, he was found in the same state. He had throughout the day assumed a sullenness of appearance; had not moved from the spot where he was left last night; refused both meat and drink; but had still a lively eye, and did not seem so much inconvenienced by the medicine, as alarmed by the operation.

On the following day he had all the character which appeared yesterday. In the course of this day he passed numerous alvine evacuations, and also urine very freely. He gradually recovered.

EXPERIMENTS BY THE MOUTH.

EXP. II.—*Eau Medicinale, with the Sediment which it forms, shaken up*.—At half past ten *a.m.* eighty drops (by measure, sixty minims) were given to a large and very strong rough terrier.

Two o'clock.—He was lying down, and looked lethargic.

Half-past four.—Pulse 95, vibrating and intermitting every five beats.

Eight o'clock.—The pulse was softer; he had vomited some frothy mucus, and appeared very languid.

Next morning.—Pulse 164, and irregular. The dog had recovered his strength.

Ten o'clock.—A further dose of 160 drops was given.

Half past two.—He looked dejected; the pulse was 104, and very irregular.

Four o'clock.—Had brought up some opaque viscid mucus.

Six o'clock.—Had vomited a quantity of frothy slime, mixed with blood, and appeared altogether very ill. Rigors; the pulse was 80, and small, with intermissions after every five or six beats.

Half past nine.—Seemed dull and languid; the pulse was 106.

On the following morning, at ten o'clock, he was found extended on the ground; had voided a quantity of urine; he was quite insensible, and now and then stretching out his limbs, and was much distressed; his inspirations were six in the minute.

At one o'clock he did not seem to suffer.

Half past two.—He was stretching himself out, as if in the act of dying. There were slight convulsions of one of the legs. The pulsation of the heart was not to be felt; and no distinct respiration could be observed.

At three o'clock he had expired.

Dissection.—The stomach was highly inflamed, and contained a dark brown fluid; the marks of inflammation increased in the duodenum, and through the jejunum—decreased in the ilium, and increased again in the colon,

which appeared in a state of general ecchymosis from venous blood extravasated under the mucous membrane. The cæcum and rectum were slightly inflamed.

EXP. 12.—*Expressed Juice of Colchicum, followed by Eau Medicinale, clear and turbid.*—Monday, at half-past twelve *p. m.* two fluid drachms of the expressed juice were given to a very powerful, lively dog, of the rough terrier kind.

Half-past two.—He was lying down, trembling; the pulse 108.

Four o'clock.—Looked dejected; had been making attempts to vomit.

Six o'clock.—Extremely languid and dull; pulse 180; had passed an alvine discharge, which was mixed with mucus, and of a putrid appearance.

Half-past nine.—There was increased lassitude, and the animal could not stand; pulse 160.

Tuesday morning—Appearances much the same as last night; refused to eat.

One o'clock.—Continued very languid; the skin hot; the pulse feeble, and 184 in the minute.

Half-past two.—There was no change.

Five o'clock.—Appeared exceedingly dull, and staggered when put upon his legs; the pulse 166.

Eleven o'clock.—Pulse 186; no further change.

Wednesday morning, eleven o'clock: the pulse 200, and very small; he appeared better. At twelve, he passed a watery alvine discharge; was thirsty, and drank freely of water.—At half past twelve, the dog took a second dose of two drachms of the expressed juice.—At two o'clock was very ill; the pulse 186; the inspirations 18.—Five o'clock. He continued languid and dull; the pulse was 180.

Eight o'clock: the pulse too feeble to be counted; he looked worse.—Eleven o'clock: the pulse was only 90; he looked better, and drank some water.

Thursday morning.—He appeared to be recovering; the pulse was 100. He continued to mend through the day; and towards evening ate very heartily, but was still dull. In the course of the day he passed a considerable number of discharges mixed with blood.

Friday morning.—The dog still looked dull; but was stronger, and had mended considerably. He ate heartily; and during the day had more of the discharges before described.

Saturday.—Had still the heavy look; and, though he ate with good appetite, appeared yet under the influence of the medicine; the pulse was 166.

Sunday morning.—He was looking much better, was more lively, and running about; the pulse was 168.

Monday morning.—Appeared quite well; he ate heartily, and was in very good spirits.

Tuesday morning.—At ten o'clock, this dog being perfectly recovered, sixty minims of eau medicinale, filtered so as to be quite transparent, were given; his pulse then being 180, and regular.

At half past twelve, no apparent change had taken place, except in the pulse, which was 168, and regular.

Half past two.—No alteration.

Half past four.—The pulse, and appearance of the animal were the same. He continued quite well throughout the evening, and without any variation in the pulse; ate with the best appetite, and was in excellent spirits.

Wednesday morning.—He was perfectly well; the pulse was strong, and 180.

Twelve o'clock.—He took a further dose of two fluid

drachms of the *clear* eau medicinale. In a few minutes he had two unnatural discharges.

Two o'clock.—The pulse was 174; no other change.

Three o'clock.—He looked lively, and there was no visible alteration; the pulse was 160, and regular.

Half-past four.—His spirits were not so good; there was no desire to move; but the pulse remained unaltered.

Eight o'clock.—He had passed several discharges containing much blood; looked very dejected, and his pulse was 156.—Half-past nine: great reluctance in moving.—Eleven o'clock: no change.

Thursday morning.—Looked better; was lively and playful; the pulse was 168; he would not eat. Continued to improve throughout the day; and at eleven *p. m.* the pulse was 168, and regular.

Friday morning.—Appeared quite well; the pulse was 160.

On the following *Monday*, at eleven *a. m.* the dog having completely recovered his strength, and being in the best spirits, was made to swallow a bottle (two fluid drachms) of eau medicinale with the sediment shaken up; the pulse being 168.—One o'clock: the pulse was 192; but he did not appear at all affected.

At half-past three, he threw up some undigested food; looked less lively; and the pulse was more feeble, but still 192.

Six o'clock.—The pulse was 168; weak and irregular.

Ten o'clock.—Had passed some discharges tinged with blood; and a considerable quantity of undigested food had been ejected from the stomach, mixed with mucus and tinged with blood; he looked dull.

Tuesday morning.—Had passed several bloody discharges of the description before mentioned; his strength

did not seem much impaired; but the pulse was very weak, and 180.

Three o'clock.—The pulse not altered; his appearance was improved, but there was no appetite.—Sixty minims more of turbid eau medicinale were now given.

Six o'clock.—The pulse was 166, and very weak; the animal looked dull.—Eight o'clock: no change.

Eleven o'clock.—The pulse and appearance the same.

Wednesday morning.—His appearance was most unfavourable; he looked dejected; was much debilitated; and his pulse so feeble and fluttering that it could not be counted.

Three o'clock.—He was altered for the worse; the pulsations of the heart were scarcely to be distinguished; he had voided some liquid bloody discharges.

At six o'clock, the pulsations of the heart could not be felt; the inspirations were only eight in the minute; all power of motion was lost, and he appeared to be almost dead; his extremities being quite cold, and the body extended on the ground.

Seven o'clock.—He was still living, but the inspirations were only four in the minute; and he was completely insensible.—Eight o'clock: was found dead.

Dissection.—The stomach contained a quantity of bloody fluid, and was in a state of gangrene. The duodenum, jejunum, and ilium were much inflamed, and here and there marked with livid spots. The inflammation increased in the colon and cæcum, and was still greater in the rectum. The auricles and ventricles contained black fluid blood; the liver and lungs appeared healthy.

EXP. 13.—*Eau Medicinale in the transparent state, followed by the Tincture of Colchicum.*—Ninety drops, measuring sixty minims, of eau medicinale, filtered and

quite transparent, were given at ten o'clock *a. m.* to a strong terrier dog.

At two o'clock he was found vomiting a quantity of thin transparent mucus; the pulse was 150.

Half past three.—He had vomited more mucus; and looked rather languid.

Half past four.—Had again vomited mucus.

Eight o'clock.—More frothy mucus from the mouth.

On the following morning, it appeared that he had in the night passed a quantity of liquid alvine discharge, tinged with blood, and mixed with mucus; the pulse was 168, and irregular.

Half past two.—His looks were dejected; the pulse was 104, and very irregular.

Four o'clock.—Had vomited some dark viscid mucus.

Six o'clock.—He had brought up a quantity of frothy slime mixed with blood; seemed greatly affected; had rigors; the pulse was 80 and small, with an intermission at every fifth or sixth beat.

Half past nine.—The pulse was 106; he looked dull and languid.

On the following morning he appeared quite recovered.

The next day, at half past twelve, six drachms of the tincture of colchicum*, of the shops, were given to this dog; he soon became languid, and looked exceedingly heavy and dejected.

At two o'clock he was very ill; the pulse was 96; the inspirations were laboured, and only 24 in the minute. There was a copious discharge of viscid mucus from the mouth.

At five o'clock the pulse was 120, and small; the breath-

* Prepared in the proportion of four ounces of the dried roots to a pint of proof spirit.

ing was laborious; and he appeared to be in pain. He shewed distress on pressure of the bowels.

Eight o'clock.—Moaning, as if suffering pain; the inspirations were 48 in a minute, and were quick and difficult.—Eleven o'clock: the pulse was extremely small, and 180; he still appeared to be in great pain, and very languid; he was uneasy, more particularly on pressure being made on the sides of the abdomen.

On the ensuing morning he was found dead and quite cold.

Dissection.—The stomach was much inflamed, as was also the œsophagus, and the whole of the intestines. The inflammation was most marked in the jejunum and colon; the cæcum and rectum were also inflamed, and interspersed with venous ecchymosis. The bladder was contracted; the gall-bladder was much distended with bile; the stomach contained grumous and offensive bloody serum; a layer of coagulable lymph covered the internal surface of the large intestines very generally.

Exp. 14.—*Wilson's Tincture.*—At half past one *p. m.* two fluid drachms were given to a common-sized terrier.

At half past two he looked languid; the pulse was 107, and irregular.

Four o'clock.—The lassitude had increased.

Six o'clock.—He had thrown up some frothy mucus; the pulse was 120.

Half past nine.—He appeared better; the pulse was 140, and soft, but rather irregular.

The next morning he was on his legs, and looked much better, but refused to eat.

At eleven o'clock he appeared altered for the worse.

One o'clock.—The pulse was 180 and very hard; he looked very languid, and still refused to eat.

Five o'clock.—The pulse was 90; the animal appeared more dejected.

Eleven o'clock.—Excessive lassitude; the pulse 86.

The following morning he was much better; the pulse 164, and regular, but hard.

At twelve, a second dose of four fluid drachms of this tincture was given to him.

At two, he appeared exceedingly ill, was very sick, and had brought up a small quantity of mucus tinged with bile; pulse 180.

Five o'clock.—The pulse 170, and small; he moaned on pressure being made upon the abdomen.

Eight o'clock.—The pulse 160, and very languid; he was suffering great pain.

Eleven o'clock.—The pulse 80; the animal was incapable of standing; the respiration was slow and very laborious. The next morning he was found dead.

Dissection.—The stomach was highly inflamed. Inflammation also appeared throughout the intestines, and particularly in the colon. The stomach contained a brownish fluid; and the ilium, some offensive bloody serum. The bladder was distended with urine. There were livid spots distributed on the surface of the lungs; the gall-bladder was distended with bile.

Exp. 15.—*Reynolds' Specific.*—At eight *a. m.* four fluid drachms were given to a strong lively terrier, of middle size; his pulse was 160, and very regular.—Ten o'clock: he had thrice vomited some frothy mucus; looked rather dull, and trembled universally; the pulse was become more feeble and indistinct.—Eleven o'clock: he had again twice vomited more matter of the above description.—Twelve o'clock: this had again happened three times, together with two or three alvine discharges, one

of which was slightly streaked with blood; the animal looked very dejected.—Half past 2 o'clock: his pulse was fluttering, feeble, and 182; there was another bloody alvine discharge.—Six o'clock: his pulse was 216; there was great debility, and occasional moaning; he loathed food.—Ten o'clock: no discharge; his pulse was 200; and the animal still looked very dejected.

Tuesday morning.—His pulse was 160, hard and fluttering; he had passed in the night two very bloody discharges; his appearance was improved; but he refused meat.—At 8 o'clock, two drachms more of the specific were given to him.—Ten o'clock: he passed a copious liquid discharge, almost the whole of which appeared like blood; his pulse very feeble, and 186.—Twelve o'clock: vomited the same description of matter as yesterday three or four times; the pulse 192.—Five o'clock: he seemed better; the pulse was now 172, and regular.

Tuesday p. m. 10 o'clock.—The pulse was 192, very feeble and regular; the dog appeared restless.

Wednesday morning.—He was better, but still very dejected and hanging down his head; the pulse was 192, feeble and regular. He continued to mend throughout the day, but never so far as to take food.—Seven o'clock: half a bottle more of the specific was given to him.—Eleven o'clock: he was very dull; there was no further change, nor any evacuation.

Thursday morning.—He had been very sick in the night, bringing up a quantity of bright bilious coloured frothy mucus; the pulse was very feeble, and 192.—Two o'clock: had passed two small bloody discharges, was very ill, and loathed food; he continued in this state throughout the day, the pulse remaining the same.

Friday morning, half past two o'clock.—His pulse was so feeble as scarcely to be distinguished; the breathing laborious and slow.

Saturday, eleven *a.m.* the dog much recovered, and his pulse 168. He took a further dose of five drachms. Four o'clock: the inspirations were not more than 12 in a minute; the pulse not to be felt. There was an involuntary discharge of bloody liquid matter from the rectum, and of bile from the stomach.—Six o'clock: he was found dead.

Dissection.—The stomach was distended with air, and contained some offensive bloody fluid; the mucous membrane was in the highest state of inflammation, which extended throughout the intestines as far as the colon, this being in a state of gangrene: there was an intromission of the colon and ilium into the cæcum, which was gangrenous: the intestines contained a quantity of the same fluid as was found in the stomach. The right ventricle was distended with black coagulated blood; the left was contracted, as also both auricles. The appearance of the brain was healthy.

EXP. 16.—*Vinous Tinctures of Hellebore and Opium.*—Sixty minims of this mixture, in the proportions mentioned at p. 196, were given to a fine setter dog, on Monday, at 20 minutes past twelve. There was no change in the animal till after 4 o'clock; his pulse was then only 98, and intermittent.—Six o'clock: the pulse was 80, and intermittent.—Eight o'clock: the pulse intermitted after every 5 beats, and was hard, vibrating, and irregular.

Tuesday morning.—The pulse 92, irregular and intermitting: 160 drops of the same mixture were given to him at 10 o'clock; he immediately discharged from his mouth an abundance of frothy saliva.—Half past 2 o'clock: he had an appetite; the pulse was then 76, intermitting every 3 beats, and very hard.—Six o'clock: the pulse was 108, and the dog appeared lively.—Ten o'clock: the pulse was 106.

Monday.—He looked quite well.—Eleven o'clock: half

an ounce more was given to him; the pulse was 120, and intermitting after every 5 beats; the same appearances followed this dose as the last; if at all different they were less remarkable, and he ate some food a few minutes after.—One o'clock: his pulse was 90, very irregular, and intermitting; in other respects he looked well, and at half-past two he made a hearty meal.—Five o'clock: he appeared dull; the pulse was 80, softer and still intermitting.—Eleven o'clock: the pulse was 79, vibrating and intermittent.

Tuesday morning, eleven o'clock: his pulse was 104, and very irregular, but the animal appeared quite recovered.—Twelve o'clock: he took six drachms of the tincture of hellebore; its action upon the salivary glands was instantaneous and profuse, the animal throwing out large quantities of frothy viscid saliva in all directions.—Two o'clock: his pulse was 76, irregular and intermittent. This dog appeared better towards evening, but having made his escape at dusk, the result of the experiment could not be ascertained.

EXP. 17.—*Tincture of Digitalis, followed by Laudanum and Elaterium*.—At half past 1 p.m. 40 minims of tincture of digitalis were given to a terrier dog of middling size; there was immediately a copious foaming at the mouth.—Half past 2 o'clock: a great flow of saliva; he was staggering and looked very dull; the pulse was 125, and irregular.—Four o'clock: he was much better.—Six o'clock: looked stupid, and was extremely feeble; the pulse was 104, and very irregular.

Monday.—He appeared quite well. At 12 o'clock 40 minims of laudanum and 4 grains of elaterium were given to him.—One o'clock: the pulse was 228.—Half-past two o'clock: had vomited some thick mucus mixed with bile, and there was reason to believe that the medicine was also

discharged from the stomach; he ate freely of meat.—Five o'clock: the pulse was 180.—Eleven o'clock: the pulse was 190; he had vomited a quantity of undigested food.

Tuesday morning, 11 o'clock: his pulse was 200, and regular; he appeared recovered.—One o'clock: five drachms of the tincture of digitalis were given to him; this instantly produced a flow of viscid frothy mucus, and he made many attempts to vomit.—Two o'clock: he had vomited some undigested food mixed with mucus, and which appeared coloured with the tincture; the pulse was 186, and irregular; there continued an abundant discharge of viscid mucus from the mouth.—Five o'clock: pulse 200; respiration quick and laboured; he had vomited more mucus tinged with blood; great lassitude.—Eight o'clock: the pulse was feeble and rapid, beating 210; the animal made continual attempts to vomit.—Eleven o'clock: the pulse was 200: there were incessant attempts to vomit; the respiration was very quick.

Wednesday.—Was much better; the pulse 160; towards evening he had an appetite for food, of which he ate freely; there was a constant dry cough, attended with an effort, as if endeavouring to bring something from the throat.

Thursday.—He appeared lively, and had a good appetite, but was much distressed with the cough before mentioned, and his pulse in consequence could not be counted.

Friday.—He had in a great measure recovered his looks; the cough was better and the appetite natural; the pulse varied from 150 to 164.

Saturday.—He was lively and still better; the pulse was 168.

EXP. 18.—*Want's Gout Remedy*.—At two p. m. two of Mr. Want's powders were given to a large mongrel dog. The frequency of his pulse could not be very satisfactorily ascertained, the dog being particularly timid,

and trembling as soon as he was approached.—Six o'clock: his pulse was certainly slower; but he ate with a good appetite, and was not altered in his appearance. Ten o'clock: no alteration.

Friday morning.—His pulse remained the same; the dog appeared quite well. There had been no unnatural discharge, and he ate very heartily. At ten o'clock, two more of the powders were given to him. Two o'clock: he was trembling, and seemed for the first time affected by the experiment. Six o'clock: looked dejected; was still shaking, and refused both meat and drink; no sickness had been produced; the pulse was slow and irregular. Later in the evening, his pulse was quicker, and much more feeble; but there was not any remarkable lassitude.

The following morning, Saturday, it was observed that he had passed several discharges, two of them bloody and of the worst appearance; his pulse was feeble, and occasionally he moaned; he refused food the whole of the day; but towards evening was better, and his pulse became rather stronger.

Sunday morning.—Appeared to be recovering; ate a little meat, and walked steadily and without difficulty: he continued to improve throughout the day, and at night made a tolerable meal. No further trial was made with this medicine.

EXP. 19.—*Acetic Preparation of Colchicum.*—An ounce, by measure, of this preparation, neutralized by pure magnesia, was given to a large mongrel dog. It produced no visible effect, but his pulse was rather accelerated; he ate and drank freely, and was in good spirits on the following morning.

A few days after, at two o'clock, this dog was made to swallow two fluid ounces of aet. colchici, evaporated

down to half an ounce, mixed with magnesia. This, like the other dose, did not occasion any change in the appearance of the animal. His pulse did not undergo any alteration, and his strength was in no way apparently affected.

On the following morning it was found that he passed two or three copious discharges; but these were quite free from blood: and his pulse was found to be the same as at the commencement of the experiment.

EXP. 20.—*Clear and Turbid Aqueous Decoction of Colchicum.*—On Tuesday, at noon, gave to a young dog of mixed breed, two fluid drachms of a clear decoction of colchicum, prepared by boiling two ounces of the fresh root, in a sufficient quantity of water, down to *two ounces* by measure; his pulse was 132, regular and full.—Half-past two: the pulse was feeble, irregular, and vibrating.—Six o'clock: there had been no apparent alteration; his pulse, however, was increased to 164; he looked lively, and stood erect without inconvenience.—Eight o'clock: no change.—Eleven o'clock: appearances the same; pulse 156; there had been two discharges, streaked with blood.

Wednesday morning.—He had passed, in the night, two further discharges streaked with blood. He now appeared quite well, eating with great appetite, and in good spirits; the pulse was weak, but regular, and 174.—At half-past one, gave him half an ounce of the same decoction, with its sediment shaken up.—Three o'clock: pulse 180, but so feeble as scarcely to be perceptible; and he looked dull.—Six o'clock: he had brought up a large quantity of undigested food; the pulse was extremely feeble and rapid, being about 240 in the minute. He seemed very weak, but was sensible and apparently free from pain.—Two o'clock: he was greatly debilitated, and again

vomited; the pulse could not be counted.—At eleven o'clock he was scarcely able to stand, and looked very heavy; the pulse still so frequent that it could not be counted.

Thursday morning.—He had lost all power of voluntary motion, and was lying extended on the ground, as if in *articulo mortis*. There had been an excretion of blood from the rectum, and in the night some more bloody discharges.—At half-past twelve he died.

Dissection.—The stomach was empty, and slightly inflamed. The inflammation was observed in the same degree through the duodenum, jejunum, and commencement of the ilium, increasing towards its termination, and also throughout the colon and cæcum; the rectum was slightly affected; and the left ventricle of the heart contained a small quantity of dark fluid blood.

EXP. 21.—*Extract from Eau Medicinale.*—*Monday*, at half-past ten o'clock, two fluid drachms of a mixture of this extract, in the proportion of $5\frac{3}{4}$ with 100 of fluid, were given to a large spare dog of the mixed breed. The pulse before the experiment was 168, and intermittent.—Twelve o'clock: no change.—Six o'clock: he was looking dejected, and loathed food; there had been no sickness nor unusual evacuations; the pulse was rather slower, harder, and more regular.—Ten o'clock: he had vomited some half-digested food, and had passed an alvine discharge.

Tuesday morning.—His pulse was 188, vibrating and hard; there had been two or three liquid discharges; one very slightly marked with blood: his appearance was improved since last night, but he still refused to eat.—At eight o'clock he took one drachm more of this mixture.—Twelve o'clock: his pulse was 192 and irre-

gular; he made frequent attempts to vomit; there had been two loose discharges of a light appearance.—Five o'clock: he appeared to be in some pain, and was very dull; his pulse was now 208, regular and feeble.—Ten o'clock: the pulse was 216, and regular; there was no other change.

Wednesday morning.—His pulse was feeble, and 192; he appeared rather better.—At ten o'clock *a. m.* as much of the mixture as contained $5\frac{1}{2}$ grains more of extract was given to him.—Half-past three: his pulse was 168, full and regular; there was no evacuation; rigors.—Six o'clock: pulse was 204, and very feeble; he was looking very ill and dejected.—Eleven o'clock: he appeared almost unable to stand.

Thursday morning.—He had discharged, during the night, a considerable quantity of liquid alvine matter, for the most part like blood; his pulse feeble, and 168.—Two o'clock: seemed excessively weak, but did not appear to suffer much pain; he refused food.—Half-past five: gave a further dose of $5\frac{1}{2}$ grains of extract.—Ten o'clock: hitherto no remarkable change.

Friday morning.—He was found dead. Some bloody alvine discharges had been passed in the night.

Dissection.—The stomach was distended with air, and its coats were much inflamed; there was abundance of venous coagula under the mucons membrane, distributed in patches; the stomach contained a small quantity of yellow putrid fluid; there was an intromission of the colon into the cæcum, which latter intestine was in a state of mortification; the right ventricle contained a small quantity of coagulated blood, and also some fibrine; and the aorta, for the space of two inches, was filled with fibrine, completely free from the colouring part of the blood. The brain was found in a natural state.

EXP. 22.—*Extract from Wine of Colchicum* (obtained by evaporation from the wine of the shops, prepared in the proportion of four ounces of the fresh roots to a pint).—At 11 o'clock, a quantity of this extract, equivalent to that furnished by two fluid drachms of eau medicinale, was given to a fine healthy dog of the mongrel breed; the pulse before the experiment was about 150, regular and full.—Four o'clock: the pulse was less full, and 148; he ate freely and appeared lively.—Ten o'clock: the pulse so exceedingly feeble, as scarcely to be felt; he had been very sick, bringing up a large quantity of undigested food and frothy mucus; looked dull and unwilling to be moved. On the following morning he was quite well, and the appetite was good.—Twelve o'clock: the pulse was 168, feeble and rather irregular: we gave him double the quantity just now mentioned of this extract.—At 4 o'clock his pulse was harder, and 156; he refused food, and looked less lively.—Six o'clock: the pulse was very feeble, and could not be distinctly counted. He had twice vomited a quantity of viscid frothy mucus, and looked dejected.—Ten o'clock: he had passed two very bloody discharges; his pulse was now so feeble as scarcely to be felt, and the animal looked drooping and very ill.—On the following morning he continued very languid; the pulse much as on the preceding night; he refused food. Towards noon, he improved in appearance and spirits.—Four o'clock: the pulse was more distinct, but irregular and slow; he still refused to eat.—Seven o'clock: he was much better; standing, and looking very lively; he now ate with a tolerable appetite.—The next morning he appeared less lively, and would not eat. In the night he passed a watery discharge, marked with blood; the pulse was still very feeble and indistinct. He was better towards the evening, and then made a good meal.—On the next morning, he seemed quite well.

EXP. 23.—*Extract from Hellebore, followed by Elaterium, and a mixture of Extract of Colchicum from its Tincture in Cape Wine.*—On Friday, at eleven *a. m.* a drachm of extract of hellebore, the whole quantity furnished by half an ounce of the root in decoction with water, was given in the form of a bolus to a very strong lively dog, of a mixed fox-hound breed; his pulse being 160, full and regular. In a quarter of an hour, he vomited a quantity of transparent mucus, and with it the bolus; his pulse was now 172, and very feeble: he continued sick, and retched most violently during an hour and three quarters.—Two o'clock: the pulse was only 90, hard and regular.—Half-past three: appeared much recovered, and ate with a good appetite.

Saturday morning.—The pulse was 116, full, and irregular. We gave him half a drachm of the extract at half-past eleven.—At a quarter past twelve, the pulse was 126, but more feeble; he was very uneasy, and had made two slight attempts to vomit.—At twenty minutes past one, he brought up some undigested food; but there was no appearance of the hellebore.—Three o'clock: he had thrown up some more undigested food, but there were no marks of the extract; pulse 104, full and irregular.—Five o'clock: no remarkable change in the animal; his pulse had improved, and there had been no further sickness.—Seven o'clock: he appeared quite well; the pulse was 154, strong and regular.

On the following morning, he continued well; the pulse was strong, and 150.—At seven o'clock we gave him a drachm of the extract wrapped up in thin paper. In little more than half an hour, he vomited some frothy mucus, mixed with undigested food, but there was no appearance of the hellebore; the pulse was 152, hard and intermittent.—Half-past ten: had brought up some more undigested food, and with it a great part of the hellebore.

Monday morning.—His appearance was quite good; but the pulse was only 108, and very irregular.—At nine o'clock, he was made to swallow a drachm of elaterium; there was instantly a copious salivation.—Eleven o'clock: the pulse was intermittent, and 136; the dog trembled much, and looked very dull; he had vomited some viscid mucus; we could not ascertain whether any of the elaterium was amongst it.—One o'clock: he had brought up from three to four ounces of bright bilious fluid; his pulse was now 228, and very irregular; he drank freely of water.—Four o'clock: had a liquid alvine evacuation, with great effort in passing it; the pulse was 240.—Six o'clock: no visible change; the pulse was the same.

The following morning he was quite well, and ate with a good appetite.—At half-past one, two fluid drachms of an extract of colchicum, obtained *from its tincture*, in Capewine (100 parts containing $5\frac{1}{4}$ of the extract, the quantity yielded by eau medicinale, see p. 218), were given to this dog, his pulse being then 192, and regular.—Three o'clock: the pulse quicker.—Half-past four: it was upwards of 200, and so rapid as to be counted only with difficulty; he had brought up a quantity of undigested food.—Eight o'clock: had evacuated some very viscid bloody mucus; was uneasy, and looked very dull; the pulse was the same in frequency, but rather more feeble. Half-past ten: had evacuated more bloody mucus; suffered pain on pressure of the abdomen; looked very languid; the pulse continued to be upwards of 200; it was in a great measure regular, but feeble.

Wednesday morning, June 13.—He appeared better; the pulse was 212, and more feeble than last night.—At nine o'clock, he took sixty minims more of the solution.—Half-past ten: had vomited some frothy mucus, tinged with blood. There were some appearances of a part of the mixture amongst it; but this could not be ascertained

with certainty. He was crouching, and seemingly very weary; he refused food.—Half-past eleven: the pulse was 200, and rather more distinct; was much distressed on having the abdomen pressed.—Half-past one: the pulse was more feeble, and 210; he looked dejected, and his strength was much impaired. He staggered as he walked, and moved with evident difficulty.—Three o'clock: the pulse as before, but more irregular; he had vomited a considerable quantity of transparent frothy mucus.—Seven o'clock: appeared very dejected, and unwilling to move. These appearances continued till midnight, when he was last visited.

Thursday morning.—At two o'clock, life appeared to be nearly extinct; the animal was extended on the ground, and quite insensible; his inspirations were about twelve in the minute; and the pulsations of the heart so languid as scarcely to be felt. At seven o'clock he was found dead.

Dissection.—The stomach was distended with air, and contained about an ounce and a half of a yellow, viscid, fluid. It was highly inflamed throughout, as was also the omentum. The whole intestines were slightly inflamed; but the rectum bore the strongest marks of inflammation, and under its mucous membrane were many small coagula of venous blood. The bladder was distended with urine of a saffron colour; the gall-bladder was also much distended with bile; the lungs were slightly inflamed. The ventricles of the heart contained fluid blood of a very dark colour.

EXP. 24.—*Extract from the Acetic Preparation of Colchicum.*—Ten grains of this substance, mixed with water, were given to a large strong mongrel dog. This had the effect of raising the dog's pulse considerably at

first; but it afterwards became depressed, and towards evening he was greatly affected. On the following morning he appeared better, but was still much dejected and refused food. He had, in the night, evacuated some bloody discharges. He recovered towards the close of the day; and on the succeeding morning was quite well, and ate and drank freely.

This dog passed an extraordinary quantity of urine, whilst he was under the influence of the above experiment.

EXP. 25.—*Extract from the Acetic Preparation of Colchicum.*—March 5, at 11 o'clock, ten grains of this extract (obtained as before by evaporation of the Acetum Colchici), and twenty grains of calcined magnesia mixed with water, were given to a very lively, strong, mongrel dog; his pulse being regular, and about 144 in the minute.

There was no alteration in the animal till six o'clock; he then drank some water, which he instantly threw up, together with a quantity of frothy mucus and undigested food. This was followed by a copious alvine liquid discharge. The pulse remained unaltered. At this time he passed an unusual quantity of high-coloured urine; he refused meat.

March 6.—He seemed in good spirits this morning, was jumping about, and apparently free from all inconvenience; but his appetite had not returned. He passed, during the night, two very copious fluid evacuations, only slightly marked with blood. Before night, he made a very plentiful meal, and was in all respects quite well.

EXP. 26.—*Extract from the Proof Spirit Tincture of Colchicum, prepared with the dried roots, in the proportion of four ounces to a pint.*—At half past one, two

fluid drachms of a mixture of this extract and water, in the proportion of $5\frac{1}{4}$ grains in one hundred, were given to a strong full-sized dog of the bull species. The pulse before the experiment was 156, hard and rather irregular. Seven o'clock: no change was perceived till now, when it was observed that he had brought up a small quantity of viscid mucus, and the pulse was considerably quicker. Later in the evening, the pulse was more indistinct and very quick, and the dog had lost his spirits.

The next morning he had passed two very bloody discharges, and had been very sick, bringing up a quantity of mucus, which, in one instance, was of a bright yellow colour. His pulse was very variable in frequency, vibrating and extremely irregular; the animal looked greatly dejected, the skin was very hot to the touch, and he seemed to feel increased pain on pressure of the abdomen. At half-past eleven, we gave him one fluid drachm in the same proportion. Three o'clock: he had twice thrown up some frothy yellow mucus, was very dull, and unwilling to walk.—Seven o'clock: he appeared much the same, and had vomited mucus similar to that before described. Eleven o'clock: a change had taken place for the worse; he lay motionless on the ground; and on being urged, walked with great difficulty. His pulse could not be felt, and he appeared to be dying.

On the following morning the animal appeared stronger, and in other respects better than before, but was still unwilling to move; the pulse was 126, very irregular and fluttering. He continued to improve throughout the day, but appeared uncommonly lethargic; his pulse was very irregular; the skin was extremely hot. At six o'clock he took some meat with a tolerable appetite. In the course of the day he passed a bloody discharge.

Thursday morning.—He was nearly recovered in all

respects, eating heartily; and his spirits were as good as before the experiment.

Exp. 27.—*The same Extract inserted into a wound.* Twenty grains of this active extract (obtained from the tincture mentioned at p. 510) were inserted into the inside of the thigh of a small dog, at a quarter past nine o'clock *a. m.* The pulse, at the time of the experiment, was 140 in a minute, and regular.

Eleven o'clock: he had vomited some undigested food, and passed two alvine evacuations, one of which was mixed with bile of a bright yellow colour: the pulse was more rapid and less distinct.—Two o'clock: the pulse was 154, hard and regular. He had passed two other loose discharges, one of a pale colour; the other much marked with blood.—Three o'clock: he had passed another small but very bloody discharge; the pulse remained the same.—Six o'clock: the pulse was scarcely perceptible; the dog appeared lethargic and insensible, and the inspirations were short and quick.—Nine o'clock: the inspirations were deep and very laborious; the pulse was not to be felt; he was in evident distress, and life appeared nearly extinct; there was a discharge of blood from the bowels, and the pain, which was seemingly considerable, increased greatly on disturbing the animal, and especially when the abdomen was pressed with the hand. In a quarter of an hour more the inspirations were only seven in a minute, accompanied with marks of distress.—Half past nine: the inspirations were only four in the minute. At half past ten he died.

Dissection.—The cardiac orifice of the stomach was much inflamed, but the pylorus had not any appearance of inflammation. The stomach contained a quantity of blood and mucus; the intestines were highly inflamed

throughout. The thigh upon which the operation was performed, was black and almost approaching to mortification. The appearances of the heart, liver, and lungs, were natural.

EXP. 28.—*Sediment from Wine of Colchicum.*—At half past eleven *a. m.* ten grains of this sediment, procured from the vinous preparation of Colchicum, which had been set by for use in the transparent state, were given to a large and very strong terrier. The Pulse at the time of the experiment was 114 and intermittent. Two o'clock: no visible effect had been produced; the pulse continued slow, irregular, and intermittent. Three o'clock: the pulse was 94 and rather fuller. Five o'clock: no alteration. Seven o'clock: he appeared lively and as well as before. Ten o'clock: the pulse was 90, fuller, and more regular.

On the following day, ten grains more were given; but this quantity, like the former, had no remarkable effect.

Two days having elapsed, ten grains more of sediment were given him; his pulse at the time of the experiment was 108 and very irregular. Three hours after, he appeared quite well, was in good spirits, and running about the yard. Two days after, a further dose of ten grains of sediment was given; but even from this additional quantity no apparent change took place in the animal.

EXP. 29.—*With the Macerated Roots left by the Vinous Infusion**, prepared according to the direction of Sir Everard Home —On Friday at 12, *m.* the produce

* See p. 195.

of an ounce of these roots dried, boiled in water, and reduced by evaporation to the state of an extract, which weighed 96 grains, were given to a large strong mongrel dog. His pulse before the experiment was 156, weak, and irregular.—Three o'clock: the pulse was weak, slower, and more irregular; his appearance not changed.—Six o'clock: the pulse was quick, but more feeble, and 192 in the minute; he refused food, but drank freely of water, some of which was immediately rejected from the stomach; he looked dull.—Ten o'clock: he appeared considerably weaker, and looked very dejected; he again drank most freely of water, some of which he brought up immediately. His pulse seemed to be more regular, yet less distinct, and in frequency 198; he refused meat.

Saturday morning, nine o'clock, his pulse was very feeble, and still rather quicker; his thirst was quite insatiable; and he again brought up a great part of the water he had been drinking. There has been this morning a considerable alvine evacuation of blood, both liquid and coagulated.—Twelve o'clock: his pulse was with great difficulty ascertained to be 180, and extremely feeble; he was evidently distressed, and had passed another bloody discharge.—Four o'clock: two more bloody discharges were observed; he looked exceedingly heavy.—Seven o'clock: he continued in the same state. Eleven o'clock: was lying extended on the ground, and unable to raise his head; his pulse could not be counted, and the respiration was very slow. At twelve o'clock he was dead.

Dissection.—There was a slight inflammation at the pyloric orifice of the stomach, which continued throughout the mucous membrane of the intestines. The stomach was distended with air. Some coagulated albumen was observed in different parts of the intestines.

EXP. 30.---*Macerated Roots left by the Vinous Infusion, in the proportion of an ounce to the pint* (the same, namely, as the Pharmacopœia directs for the acetic preparation).---One ounce of the dried roots, which had been digested as above, was boiled in a sufficient quantity of water down to an extract weighing 95 grains. This was given to a very healthy good-sized dog of the mongrel kind, at ten o'clock; his pulse at the time of the experiment being 126, strong, and rather irregular.—Four o'clock: the pulse was 180, full, and more regular; he drank freely of water; appearance unaltered.—Seven o'clock: he vomited some frothy mucus and undigested food; the pulse became more feeble.---At ten o'clock, the pulse was so feeble and rapid that it could not be exactly counted.—At eleven, he looked very lively, but refused to eat; his pulse was very quick, and it still could not be counted with accuracy.

Wednesday morning.---The dog looked as if recovered, but refused to eat; his pulse was regular, and 150 in the minute.---About one o'clock he vomited some bile and mucus. He continued, to all appearance, tolerably well throughout the day, and at ten in the evening took a small quantity of meat, and drank freely of water.

Thursday morning.---He was quite well, and had his appetite as usual.

EXP. 31.—*Macerated Roots left by Proof Spirit, in the proportion of an ounce to the pint.*—One ounce of the above roots, when carefully dried and powdered, was boiled in a sufficient quantity of water down to an extract weighing 54 grains. This was given to a very healthy good-sized dog of the mongrel breed, at twelve o'clock; his pulse being before the experiment 156, and regular.—Seven o'clock: the pulse continued unaltered

throughout the day; nor could any difference be observed in the animal's appearance. At this time also he ate heartily of meat. On the following day he seemed to be quite recovered

EXP. 32.—*Macerated Roots left by Acetic Acid, in the proportion of an ounce to the pint, as the Pharmacopœia directs.*—The produce of an ounce of the macerated roots, treated as in the preceding instances, was given, in the form of extract, to a large mongrel dog, at two o'clock. The pulse before the experiment was 138, and regular. There was a copious secretion of frothy saliva; but no other apparent effect resulted; and at seven o'clock he ate very heartily.

EXP. 33.—*Clear Vinous Infusion of Colchicum.*—At Eleven o'clock *a. m.* sixty minims of the clear vinous infusion of colchicum, prepared according to the direction of Sir Everard Home*, were given to a large and very strong rough terrier.—At half past three he had passed an abundance of urine.—Half past four: the pulse was 150 and irregular.—Eight o'clock: no further visible change.

On the following morning he looked quite well; the pulse was 144, and regular. At half past one, two fluid drachms were given; this, in the course of the day, produced several copious evacuations; but the dog did not appear much inconvenienced at any period by it.

On the ensuing morning he was apparently well, but refused to take any food.—At 11 o'clock three drachms more were given. The pulse was 130.—Twelve o'clock: the pulse was 144, and regular.—One o'clock: he seemed

* See p. 195.

affected by the medicine, and was considerably dejected; the pulse was 132, and irregular.—Two o'clock: the pulse was 144, irregular and intermittent; he had passed a foetid loose discharge, and a plentiful quantity of urine.—Half past two; he had thrown up some thick bilious-looking mucus, was better, and running about. He drank freely of water, but had no appetite for food.—Five o'clock: he looked lively; his pulse was 190 and hard, intermitting every five beats.—Eleven o'clock: he was lively; the pulse was 188.

On the following morning, at eleven o'clock, his pulse was 140, still intermitting every two or three beats.—At twelve, five drachms more of the infusion were given to him. Two o'clock: he looked heavy, trembled much, and was altered greatly in his appearance.—Five o'clock: the pulse was 120, the respiration laborious; and he was dull and languid.—Eight o'clock: he was in the yard running about, appeared restless, and much under the influence of the medicine; his pulse was so weak as not to be counted. Eleven o'clock; the pulse was 120; he appeared much worse; the respiration was slow, and scarcely perceptible; he had vomited bloody mucus, and had passed some bloody liquid discharges. On the following morning he was found dead; and evidently had expired several hours.

Dissection.—The stomach was much inflamed, and contained a quantity of offensive bloody serum; the pylorus was nearly gangrenous; the internal coat of the duodenum, jejunum, and ilium, was highly inflamed, and almost entirely lined with coagulated albumen. The external as well as internal coats of the colon were more inflamed than the small intestines; and there was effused venous blood under the mucous membrane. The bladder was contracted, and its mucous membrane slightly inflamed.—There were appearances of partial inflammation on the

liver; the gall-bladder was distended; the heart and blood-vessels contained a quantity of black fluid blood. The rectum was much inflamed.

EXP. 34.—*The same Vinous Preparation of Colchicum, with its deposit shaken up.*—At eleven o'clock a.m. sixty minims of this infusion in the turbid state were given to a common-sized shepherd dog. At two o'clock he appeared dejected, and was extended on the straw.—Half past three: no visible change.—Half past four: the pulse was 112, and very irregular.

On the following morning, his pulse was 100, and irregular.—Twenty minutes past one: two fluid drachms were given.—Four o'clock: there was no apparent alteration in the animal.—Six o'clock: the pulse was 120, and intermittent.—Half past nine: the pulse was 160; there was no other change.

On the ensuing morning he looked languid, but there did not appear any notable alteration.—At eleven o'clock, three drachms more were given; the pulse was then 120. Twelve o'clock: he appeared strongly under the influence of the medicine, and his inspirations were only fourteen in the minnte.—One o'clock: he had vomited some frothy mucus; the pulse was 84.—At half past one, he was running about the yard, when he passed a dark, bilious, and very foetid discharge.—Half past two: continued the same; there was no appetite.—Five o'clock: the pulse was 90; he looked dull.—Eleven o'clock: there was great lassitude; the pulse was 86.

The next day, at eleven o'clock, he appeared better; the pulse was small, and 180.—At 12, five drachms more were given.—Two o'clock: the abdomen was hot to the touch; the pulse 228.—Five o'clock: the pulse was 194, and small. He appeared heavy and lethargic, and had vo-

mitted some bloody mucus.—Eight o'clock: he vomited blood; there was no alteration in the pulse or general appearance. Eleven o'clock: the pulse was 100, and extremely small; he suffered great pain, which seemed to increase on pressure of the abdomen and bowels. He was not altered for the worse in his appearance.

On the following morning he was found dead; but had not long expired.

Dissection.—The stomach was found highly inflamed, and approaching to gangrene. The duodenum, jejunum, and ilium, were also much inflamed; and the appearances of inflammation increased in the colon. The rectum was also inflamed, exhibiting an effusion of venous blood in small coagula under the mucous membrane. The stomach was distended, and contained about 3ij of thick yellow serum; in the intestines, grumous blood was mixed with layers of coagulated albumen.

EXP. 35.—*Elaterium.*—Five grains of this active substance, mixed with water, were injected into the jugular vein of a mongrel dog. There was instantly a flow of urine, and also an alvine discharge. In a few minutes he vomited some dark mucus and undigested food. In ten minutes he exhibited great prostration of strength; the pulse was feeble, regular, and 72; the inspirations were alternately quick and slow, and 27 in the minute; the eyes were fully dilated, and appeared almost starting from the head. In half an hour, he appeared distressed; there were frequent convulsive motions of the whole body; the tongue was protruded, and there was a discharge of viscid saliva from the mouth; the inspirations were very laboured, and only twelve in the minute; the pulsations of the heart could not be felt.—In

three-quarters of an hour he appeared to have expired; but, in four minutes, he made more inspirations, quicker, and not so deep. In two minutes, they were slower, exceedingly laboured, and only 7 in the minute. In fifty-five minutes after the operation, he expired.

Dissection.---The stomach was found slightly inflamed, as were the lungs; and marks of slight inflammation appeared partially throughout the intestines.

EXP. 36.—*Resin of Opium.*—At twelve o'clock, a drachm of the resin of opium, with which I was furnished by Mr. Battley of Cripplegate, was given to a terrier dog, his pulse then being 168, and regular. At 20 minutes past one, he threw up a quantity of transparent thin mucus, and made a whining noise, appearing very uneasy; the pulse being 96, irregular and intermittent. In ten minutes more, there was a copious discharge of frothy saliva from the mouth, and the whining continued.—At half-past two, he had several times thrown up more fluid of the above description.—At a quarter before three, he was found staggering, and seemed to have great difficulty in supporting himself; the pulse was 84, hard, irregular, and intermittent.—At half past four, he continued to throw up abundance of mucus; had not slept since he took the resin; the pulse was 114, and intermittent.—Half-past five; he looked dejected; the pulse was only 90; the pupil of the eye was much dilated; he was quiet, and seemed disposed to sleep. Slept till half-past seven; then awoke uneasy, moaning, and very restless; the pulse was 90; the inspirations were 18; and he looked very dull.

Tuesday morning.—At eleven o'clock, this dog appearing to be quite recovered, his pulse regular, and 144 in the minute, fifteen grains of Mr. Battley's sedative

extract of opium were given to him. It had the effect on the salivary glands before described, as produced by the resin, and he made the same plaintive noise.—Two o'clock: he looked dull; the pulse was 84, irregular, and intermittent.—Three o'clock: the pulse was 84; he was lying down, and very dejected.—Half-past four: he looked stupid; the abdomen was remarkably tense; the pupils were much dilated; the animal appeared to suffer great inconvenience; the pulse was 72.—Six o'clock: the pulse was 72; the appearance of the dog was very dull; there was great prostration of strength, and he made a continual whining noise.—Eight o'clock: the pulse was 126; he was standing on his legs with great difficulty, and made much noise.—Half-past nine: he was quiet, and appeared in much better spirits; the pulse was rather slower than before. He now stood with less difficulty.—Eleven o'clock: he was very restless and uneasy; the eyes were bright.

Wednesday morning.—The dog seemed quite recovered, and ate with a good appetite; the pulse was 192.—Eight o'clock *p. m.* he continued to mend; had passed several dark-coloured discharges; the pulse was 192, and irregular.—Eleven o'clock: no alteration.

Thursday morning.—He appeared quite well; the pulse was 160.—At eleven o'clock, fifteen grains of crude opium were given; there was an immediate slight frothing of the mouth, which was not followed by any remarkable appearance in the animal; but he looked rather dull. At three o'clock, his pulse was 120; he had passed a dark bilious discharge. On the following day he was perfectly well.

Saturday, Jan. 2, 1819.—Gave him a drachm of crude opium at twelve o'clock, the pulse being 148. At half-past one o'clock, the pulse was 180, and the appear-

ance of the animal was scarcely changed. At four o'clock, the pulse was considerably lowered, but not to be counted with accuracy, in consequence of frequent shiverings with which the animal was affected. He looked dull, frothed at the mouth a little, was almost incapable of standing, and was sensible.—Six o'clock: the pulse was much the same, and the same difficulty existed of ascertaining its number. He appeared very feeble, and uttered faint plaintive cries.—Ten o'clock: the pulse was only 66; he was extremely languid, and unable to move; the pupils were much dilated, and he seemed not to sleep throughout the day; had refused food since he took the opium, but appeared sensible.

Monday, eleven o'clock.—The pulse was the same, and the general appearances of the animal corresponded with those of the preceding night. He walked with difficulty, dragging the hind extremities, which seemed nearly paralysed; the pupils were more dilated than before; and the animal looked more stupid.—Ten o'clock *p. m.* he gradually mended throughout the day, and, in some degree, recovered his strength; the pulse was 66. The dog was still without appetite.

Monday morning.—The pulse was 168; he looked better; and very shortly he recovered.

FROM the preceding experiments I have been led to draw the following conclusions; which, for the sake of arrangement, and more familiar exposition of the subject, I shall state in the order of the experiments.

EXP. 1.—The symptoms which quickly followed the introduction of the eau medicinale into the circulation,

shew its powerful influence on the nervous system, and would appear sufficient to warrant the conclusion, that by means of this active agency, it produced death; but the extensive morbid changes, found, by dissection, on the alimentary canal, must lead us to consider, that the disturbance of the nervous system, as well as of the circulation, would, in a great measure, arise from the diseased action taking place so rapidly in the stomach and intestines; and therefore the fatal exhaustion produced may in part only be referred to the specific and primary agency of the medicine upon the nerves. This dog survived the experiment twenty-seven hours.

The action of this medicine, in common with other vegetable medicines through the medium of the circulating blood, has already been demonstrated by Sir Everard Home, in his experiments related in the *Philosophical Transactions* for 1816; and the experiments of earlier physiologists to the same effect, are related in the *Edin. Med. & Chir. Journal* for April 1817.

EXP. 2.—The symptoms in this case were fewer, but not very different in character from the former; and the appearances on dissection bore a near resemblance, although they were less in degree. The loss of blood which this animal suffered in the operation, may be considered as having hastened his death. He survived the experiment only seven hours and a half.

EXP. 3.—A larger quantity of this medicine was used, than of either of the two former preparations, as it is directed so to be administered. We are here struck by the great resemblance of the symptoms which were produced. In each instance the action of the liver was excited; but the mucous membrane of the stomach

and intestinal canal was the texture most affected. This animal survived the experiment only seven hours and a half.

EXP. 4.—The very opposite manner in which this compound medicine acted, sufficiently proves its distinct nature from the eau medicinale. The instantaneous and chief influence of this preparation on the nervous system, and the urgency of the symptoms during two hours, with the subsequent speedy recovery of the animal, present a strong contrast to the result of the former experiments, and strengthen the supposition, that death from the other medicines is chiefly to be attributed to the inflammation produced on the alimentary canal.

EXP. 5.—Supports the conclusions just stated.

EXP. 6.—The immediate action of this compound is alone sufficient to prove its difference from the eau medicinale. The first symptoms were very similar to those in Experiments 4 and 5, affecting the nervous system powerfully; but the progress was that of active inflammation, the effects of which appeared very manifest upon dissection. This animal survived the experiment about twelve hours.

EXP. 7.—This preparation of colchicum produced very active effects in a short time. The dog survived only five hours and a half. The nervous system was much less affected than by the eau medicinale; but the appearances found on dissection were very similar from the two medicines. The lungs were affected in the present instance; which did not happen from the eau medicinale.

EXP. 8.—In this experiment the tincture was made of the dried roots, these being used in the proportion of four ounces to a pint of spirit. This is the most active preparation of colchicum which can be made, and is, in my opinion, a very unfit medicine. I have known it to operate as a violent drastic, when added to a simple purgative. In this experiment the effects produced on the heart were quite remarkable, and appear to have been the cause of death. The dog survived nine hours.

EXP. 9.—The effects of pure colchicum are here distinctly shewn; and the blended influence of the medicine upon the nervous system and alimentary canal is well demonstrated. High inflammatory action was produced. In each of the foregoing experiments, the liver was excited to increased action; but this effect may probably be rather referred to the combined derangement of parts, than to any specific influence of either of the medicines in increasing biliary secretion. This dog survived only eight hours and a half.

EXP. 10.—The comparative mildness of this preparation is rendered very evident by this experiment; the nervous system was slightly affected; and the intestinal canal and kidneys were stimulated; but the mucous membrane of the intestines did not appear to have suffered any degree of inflammation.

EXP. 11.—The immediate symptoms appear to have arisen from the disturbance of the nervous system; and those which followed, from the inflammation of the stomach and intestines. The animal survived this experiment twenty-eight hours and a half.

EXP. 12.—The greater activity of medicine, when introduced directly into the circulation, than when administered by the mouth, is here shewn; more than four times the quantity of the expressed juice being used than in the experiment by injection. The animal, although rendered very ill, recovered in a short time.

In the first edition of this Treatise, I noticed, at p. 164, “that in an old bottle of eau medicinale, the sediment which adhered tenaciously to the bottom, was considerable;” and I suggested, that this might serve to explain, in some measure, the very different activity of the medicine at different times. Sir Everard Home has since noticed this circumstance, and verified the opinion by the comparative use of the medicine, with and without the sediment, in his own case. In the present experiment, the dog recovered in a short time from the large quantity of three drachms of clear eau medicinale given in two doses.

The animal was so completely recovered at the end of five days, from the last dose, that he was made the subject of a comparative experiment with the eau medicinale, having its sediment shaken up. The immediate symptoms were very severe; and from the same quantity as before employed (three drachms in two doses), death took place in nine hours. Very similar appearances of inflammation were found in the alimentary canal as in the experiment by injection.

EXP. 13.—It appeared desirable to repeat the experiment with the clear eau medicinale, upon a fresh dog. The symptoms produced, by a dose of 60 minims, were more urgent than I should have expected; and the effect on the mucous membrane of the intestines was very con-

siderable ; but the animal speedily recovered. The stimulating properties of the tincture of colchicum are strongly exemplified in the appearances, on dissection, of violent inflammation throughout the alimentary canal, even from the œsophagus ; but the heart was not in any degree inflamed ; a result which was produced by this medicine in the experiment by injection. The dog survived the second part of this experiment probably about 16 hours.

Exp. 14.—The effects produced in this experiment, shew this tincture to be a very active medicine, although less so than the eau medicinale. Of the latter medicine, three drachms (Exp. 11) proved destructive in twenty-nine hours : whereas, of Wilson's tincture, six drachms were given ; and we may reckon that death did not take place until at least forty hours had elapsed. By the second dose of the eau medicinale, the pulse was rendered slower, while the opposite effect was produced by the second dose of Wilson's tincture.

The appearances found on dissection were not in any material degree different.

Exp. 15.—By this medicine, the first action upon the pulse was that of rendering it quicker ; and the same effect was produced by the different preparations of colchicum, and by Wilson's tincture ; but in each instance the pulse appears to have been at first rendered slower by the action of the eau medicinale. This tincture affected the mucous membrane of the intestines very severely. It is an active preparation, but more diluted in its menstruum than the other quack medicines. Twelve drachms were administered in four doses ; and the animal survived the experiment four days and a half. The ap-

pearances on dissection very much corresponded with those produced from Exp. 3.

Exp. 16.—As from the eau medicinale, the pulse was rendered slower from this compound preparation; but the different result of the experiment, as in that by injection, completely negatives the opinion of Mr. Moore, of the identity of this medicine and the eau medicinale. It seems remarkable that the animal should bear such large doses of hellebore with so little comparative injury.

Exp. 17.—The chief intention of this experiment was in reference to the opinion which seems to have been conceived*, that the eau medicinale is a composition of elaterium and laudanum; and further, to ascertain, by the administration of that active medicine the tincture of digitalis, how far the dog is susceptible to the influence of the powerful articles of the materia medica; enabling us in this way to reason more correctly in regard to the doses, and their effects, of the gout *specifics* on the dog. The inferences suggested by the results of this experiment appear to me sufficiently obvious. See also Exp. 6. and observations on the same, p. 524.

Exp. 18.—This medicine is, from its sensible properties, very evidently a powder of colchicum mixed with some other substance. So far as it was tried in this experiment, the effect upon the intestinal canal was similar to that produced by the other preparations of colchicum. The other preparations, being in solution, would of necessity operate in a much more active manner.

EXP. 19.—The extraordinary quantity of this preparation, amounting to three ounces in two doses, administered not only without fatal result, but even without serious symptoms being produced, is a striking proof of the modification effected in the properties of the colchicum by means of the acetic acid as a menstruum. It is worthy of observation, that, although a considerable purgative operation was produced, the mucous membrane of the intestines did not appear to have been irritated. No blood was discharged.

EXP. 20.—The intention of this experiment was to ascertain the power and fitness of water as a menstruum in the preparation of colchicum. The proportions of the fresh roots were the same as directed by Sir Everard Home in his preparation. The portion poured off clear, in the dose of two drachms, produced considerable irritation of the mucous membrane of the intestines; and a second dose of four drachms of the turbid liquor proved fatal in twenty-two hours and a half. In the turbid state, the colchicum was very much concentrated. The superiority of acetic acid as the menstruum, is sufficiently apparent when brought into contrast with the present simple preparation. The influence of the colchicum in increasing the frequency of the pulse, is again worthy of observation, as illustrated by this experiment.

EXP. 21.—In this and the following experiments with extracts, my chief object was to ascertain the possibility of making a preparation of colchicum equally active with the eau medicinale. At page 218 I have stated that 100 parts of eau medicinale furnished $5\frac{3}{4}$ grains of extract; and such was the exact quantity from that specimen. But other vials have furnished a larger

quantity, and hence another explanation is afforded of the uncertain activity of the medicine administered in the same given doses. In these comparative experiments, however, no fallacy arises from this circumstance, as the proportions employed were of extract itself, and of the same consistence. The usual liquid contents of a bottle of eau medicinale are 120 grains. It appears that in the present experiment $16\frac{1}{2}$ gr. of the extract were administered before death was produced. Now, as one vial which was evaporated for the purpose, furnished eleven grains of extract, and another only eight, we may calculate the $16\frac{1}{2}$ gr. to be equivalent to two bottles of the liquid; and this was the quantity employed in experiment 11. In that experiment the animal survived 28 hours and a half: in the present, death did not take place until nearly four days had elapsed; but the quantity in this last case was given in three doses instead of two, as in the first; and therefore we are not led to the conclusion, that any of the virulence of the medicine was dissipated in the careful evaporation of it. The peculiar odour of the eau medicinale comes over by distillation. With regard to the symptoms in this experiment, it may be noticed, that the pulse was rendered quicker by the repeated doses of the medicine. The reverse happened in experiment 11. Is this to be imputed to the difference of constitution in different dogs? Or is it probable that the eau medicinale is a compound preparation, and that its more sedative part becomes dissipated by evaporation? The morbid changes of structure were here rather more remarkable than in experiment 11. The insulated state of the fibrine in the aorta, moulded by the vessel, and which, occurring in the human subject, was formerly called a polypus of the vessel, seems connected with the laboured death of the animal; and appears to demonstrate the powerful influence of the medicine upon the circulation.

The intromusception of the intestine, marks its severe action upon the muscular function of the bowels, together with its destructive effects upon the mucous membrane.

EXP. 22.—The animal recovered in a short time from the administration of twenty grains of this extract. It is true that the immediate symptoms were urgent, but we again see the proof of the superior activity of the *cau medicinale*.

EXP. 23.—I was desirous in this experiment to compare the effects of hellebore, elaterium, and colchicum, in the solid form. The hellebore rendered the stomach extremely irritable. Less of sedative effect was produced on the pulse than in experiment 16, in which opium was joined with the hellebore. It is evident, from all the experiments with hellebore, that the dog, although seriously disordered by it, can bear very large doses without fatal consequence.

The dog was also able to take a very large dose of elaterium without death being produced. Indeed the recovery was so quick that it seems necessary to suppose that the medicine was rejected from the stomach. The influence upon the pulse was very different from that of hellebore, the elaterium producing excessive frequency. This happened also in experiment 17. The destructive properties of colchicum are shewn in the latter part of this experiment. Ten grains of extract, obtained from a vinous preparation, given in two doses, proved fatal in forty-eight hours. That this smaller quantity than what was used in experiment 22, should kill the animal, is to be considered in connection with the previous administration of two powerful medicines, which must have ren-

dered the animal more susceptible of injury, independent of any mischief which they might have produced.

EXP. 24 and 25.—These experiments afforded a further and still more convincing demonstration in how very favourable a manner the acetic acid modifies the properties of the colchicum. The kidneys were excited to increased action in a remarkable degree. The mucous membrane of the intestines was only slightly irritated, if we compare its effects with those produced by the other preparations of colchicum; and this effect is more clearly shewn in the result of the following experiment.

EXP. 26.—We have here further to observe, in how very different a manner the colchicum acts, according to the menstruum which is employed in its preparation. About seven grains of this extract produced very severe symptoms, and a second dose of four grains almost proved fatal. Indeed symptoms of almost equal severity followed the use of this substance, as from a similar quantity of extract obtained from the eau medicinale. The first dose of the eau medicinale extract rendered the pulse slower; the opposite effect was produced by each dose of the colchicum extract. The animal recovered very quickly from this experiment, when we consider the intensity of the symptoms which he suffered.

EXP. 27.—This experiment points out that the colchicum is very quickly absorbed into the circulation; and the rapid progress of the symptoms demonstrates the virulence of the medicine. This is further shewn by the local action of the extract. We may here again notice the immediate influence of colchicum in quickening the pulse. The animal survived the experiment only twelve

hours. The morbid changes discovered on dissection were similar to those produced by injection of the medicine into the circulation, and its administration by the mouth.

EXP. 28.—Sir Everard Home having found, as I have stated at p. 526, that the eau medicinale was a much more active medicine when used with its sediment, than if poured off clear, and considering the vinum colchici to be substantially the same medicine as the eau medicinale, he was induced to make trial of the sediment of the colchicum administered in a separate state. He relates, *Phil. Trans.* Part II. that “six grains of the deposit of the vinous infusion of colchicum were given to a dog in bread and milk; in three hours it produced vomiting and purging, which lasted twenty-four hours; during the latter part of that time there was blood in the stools, as well as in what was brought up from the stomach.” The result of this experiment led Sir Everard to recommend the use of the vinous infusion of colchicum in its transparent state, after having by degrees deposited its sediment. The latest observations of Sir Everard are contained in a letter to Mr. Brande, published in No. XI. of the *Journal of Science and the Arts*; and he states, that the deposit which is separated from the infusion by keeping, when given by itself in the dose of a few grains, produced inflammation and ulceration on the coats of the stomach and intestines;” and “that the infusion removes the paroxysm of gout equally readily, whether given without the deposit as with it.” Being desirous to ascertain to what extent the deposit might be administered to the dog, I carefully collected, from genuine preparations, a large quantity of that sediment which the vinous infusion of colchicum very slowly deposits after its last filtration, and being put by for use in the transparent state. My surprise may be

conceived on finding, that even forty grains* were administered in four doses, without any appearance of inconvenient effects to the animal.

I am wholly at a loss to account for the very different result of my experiments; for I must necessarily conclude that the deposit of the vinous infusion of colchicum is an inert substance. It appears to me to consist chiefly of insoluble extractive matter. It cannot be dissolved either in wine or diluted spirit, even with the assistance of heat. On the contrary, the sediment of the *can medicinale* is perfectly soluble in its own liquor, by the application of heat. Hence a very striking difference is manifested in the nature of these two medicines.

EXP. 29.—My intention in this and the three following Experiments, was to ascertain the comparative degree in which the respective menstrua had extracted the properties of the colchicum. In the formula directed by Sir Everard Home, the alcohol of the wine is first carried off by distillation, and therefore the remaining fluid may be stated to be, little else than mere water, holding in solution portions of saline and vegetable matter. This I conceive to be a very unfavourable menstruum, as from the quantity of mucilage of the colchicum taken up by the water, it is almost impossible to obtain a clear liquor. On distilling some of the infusion procured from Fisher's, a considerable portion of spirit was obtained; so that I presume it has been found expedient to deviate from the

* This sediment was freed from its moisture by being placed in the cavity of a piece of chalk-stone (no heat being used); and the dose expressed was of dry sediment. Hence it is readily seen that the quantity administered was very large. It was given in mixture with a little water.

formula in question, either in using wine itself, or in adding some alcohol to the aqueous liquor. It is demonstrated by the present experiment, that the menstruum in question had only partially extracted the active properties of the colchicum. I think, however, that this is chiefly to be imputed to the very large proportion of roots employed; no less, namely, than 2 pounds to 24 oz. of menstruum. The animal was affected with very severe symptoms, and died in thirty-six hours.

EXP. 30.—The result of this experiment, with a vinous infusion of a much smaller proportion of solid material, confirms the inference just now stated. More symptoms, however, arose than might be expected; but the animal quickly recovered.

EXP. 31.—This experiment points out that the proof spirit is a very successful menstruum in depriving the colchicum of all its active properties; but I have before* offered my objections to its employment as a medicine.

* See p. 194, 198, and Exp. 8, 26, 27. At p. 219, I have erroneously described the pure alcohol menstruum as dissolving most of the principles of the colchicum; for in a subsequent Exp. with proof spirit, I found that 100 parts yielded six grains of extract, proving that it is by far the most active of all the menstrua which can be employed. I may here take occasion to mention that one of the fluid preparations of colchicum which I employed proved almost wholly inert. I obtained it from a very respectable chemist. On examining the roots which had been used, I found that, instead of being firm and mealy, they were very porous and fibrous. Hence a lesson is afforded to the chemist, of the necessity of examining the condition of the roots before they are employed for medicinal purpose.

EXP. 32.—The large proportions in which the acetie preparation can be used with impunity, as shewn in Exp. 19, 24, and 25, would naturally suggest the question, Does the acetie acid really extract the virtues of colchicum; for otherwise little praise would be due to it for the mildness of its operation? The present experiment answers this question most satisfactorily; and appears to me to lend a full support to the preference which I have assigned to the acetum colchici, over every other preparation of this medicine.

EXP. 33.—The result of Exp. 28 having been so directly opposite to what I expected, I was desirous to ascertain the comparative activity of clear and turbid portions of the infusion of colchicum, prepared according to the formula of Sir Everard Home. It appears that 11 draehms of the clear infusion were administered in four doses, and that death took place in 88 hours. The appearances on dissection indicated very high inflammatory action.

EXP. 34.—This experiment serves to shew that the clear infusion possessed as much of the active properties of the colchicum, as the turbid. The terrier dog was even the strongest in appearance. In this experiment the animal survived a few hours longer than in the preceding instance. Hence, according to each of my experiments, it follows, that the deposit of the vinous infusion of colchicum is an inert substance; and, consequently, that the clear infusion is as active a medicine as that which is turbid from the admixture of the sediment. It is worthy of observation, how much weaker a preparation this infusion is, than the eau medicinale, Wilson's tincture, and Reynolds's specific; which, in the ratio of their strength,

may be ranked in the present order ; but notwithstanding this fact, we see its destructive influence upon the stomach and alimentary canal of the dog, in doses very far short of those of acetum colchici and magnesia administered with perfect impunity. I must add, that again and again I have prescribed the latter medicine with every benefit, to those patients who had previously taken the vinum colchici with ill effect and disappointment. I am quite aware that some gouty persons do take this medicine with much *apparent* advantage, but I contend that such instances form a strong exception to the general rule. I have already in this Treatise, detailed my objections to the *principle* of arresting gouty action, by a few *specific*-acting doses of this or any other medicine.

It appears to be the object of those who prescribe the gout *specifics*, to avoid producing any effect upon the secreting organs, and to treat the gout as a painful nervous irritation. The following is the rationale expressed by Sir Everard Home in the letter to which I have already referred. “ The infusion (of colchicum) when clear of the deposit, can be taken in doses of 60 or 70 minims without producing any disturbance in the stomach, increasing any of the secretions, or bringing on an irregularity of pulse ; effects which commonly occur when that dose is given with the deposit mixed with it.” On the contrary, it is professedly my object to remove vitiated secretions which are accumulated, by exciting the bowels and kidneys to increased action ; to allay the irritation of the nervous system, and relieve pain by sedatives ; further to mitigate the local gouty action, and preserve the limbs from lameness, by local treatment ; and, lastly, not to consider the disease as cured, when all the external symptoms have passed away ; or, to speak

more familiarly, when the fit of gout seems completely over; but to follow up an alterative method of treatment, both by a mild course of medicine, and attentive regimen, so that the constitution shall by degrees be radically changed.

EXP. 35.—In the administration of elaterium by the mouth, Exp. 17 and 23, so large a quantity was given without death being produced; and as in Exp. 6, by injection, laudanum was joined with it, I thought it desirable to make a distinct experiment by injection, which might unequivocally shew the influence of elaterium upon the dog. The morbid appearances found on dissection were not sufficient to account for death, which took place in the short space of three quarters of an hour. We ought therefore to conclude, with Orfila, who classes elaterium with the acrid poisons, “that death, which is the consequence of the administration of this substance, must be attributed to the disorder of the nervous system sympathetically affected*.” I have no doubt, that in Exp. 17 and 23, the animal rejected the medicine from the stomach. Orfila tied the œsophagus, in his experiments of administering elaterium by the mouth.

EXP. 36.—The sedative action of the resin of opium was much more remarkable than might be expected, and serves to shew that the anodyne properties of opium must be in some degree weakened by the separation of its resin, although its stimulating qualities may thus be lessened. The dose of sedative extract was in the proportion of a fourth to the resin, but it acted in a very similar manner and degree. The dose of 15 grains of crude opium had

* Wallis's Translation, vol. ii. p. 16.

scarcely any effect, except that of rendering the pulse slower. One drachm produced powerful effects on the nervous system: but we see that from this large dose, in addition to what had been before administered, the animal shortly recovered. How well does this experiment answer the objections of those who may be disposed to doubt the instruction afforded by the present experiments with the gout medicines, as applicable to the administration of them to the human subject. The dog can take, without fatal injury, very large doses of those powerful agents, hellebore, elaterium, opium, and digitalis; whereas, of the several gout medicines, the repeated portions employed in the present experiments were such as are directed for the human subject, or nearly so; and we witness the destructive effects produced.

In regard to the probable nature of the eau medicinale, it appears to me rendered sufficiently evident that it cannot possibly be a preparation from the bulbs of colchicum. Orfila makes mention of the active qualities of the seeds of colchicum. I shall take a convenient opportunity to ascertain whether a preparation can be made from the seeds, agreeing more with all the properties of the eau medicinale, than the medicines which have been made the subject of these experiments.

In addition to what I have mentioned of Wilson's tincture, at page 217, I have to observe, that it appears, by its action on the dog, and from other circumstances, to be a concentrated preparation of colchicum. It resembles a mixture of the extract obtained by evaporation with diluted spirit, and does not become transparent on being heated.

Reynolds's tincture appears, from good evidence, to be a preparation of colchicum in rum, with colouring matter, more diluted than Wilson's tincture, but most

probably a concentrated warm infusion. Its sediment, like that of colchicum in the other preparations, is insoluble.

I shall here bring to a conclusion the details of this experimental inquiry, which I have endeavoured to make as demonstrative as the nature of the subject would allow; and which, I trust, will be found to contain a considerable share of useful information.

SOME
GENERAL CONSIDERATIONS
ON
MORBID STATES OF THE DIGESTIVE ORGANS;
WITH
PRACTICAL OBSERVATIONS ON GRAVEL.

WHEN we consider the importance of the functions which each of the digestive organs respectively performs, from the first step of the process to the last in the office of nutrition; and how constantly we exceed our natural wants, and commit every sort of indiscretion in our daily habits of diet; it cannot, on the least reflection, be matter of surprise that some disorder of function should be a frequent consequence. So much indeed are the derangements of the digestive organs connected with all chronic diseases, that if a division of them were to be founded on the most successful mode of treatment, the labours of the Nosologist would be reduced within a small compass. The confidence in specific remedies for particular diseases is gradually dying away, as sound science advances; and of late years, true philosophy has begun to sway the practice of medicine, in making careful experiment and rational induction the basis of our theory, instead of vague hypothesis and popular prejudice.

The important consideration which is due to the state of the digestive organs in the treatment of various diseases of serious magnitude, is exemplified in the most interesting manner by Dr. Hamilton and Mr. Abernethy, in their respective publications on the subject; and the

original views which they introduced must, I conceive, be ranked amongst the greatest improvements which the art of practical medicine has received. We no longer superficially allow of the application of tonics and nervous medicines for a weakened stomach and its attendant nervous sympathies, or for nervous forms of complaint; but we make it our care strictly to investigate the causes of such debility. Obstructions may exist below the stomach, and Nature may prudently have denied an appetite, when the powers of digestion are oppressed and interrupted. The nervous sympathies which attend these derangements, are now justly regarded as secondary forms of complaint, and are often found to yield, as by a charm, to the removal of the primary causes of irritation. The general doctrines of disease having thus received a legitimate improvement, it might be imagined that the consequent practice of medicine in individual diseases, and anomalous cases, would become equally clear and defined. Some improvement, however, in the laying down of our principle appears to me to be yet wanting. Rules of exact treatment, it is true, can only belong to experience, and do not admit of graphical delineation; but I cannot help believing that the principle of analysis upon which I have founded the preceding Treatise, may also be brought into useful application on my present subject. A great difference of opinion appears to prevail, both as to the particular viscus most commonly affected, and also as to the ratio medendi. By some, whatever organ or organs in the series may be affected, calomel is administered with the same freedom and frequency as the most simple purgative in the *materia medica*; and, doubtless, serious and irreparable mischief is often inflicted on the constitution by the abuse of this powerful agent.

Even simplicity, which to a certain extent belongs equally to philosophy and truth, may be carried too far, and lead us too much into a principle of generalisation. Dr. Hamilton's peculiar practice appears to consist in the continued employment of purgative medicines; so that not only a vitiated accumulation in the alimentary canal is prevented by the constant excretions which take place; but healthy action becomes excited by the total change of function which is thus produced. This, and not merely the effect of the removal of any particular accumulation, appears to be the necessary explanation of the mode of cure. In the space of time occupied in the treatment, the quantity of mere accumulation cannot bear any proportion to the ratio of total effect which is produced by the purgative plan.

Mr. Abernethy's method of treatment seems to be founded on the belief that simple error of secretion in the course of the alimentary canal, but in the liver more particularly, is the cause of the symptoms; and and that this is to be obviated by means of treatment which tend rather to sooth and correct, than to act with any considerable degree of activity in promoting excretion. In the one case, a dose of calomel in combination with a purgative is chosen as the remedy; in the other, a milder oxide of mercury, and in a small dose, so as professedly not to produce any very sensible operation; and the action of the bowels is to be gently assisted by some very mild aperient.

The diseases treated of by these different authors are certainly very different in character. Dr. Hamilton brings under consideration a variety of important diseases, acute and chronic, successfully treated by purgative medicines. Mr. Abernethy has, in a most ingenious and philosophical Essay, treated of those general derangements of the

system, which have appeared to owe their origin and support to a faulty state of the digestive organs, and are almost non-descript in their exact character. As it is well known, he was led into his examination of the subject, by finding, in his practice as a surgeon, that local diseases often became evidently aggravated and extremely untractable from the disordered condition of the constitution. The practice of these eminent authors, as just observed, relates to diseases of a different class; but it also differs considerably, as it appears to me, in principle. For the sake of distinction, I shall call the respective methods, purgative, and alterative.

I am not prepared to enter upon the arduous task of pointing out all the respective indications, which may call for one or the other mode of practice; but I conceive that an arrangement of this sort is susceptible of some useful rules, and these I shall briefly attempt to offer. Not only is our principle of treatment to be varied in correspondence with the individual morbid diathesis; accordingly as the stomach, the liver, the spleen, the kidneys, a particular part or the whole of the alimentary canal may be affected, and whether separately or together; and further, according to the kind and degree of symptoms, the modifying influence of difference of constitution, and the nature of the constitutional disease under which the patient may accidentally labour; but also, the particular remedy chosen from the same class of medicines must be adapted to the particular intention, or set of intentions, which we wish to fulfil. This, indeed, is an extensive subject, and would demand a volume rather than a few pages for its discussion*.

* For much valuable information "On various Diseases of the Abdominal Viscera," see Dr. Pemberton's well-known Practical Treatise.

When the constitution has a tendency to gout in connexion with a morbid state of the digestive functions, as I have already stated, diuretic aperients taken in moderate yet regular doses, together with the mild alterative use of mercury, usually constitute the most efficacious treatment; because, in addition to a faulty condition of secreting functions, there is a strong disposition to repletion and plethora, which is to be guarded against. Also, where the gouty disposition does not exist, if apoplexy or severe inflammatory erysipelas be the threatening disease; or if our patient be corpulent and of a full habit, and is affected with anomalous symptoms bordering on paralytic tendency; the same united principle of treatment is to be kept in view.

The modification of practice, according to the particular organ affected, is now to be considered.

When the stomach is the most affected part, as indicated by the symptoms described at p. 80, very decided relief will in many cases be afforded by an emetic of ipecacuanha; and the prompt removal of the offending contents of the stomach will often afford more immediate and decided advantage, than the continued use of a mere corrective. If mercury be administered otherwise than as a purgative, it is admissible only in very small doses, and precisely according to the rules of Mr. Abernethy. If, even with such doses, the fur of the tongue increase, the saliva become more viscid, and foetor of the breath take place, it should be immediately abandoned. Eructations, or flatus of the stomach, constitute the most distressing symptom of dyspepsia*, and are the immediate

* I lately saw a gentleman, who had for a long time suffered, in a distressing degree, from dyspepsia. Flatus, and an irregular action of the bowels, were his most troublesome symp-

effect of fermentation in the stomach, which arises from the incompetence of its powers to produce the first changes in the food after mastication. The use of tonics is certainly indicated ; but they must be managed with great caution. Strong bitters irritate and disagree. The neutral carbonate of ammonia alone, in warm water or any medicated vehicle, or with the addition of lemon juice, so as to be taken in a state of effervescence, is an useful remedy. If this salt, however, be too stimulating, the carbonate of soda should be preferred ; and either may be given in conjunction with infusion of cascarilla or columbo, and a small portion of aromatic tincture. Care should be taken not to produce nausea by any kind of medicine ; and the bowels should for the most part be regulated by pills, rather than by any liquid aperient. The wasted state of body and debility which are consequent to urgent dyspepsia, must be treated cautiously with respect to purgatives, which, if improperly employed, carry off too much of the chyle, that is but scantily prepared. Thus saline purgatives, which operate on the whole tract of the alimentary canal, are less suitable than purgatives of slow solution in the form of pills. For this reason I am persuaded that many dyspeptic persons are injured by a course, and especially if long continued, of Cheltenham water. For the state of stomach now under consideration, proper preparation being made, and in the

toms. Upon investigating the case, I discovered the presence of an inguinal hernia, and of which he had not been in the least aware. He derived such immediate benefit from the use of a truss, and so much improvement took place in the action of the bowels, that he almost derived his cure from the use of the instrument ; the regular continuity of the canal being thus restored.

fit season of the year, a course of the pure chalybeate water of Tunbridge Wells exerts the most beneficial effects*. Again and again have I seen the dyspeptic invalid receive renovated powers from the use of this water, when proper regimen has also been observed.

When the symptoms correspond with the description at p. 81, the stomach is only secondarily affected by the indigestion which prevails in the alimentary canal, and mostly in the duodenum. It is to be kept in view, that in this case the appetite is not deficient; and that the patient takes every day an excess of food, in proportion to the energy of the digestive and assimilating process. Here, I apprehend, we should blend the practice of Dr. Hamilton and Mr. Abernethy. It is advantageous to stimulate the bowels occasionally to full action by a mercurial purgative; and even with the alterative doses of a mercurial preparation, it often succeeds better to join with them a small portion of some purgative medicine,

* I may take the present occasion to observe, that in a very wet summer, as, for example, in that of 1816, the impregnation of this spring is very much weakened, although it endeavours to make up in quantity of supply in a given time, what it wants in actual strength. Thus, in the beginning of November 1815, after an unusually dry summer, the spring yielded only one quart in a minute. In October 1816, after a singularly wet season, the supply in a minute was no less than three gallons and a half. Its impregnation was proportionably weakened. I find, that by comparing the effect of re-agents with the water, both as to the time and degree in which they act, with the results from the same re-agents at the time when I made my analysis, I can form a very good estimate of the strength of the water at any particular time. This is convenient, as pointing out whether or not some pharmaceutical preparation of steel should be added to the use of the water.

as the pulvis alöes compositus et pulvis rhei, with sapo durus, than to administer them separately; because it is our particular object to stimulate the bowels, rather than to favour mercurial absorption. A draught, such as is described at p. 265, or 266, or p. 398, in conjunction with either description of pill, or that suggested by Mr. Abernethy, with the infusions of senna and of gentian and some aromatic tincture, will be found advantageous. The three first-mentioned forms of medicine will probably deserve the preference, when acidity and flatus prevail, and especially when the stomach is in a feeble state; but for the removal of habitual costiveness, the last description of draught, when joined with the use of the alterative pill, is often remarkably successful. If, with a debilitated state of stomach, the tongue has a pale appearance and a cheesy sort of fur, with a state of gum which looks spongy, and is familiarly known by the name of scorbutic, the administration of nitric acid diffused in the cascarilla or columbo infusion, is much to be recommended, and, for this condition of dyspepsia, may prove a better tonic than the alkaline bitter. In such cases it will be questionable whether the mercurial alteratives should be employed; not on account of the use of the nitric acid, for when the two are taken at distinct intervals, I have seldom seen any disagreement produced; but because mercury, in the mildest form, usually proves unfriendly to the stomach of a patient of this description.

In the case to which I have alluded at p. 86, I found the treatment which I have just now described, after due perseverance, and with the union of exact regimen, succeed perfectly to my satisfaction. Nervous medicines, as intended to apply to the state of hypochondriasis, from which the patient suffered in a most distressing degree; and steel tonics with Peruvian bark, directed in reference

to the debility and languor of the stomach and the whole system, which also prevailed, had been previously tried for many weeks. From such injudicious practice, slight palliation only, by means of the volatile stimulants, was afforded; and an aggravation and confirmation of all the symptoms had resulted from the misapplied tonics.

In the treatment of the symptoms enumerated at p. 89, we are to consider that the liver itself is the chief seat of disease. It is obstructed, and perhaps affected with some degree of chronic inflammation. Here then we find that a mercurial course of medicine, with the regulated daily action of some purgative, properly adapted in its activity to the particular case, proves of the most manifest advantage. In many cases, and particularly when we can distinguish the seat of disease, either from the tenderness of the part on pressure, or from any enlargement, the use of mercury, principally by inunction on the affected part, will be found more efficacious than its internal use alone. The warm bath is an excellent auxiliary. Even in this form of disease, I am persuaded that we should avoid the severe employment of mercury, which, so administered, rather does mischief as a poison, than renders benefit as a supposed specific. When much pain prevails in either hypochondrium, the use of blisters, and sometimes also of cupping, should precede the inunction; and mercurial purgatives should be freely administered. In slighter degrees of uneasiness in the side, the emplastr. ammon. cum hydrarg. may afford considerable service.

The terms *bile* and *bilious* are, doubtless, too indiscriminately applied, both in popular language, and by the profession. If the patient suffer an attack of indigestion from ineaction in diet; or if he labour under dyspepsia, whether derived from continued irregularity

in regimen, or from the influence of other causes, the complaint receives the convenient name of bilious. When from such association of terms, the treatment is made to consist in a free use of calomel, it is obvious that a practical evil results from this error of language. If the seat of complaint be in the stomach alone, any active use of calomel should be a questionable point of treatment; and I may lay it down as a general rule of practice, that if the administration of mercury be at all required, it should be given only as an occasional purgative; or, if more repeated, it must be prescribed in its mildest alterative form.

There are two very opposite states of a faulty condition of liver, which I shall briefly point out. In the one, the biliary vessels are surcharged; and the bile, not passing off with adequate freedom by the intestinal canal, is in part absorbed into the circulation, and produces more or less of jaundice. The discharges are almost invariably dark, and highly bilious in appearance. In very chronic cases of this description, the liver acquires an increase of bulk, and may be, in many instances, discovered by manual examination, projecting below the ribs in the right hypochondrium, and considerable tenderness is almost with certainty felt at the ensiform cartilage. In the examples to which I have alluded at pages 91 and 94, I had full opportunity to study this species of hepatic complaint. I considered that from the misapplication of tonic medicines, which had been freely given, and the neglect of purgatives and alteratives, these patients had, for a long time, been suffering the serious evil of fixing their disorder. The nervous symptoms which afflicted them had been palliated by nervous medicines and stimulating diet; whereas the correct and successful treatment in these cases consists in a judicious

employment of the aperient and alterative plan, and in a regulation of the regimen. By such means the circulation of the liver will most probably be rendered free; its healthy secreting functions will by degrees be restored, and the nervous symptoms will vanish; thus affording a proof how entirely the symptoms were sympathetic, and of a secondary nature. I must state, however, that the occasional use of anti-spasmodic medicine is proper, and affords great relief to the urgent symptoms; but this constitutes only the palliative part of the treatment. In the case of symptomatic asthma, mentioned at p. 86, various tonic and antispasmodic medicines had been administered before I saw the patient, not only without relief, but with a material aggravation of the symptoms. The aperient and alterative course of treatment, together with regimen, completely effected a cure. In these cases, ass's milk proves a very useful adjunct.

The opposite description of case is that in which the liver is remarkably torpid in its action, as is shewn by the inert state of the bowels, and the absence of bile in the alvine discharges. The complexion is sallow; the appetite is deficient; the spirits are depressed. As comprising a course of medicine, my experience leads me to think more favourably of *pilula hydrarg.* joined with an aromatic aperient, as the *pulvis aloës compos.* than of calomel; and as an aromatic tonic aperient, I would particularly recommend the *decoctum aloes compos.* which may be joined with other ingredients, as at p. 266, the gentian root or cascarilla bark being used, as the physician may prefer. Chlorine, of which I shall presently have occasion to speak, will particularly deserve a trial, should other medicines fail. Horse exercise and the air of the country, form powerful aids to the influence of medicine. In the list of remedies for the torpor of functions,

now under consideration, I may mention the agency of galvanism, as a remedy entitled to recommendation. I do not profess to have any considerable experience of its effects; but I should conceive that it would not be found equal in efficacy to the judicious use of medicine, except on those occasions in which, from the great want of energy in the organs of digestion, the action of medicines is found very imperfect and unsatisfactory. I am acquainted with some good results from its administration* under such circumstances; but also, in a case in which I hoped much from this energetic stimulant, I was wholly disappointed.

The discharge of blood from the hæmorrhoidal vessels, which takes place *ad sedem*, is very commonly connected with an unhealthy condition of the liver, and seems to arise immediately out of an obstruction to the returning circulation from the intestinal canal through the *vena portarum*. Within certain limits such a discharge usually proves a source of relief; but if it continue for a long course of time as a daily occurrence, it becomes a serious source of debility, and requires an attentive and distinct plan of treatment.

In considering the question of the particular viscus which has been the primary seat of complaint, when a complicated case of disorder, affecting the digestive organs and the nervous system, comes before us, I am induced to believe that sometimes we too much overlook the influence of the brain, and may fail in our treatment from confining our attention to the remote effects of disease. With whatever integrity the brain may be acting as an instrument of thought, its physical functions (if I may be al-

* It is but justice to M. La Beaume, of Southampton-Row, to observe, that he administers galvanism medically, with great skill and attention.

lowed the distinction) may be so disordered as to become the important source of serious secondary complaint. It is an organ, not only subject to physical causes of injury and disturbance, in common with the other viscera; but is, also, under the peculiar immediate influence of all our passions and emotions. It is highly probable, therefore, as a matter of theory, that the state of the circulation of the brain, and the condition of its nervous powers, should be frequently thrown into disorder, and have a powerful primary influence on the healthy action of the digestive organs. So far as the direct application of physical remedies to this organ is in our power, our chief assistance consists in those means of treatment which either increase or lessen the quantity of its blood, or which tend to make its circulation equal and regular. Much indeed is to be accomplished through the medium of the mind; and in this species of advice, the physician must at once become the philosopher and the philanthropist.

In the case of a severe injury of the head, we see immediately the important connexion which subsists between the brain and the digestive organs. They are thrown into disorder, and such is the series of the diseased actions, that they reflect their morbid influence upon the brain; and thus a mutual kind of disease becomes established. This is the point of treatment which Mr. Abernethy has so much and so usefully considered; but whether in the just importance he has assigned to the management of the digestive functions, in cases of local injury, he may not have recommended too exclusive a treatment, is a doubt which I myself am disposed to entertain.

In many most apposite and striking instances of disturbed action of the stomach, the liver, and the bowels, I have not succeeded in my treatment by purgatives and

alteratives, until the brain itself has been relieved from a state of congestion, and which had been rather obscurely shewn by the symptoms. Even in those cases, in which an obstructed state of the liver is the real cause of a morbid determination of the blood to the brain, at the same time that our radical treatment consists in removing such obstruction, it is important to give immediate relief to the vessels of the brain by the abstraction of blood. It must be our care to discriminate between the nervous symptoms which arise from opposite causes. Head-ache, vertigo, tinnitus aurium, pulsation, and a want of mental energy, may proceed either from an excess or a deficiency of blood in the vessels of the brain; and depletion or stimulus must form our consequent principle of practice.

Sydenham, in describing the influence of the state of the mind on the functions of the animal œconomy, observes, "The patient must likewise use his utmost endeavour to keep his mind easy; for all disquieting passions, if they once become immoderate, greatly dissolve the texture of the spirits, which are the instruments of digestion, and so of course increase the *gout*."

With respect now to the excreting function of the kidneys, if it be allowed, as the truth indeed requires, that it is a material agent in the process of healthy assimilation, it must follow that considerable importance belongs to a medical acquaintance with the various morbid changes which the urine undergoes; and that we should carefully study the true indications which its external characters present. The relative connexion which subsists between the digestive organs and the kidneys is of a very curious kind; and it seems surprising that the particular state of the one should so quickly influence the other, when we consider that the blood which goes to the

kidneys for the purpose of secretion, has first to pass in part the round of circulation. Also, as the office of the kidney is entirely excrementitious, how does it render this service to the whole of the circulating blood, when, in the first round of the circulation after the absorption of the chyle, it can only have acted on a very small part of the arterial blood of the system? I am disposed to conceive that several rounds of circulation take place before a perfect admixture of blood is effected; and that in this way only, the kidneys have the opportunity of performing their salutary action on the whole blood of the system. If a person be bled from the arm shortly after dinner, the serum is entirely milky, as I have often witnessed; and hence I derive a support to my argument.

In my previous discussion of the subject of the urine, I have dwelt at length on the nature of the pink and brick-dust sediment; and I shall now make reference to the observations which will be found at p. 96, 130, 156, &c.

In a congested state of the vena portarum system, I have always found in the urine an excess of urea, and of all the other principles; and can affirm, that whenever the pink or lateritious sediment appears copiously deposited, it may be taken as a fair presumptive proof of this fact; and may be viewed as a correct indication for the use of diuretic purgatives, and of a corrective plan of medicine and regimen. Also it will be found, in this case, that tonics can seldom be employed without disadvantage. When such sediment is copious and long continued, the conclusion follows which I have mentioned at p. 141; and the treatment just recommended must be pursued with great perseverance.

If the sediment consist of the earthy phosphates, as indicated by the distinctions described at p. 160, 162, and

163, &c. it usually happens that supporting methods of treatment are required, in conjunction with the purgative and corrective; for I have constantly observed, that the patient, under such circumstances, is suffering in some measure from debility and nervous irritation; and that, in the course of a paroxysm of gout, or of chronic inflammation of the liver unconnected with gout, the state of the nervous system gives rise to these symptoms more or less remarkably, whenever the composition of the sediment changes from the urates to the phosphates.

It is sometimes thought to be of importance, that we should be able to determine with accuracy whether the urine of a person under a disease, or disordered action of the liver, is containing bile; and also to form some estimate of its proportions. I therefore instituted some experiments, with a view to compare the delicacy of the muriatic acid as a test with that, very commonly chosen, of a linen rag. The following were the results.

Healthy recent bile added to healthy urine in the proportion of one part to 30, by muriatic acid was immediately rendered slightly green, and this effect was distinct in two or three minutes. After some time it became grass-green.

With bile and urine, one part to sixty, and muriatic acid, the green hue appeared distinct after two or three minutes.

When the proportions were of bile and urine, one part to one hundred and twenty, the green colour was not very distinguishable until after twenty-four hours.

Bile and water, one part to one hundred and twenty, instantly yielded the green hue to muriatic acid.

The proportions even being one to two hundred and forty, the change was apparent after long standing.

By means of the linen rag, the yellow stain distinctive

of bile was only just distinguishable when the proportion of bile to urine was one to sixty.

Some urine of a jaundiced person, which in a short time produced with muriatic acid a light grass green, did not communicate any yellow stain to linen after being steeped for several hours.

I have lastly to notice an opinion suggested by Mr. Rose*, referred to by Dr. Henry of Manchester as being a curious discovery, and supported by him with an additional example†, that, in hepatitis, the urine is devoid of its usual and very important constituent, *urea*. In the case of my gouty patient, mentioned at p. 297, the urine was almost destitute of urea‡; and, as I have stated at p. 299, the same fact appeared in six other examples; in all of which, however, the urine was also albuminous, and uric acid was equally deficient with urea. In these cases the stomach was weak, and the digestive powers were impaired; but I could not pronounce the disease to be hepatitis.

* See Thomson's Annals of Philosophy, June 1815.

† Ibid. November 1815.

‡ The method which I used was to concentrate the urine very considerably, and then, having filtered it, to add to respective portions, nitric acid, and a saturated solution of oxalic acid. The pearly crystals, which, when urea is present in much quantity, appear almost instantly, and form indeed an unequivocal evidence, are well known. The oxalic acid acts more slowly; and, if the urea be in minute proportion, requires about twenty-four hours for its decided effect. The precipitate consists of small transparent crystals, which, as they are viewed without being washed, appear reddish in colour. They are easily distinguished from the white pulverulent precipitate of lime and

In six very clear cases of hepatitis, however, I have not only detected urea by means of nitric acid, and oxalic acid, but have also in each of these examples found a great excess of this principle. I am compelled, therefore, to believe that the cases of Mr. Rose and Dr. Henry, ought rather to be held as exceptions from the general rule, than to be chosen as the ground of so strong a conclusion as they have formed.

In the medical management of all the forms of complaint which I have considered, the strictest rules of regimen must be observed; and without which, the skill of the physician will be abortive. I shall endeavour to include all these considerations under the head of prophylactic regimen. A modification of practice in the treatment of the derangements of the digestive organs, which is of great importance, arises from the combination which they form with other states of disease, and which may be either of a primary or secondary nature.

Abundant examples of such blended states of complaint, will be found in the authors to whom I have just alluded. I shall confine myself to some slight illustration; and probably I cannot choose any one more strik-

magneisa, which immediately falls down with this re-agent. From some comparative experiments, I have found this to be a test of still greater delicacy to detect urea, than the nitric acid; and I have also Dr. Prout's authority for the fact. The method by the distillation of the urine to procure carbonate of ammonia as the evidence of urea, and this product put to the test by its powers of decomposing muriate of lime, originally suggested by Vauquelin, and adopted as the most delicate by Dr. Henry, appears to me rather questionable. Will not other animal principles in the urine furnish carbonate of ammonia at the boiling temperature?

ing than the example of an irritable state of lungs, appearing in the form of troublesome cough; uneasy respiration, easily quickened by slight causes, and accompanied with tenderness of the chest or occasional pain in the side; which we occasionally see connected with more or less in number of the symptoms related from p. 83 to 87. Sometimes the patient makes the whole reference of the uneasy sensations which are felt, to the chest; and great care is required in our diagnosis. The epigastric region is usually affected with marked tenderness to pressure in these cases. The furred appearance of the tongue, the dyspeptic symptoms and peculiar state of the secretions, the absence of regular hectic and wasting, will be strong grounds of discrimination. Under such circumstances it is, that when our patient at first appears threatened with the quick and melancholy progress of *consumption*, the whole disease, after a patient course of treatment, will often happily yield to the co-operative plan of alteratives, mild tonics, and soothing sedatives, with such topical treatment as the case suggests. In urgent cases of this kind, the patient is said to have been cured of consumption; but the lungs have been only secondarily, and perhaps sympathetically affected.

When a scrophulous diathesis is mingled with the disordered digestive functions, it is incumbent upon us to use mercurial medicine with the utmost circumspection. If carelessly employed, it will operate as a most injurious poison; readily causing ptyalism, fœtor of the breath, universal disposition to glandular enlargement, general debility, and an aggravation of the dyspeptic symptoms. If we employ mercury in such cases of peculiar constitution, we are almost restricted to its use, merely as an occasional purgative. In some instances of this kind, I have procured material advantage to the stomach, and to the

system at large, from the use of *chlorine* as an internal medicine. I have given it in the form of a saturated aqueous solution* twice a day, in the dose of half a

* Prepared by Mr. Garden, operative and general chemist, No. 372, Oxford-street. It should be carefully kept secured from the air and light, to prevent its decomposition. I always direct that the vial may be covered over with dark paper.

Far be it from me to depreciate the skill and attention of other chemists; but it is due to Mr. Garden to state, that he is extremely successful in the preparation of chlorine, by the particular process which he uses. I requested Mr. Garden to ascertain, by careful experiment, the relative quantity of chlorine held in solution in water at given temperatures. He favoured me with the following results:

“Water, at the temperature of 50° Faht. may be made to absorb $2\frac{1}{2}$ times its volume.

“Water, at the temperature of 60° will absorb $1\frac{3}{4}$ times its volume.”

Hence appears the importance, and indeed the fact is obvious on the least consideration, that the medicine should be kept in a cool situation. The most favourable method is to place the vial, inverted, in a vessel of cold water, in a dark part of a cold cellar.

Dr. Willan (on Cutaneous Diseases, at p. 360) speaks of the use of this medicine in the early stage of scarlatina anginosa, in favorable terms. He describes the dose as half a drachm for an adult, and for children ten or twelve drops. I have not found it necessary to administer it in such small doses. He points out the care required to prevent the decomposition of the chlorine; and quotes from Mr. Allen the following test for its state of purity. “When litmus paper is plunged into the true oxymuriatic acid, it is deprived of colour; but if common muriatic be present, the paper will instantly receive a red tinge, and thus ascertains that the preparation is unfit for medical use.”

drachm, gradually increased to two drachms, in union with some simple vehicle. Diluted cinnamon water conveniently serves the purpose, and renders the chlorine sufficiently palatable. I must observe, that the chlorine readily enters into new combinations, and therefore is administered in its most pure and active form in water simply; and when the stomach accommodates itself to its taste and action, thus uncombined, it will sometimes be desirable so to administer it. In a weak and delicate state of stomach, however, I have often been much satisfied with its effects, united with a dose of tinct. cinchon. compos. and a wine glass of water. I have never prescribed the chlorine solution in a smaller dose than half a drachm, or in a larger dose than two drachms. I have administered it also in dyspepsia, and in the debility of the digestive organs which sometimes is consequent to an active course of mercurial medicine; and certainly with great good effect in most cases, and without injury in any.

With a bilious patient (I thus express myself for the sake of conciseness) it very commonly happens that tonic medicines disagree in a peculiar manner, confining the action of the bowels, and proving unfavourably stimulating. The chlorine here recommends itself very strongly, as it particularly favours the regular action of the bowels. Its solvent powers are remarkable. I digested a very compact biliary calculus in an aqueous solution of chlorine, both concentrated and much diluted, and found that by each it was almost equally acted upon, broken down into very minute fragments, and deprived of colour, in the course of twenty-four hours. No such remarkable effect can be produced by the acids or alkalies, either concentrated or dilute*. It freely dissolves calomel, converting it into

* Both nitric and sulphuric acids, if in a concentrated state, exert a considerable action upon the biliary calculus. The former

a chloride of mercury. Care must therefore be observed not to give it in conjunction with this medicine, except with a due interval of time. If it be administered immediately at the close of a mercurial course, its action in increasing the soreness of the gum, and renewing the ptyalism, is very quickly shewn.

To mention further its general agency, I have to observe that the stomach very readily accommodates itself to the medicine; and indeed I have not seen it produce sickness or nausea, except in two instances, in which the stomach was evidently in a previous state of disturbance. The patient usually experiences a sense of exhilaration and improved energy from its use. It commonly acts as a diuretic, but does not irritate the bowels, unless they are very prone to be disordered. I flatter myself with the hope that this medicine may be found a most useful auxiliary in the cure of *scrophula*; and from the good effects which I have already had occasion to witness, I can venture to recommend it to the notice of the profession as worthy of trial in this deplorable species of complaint; which so much sets at defiance all ordinary means of treatment.

To such general observations on this interesting sub-

effects a partial solution, which is of a pinkish colour; the latter acts more strongly, and produces a solution of a deep grass-green colour.

Nitro-muriatic acid, in a concentrated state, accomplishes a complete disintegration of the biliary calculus; and a considerable effect, though much more slowly, is produced by the same acid in a moderately diluted state.

Muriatic acid does not appear capable of acting upon this kind of calculus, at ordinary temperatures. The liquor potassæ, either concentrated or diluted, has only a slight effect.

ject, I must now confine myself; but, before its conclusion, I cannot refrain from taking a brief notice of the method of treatment by the nitro-muriatic acid bath or ablution, some time since introduced to our attention by Dr. Scott. Such is the difficulty of ascertaining the true character and value of any new remedy, that the investigation of its merits cannot be too general; and the communications of every practitioner, which are offered with care and candour, are entitled to be received as useful. In a science so obscure and difficult as that of practical medicine, both doctrines and remedies must pass the full ordeal of time and opposition, before they should be admitted as substantial, or dismissed as useless. It is under such impressions that I proceed to the examination of the present very popular remedy.

The earliest statements on the subject by Dr. Scott, which I have seen, appeared in the second number of the *Journal of Science and the Arts*. At p. 202, he observes, "When in India, I was most anxious to discover a substitute for the mercurial calces, less injurious and equally efficacious; and I have not been entirely without success. I knew that the nitric acid acts most readily on the resinous matter of the bile; and I was in hopes that I might communicate such an acidulous state to the living body as should produce the effects that I desired." He states, that through the medium of the stomach the operation of this acid, if given to the wished-for extent, had proved injurious, and that, from its absorption by the skin, some important effects had arisen. In the conclusion of the paper, he again speaks of the good results of charging the body with some of the mineral acids, or their elements, by means of the skin.

In Part III. of Mr. Charles Bell's *Surgical Observations*, Dr. Scott has annexed some remarks; among which

I find the following: "I am not now convinced that a particle of acid enters the system. The effects I suspect arise from chlorine alone." It is evident that the author himself has no fixed position as to the rationale of the remedy; and it might easily be shewn that the views which I have quoted are not very tenable. In the admixture of the acids with water in the first instance, chlorine is but sparingly formed. When the liquor is afterwards warmed for use, and exposed in an open vessel, the little chlorine contained must almost wholly escape. Also, if it were contained in abundance, it ought to be shewn that the absorbents are capable of taking it up from the surface. The facts, however, respecting the remedy, and not the theory, are the points most worthy of discussion, and to these I now shall proceed.

I may first remark, that we have seldom any difficulty in the internal administration either of the acids or of chlorine. A question then naturally arises—is this apparently inefficacious mode of applying a medicine, which we can, with facility, use internally, really attended with any superior effects? And what is the nature and amount of such effects? Does the remedy certainly possess an active power; and is it worthy of confidence, either by itself, or as an auxiliary to other treatment?

Of the effects of any remedy of mysterious agency, the sentiments of the patient must be received with great distrust. He usually lends his faith to its operation, from a prepossession in its favour; and the influence of a mental impression frequently stands in the place of a positive and palpable effect. Of this truth, in the history of a thousand boasted medicines, of the metallic tractors, and of animal magnetism, we have abundant illustration.

I have made trial of this remedy, both by immersion of the feet, and by ablution of the whole of the lower ex-

tremities, in several cases which appeared to me the most favourably adapted to receive all the good effects which it should be capable of affording. The proportions which I employed were one drachm of each acid to a quart of water sufficiently warm.

One lady, suffering from dyspepsia and deficient secretion of bile, but easily influenced by mercurial medicine, in having the action of the liver increased; of very irritable constitution, and with a thin and delicate texture of skin; sponged the legs and thighs very diligently for ten nights, but no apparent effect of any kind was produced.

A gentleman, whose liver had been diseased when in India, and who has since frequently suffered manifest derangement of this organ; such as degrees of jaundice, some pain and tenderness in the right hypochondrium, and a vitiated condition of the several secretions; when under the influence of such complaint in its most chronic form, gave trial to this treatment, and pursued it very steadily every night for five weeks. He was not sensible of the least influence from it, either on the bowels or kidneys. He thought that shortly after commencing its use the uneasiness of his side was increased; but at the expiration of the above period, he could not trace any positive effects, or consider that he had derived from it any advantage. He acknowledges with full approbation the decided good effects of a plan of medicine, consisting of purgatives and the mild alterative use of mercury, which he has since pursued.

A lady of the most irritable constitution, highly susceptible to very gentle doses of mercurial medicine, or indeed to the agency of any active remedies, immersed her feet for ten nights, without the least apparent operation, except that the coldness of feet, to which she was

much subject, was comfortably removed during the period of its employment.

Another lady, whose liver was evidently enlarged and in a state of bilious congestion, always readily and beneficially acted upon by mercurial purgatives, immersed the feet and sponged the limbs for twelve nights; but, with the exception that it warmed the extremities very comfortably, it did not appear to produce any effect; and she found it necessary, as usual, to resort to her aperient medicines to excite the bowels.

Lastly, I made trial of the remedy on myself, being at the time, as it is in common language expressed, slightly bilious. I found it produce in the extremities an agreeable sensation of warmth; and this was the more perceptible, as I habitually suffer inconvenience in this respect. Slight pricking irritation also followed. I continued the ablution six nights. Beyond this local action I could not trace any effect; and since, the occasional coldness of the extremities has returned. I examined the specific gravity of the urine, and also its general characters every morning, and could not detect in it any variation from the ordinary state; comparing the results with former examinations. Neither the action of the bowels, my appetite, nor my internal sensations, were in any degree affected.

It appeared to me, therefore, that it acted principally as a stimulating lotion, which by its penetrating qualities might excite the vessels of the skin to increased action; and as dyspeptic persons, and those who suffer from defect or irregularity of the bilious secretion, are usually affected with coldness of the extremities, I conceived this might be a mode in which it would do service. It has been suggested that its influence on the sentient extremities of the nerves may be similar to that of

galvanism, and that such may be the *modus operandi*. This hypothesis is ingenious, and seems entitled to some consideration.

From the close sympathy existing between the skin and the alimentary canal, we can readily believe that some individuals who have a thin skin, and who possess this sympathy in an exquisite degree, will find the action of the bowels suddenly excited by the free application of acid matter to the skin, and especially when they are in a state very susceptible to irritation. In this way the excretions of the liver, in common with the general action produced, would be increased; but such reasoning differs very much from that of assigning to the remedy the direct, and as it were specific, power of acting upon the liver. Dr. Scott observes (Bell's Reports, p. 363), "With people disposed to bile, it is necessary to keep the bowels very open during the use of the bath; for one of its effects, as I have said, and on which much of its beneficial tendency depends, is to produce a flow of bile into the intestinal canal." In opposition to my unsuccessful experience with this remedy, I hear repeatedly, from very respectable authority, the praises of its extraordinary powers. I cannot therefore presume to deny that the remedy may sometimes be one of considerable efficacy, although I have with very careful search after truth endeavoured in vain to procure useful effects from it. I admit that my experience with it has not been extensive, and therefore do not offer it as conclusive. I cannot help at the same time remarking, that when this remedy is brought forward in apparent opposition to the usual method of practice by mercury, we should proceed very carefully in our estimate of its powers; and it would be matter of regret that an active agent in medicine, and one which with *proper* use is really successful, should fall

into unmerited odium from the inferior if not unworthy pretensions of the remedy now in question, which is certainly one very troublesome in its application, both as to the mode of its employment, and the time required, as stated by the author, for its use.

Mr. Bell, with much panegyric "on the obvious constitutional effects arising from this bath," as he conceives, in ordinary disorders of the constitution, has offered an account of its use "in certain obscure cases of syphilis," which appears to me highly deserving of investigation. Introducing the subject, he speaks of this description of patient as being "half poisoned with mercury," and still suffering from his disorder or its sequelæ; and under such circumstances he appears to offer the present treatment chiefly as a substitute for a well-regulated course of mercury, or for sarsaparilla, milk diet, and country air. In referring to the operation of the remedy, he says, "We have seen salivation and sore gums produced in two instances." On reading this statement, the idea occurred to me, that, when the acid liquor is freely and repeatedly applied to the skin of a patient whose system is already highly charged with mercury, a chemical action would be produced on the surface; and hence, an active combination of the acid and vapourised metal (chloride of mercury) might be supposed to result. If this view be just, it would afford an explanation of the *modus operandi* of this agent in such cases; so that it might, on many occasions, become desirable to employ the acid bath in conjunction with a mercurial course; using less mercury on account of such combination; or to abridge the duration of the mercurial course, and follow it with the use of the bath. As far as the application of the bath is concerned, I would in any case advise the application of it for about five or ten minutes, twice or thrice a day, so as thoroughly to

saturate the skin; in preference to the tedious period of half an hour, or an hour, which Dr. Scott recommends.

The following opportunity lately presented itself to me of putting the idea, which I have just now stated, to the test.

A young man, who had been pursuing a regular mercurial course, with the effect of removing his primary symptoms, applied to me with a cutaneous eruption, which appeared to be the syphilitic psoriasis. His gum was very sore and foul, his tongue was furred, and the salivary glands were in considerable action. I directed such means as I judged the most calculated to restore the natural appearance of the gum. In about a week, this change being sufficiently accomplished, and when the silver watch which he wore was still entirely tarnished, I directed the acid bath and sponging, night and morning for ten minutes. At his next visit, three days after, the gums were very much swollen, of a luxuriant red, and the salivation was considerable. He was much better both in his feelings, and in appearance. Pursuing the treatment, the eruption died away, leaving the skin scaly. After about eight days of the bath, the gum began to mend, although the acid remedy was still continued. No increased effect took place either on the kidneys, or the bowels, in this case.

This was the report which I formerly offered respecting the nitro-muriatic acid bath of Dr. Scott. I have since followed up my investigation of the remedy, both by inquiry of those professional friends whose judgment I most esteem, and by watching its effects in the occasional instances in which patients have wished for a trial of it. From the profession I have received a varied testimony, but certainly the decision against the claims of the bath as a certain remedy; I must even add, to any positive

effect which can be held important—has greatly predominated. A few, whose opinion I must consider as very respectable authority, certainly do continue to praise the remedy as often effecting that for the patient which is intended by mercury, with the additional advantage of being restorative to the constitution. For myself, I must confess that I have not seen sufficient occasion to alter my former general opinion on the subject. I have again witnessed a fair trial of the bath in many favourable examples for its operation, and have encouraged the faith of the patient in the expectation of benefit. I have used the proportions which Dr. Scott prefers in the following admixture. Add to 16 oz. of water, separately, 10 oz. of muriatic acid, and 6 oz. of nitric acid. The usual degree of dilution which proves convenient to the skin of the patient, is, two ounces of this liquid to a gallon of warm water. The bath is conveniently renewed by throwing away a fourth, and adding the necessary measure of acids and boiling water, so that the whole liquor is immediately brought to the necessary comfortable temperature. With two young children, of thin skin and delicate constitution, indisposed with diarrhœa, unhealthy secretions in the bowels, deficiency of appetite, and loss of flesh, the bath has, under my own observation, appeared to do real service, in correcting the error which I have stated, and in improving the constitution.

I have repeated the experiment just now stated, upon a patient strongly under the influence of mercury, and with a similar result and very good effect; so that upon the whole I have to offer the same sentiments regarding this remedy, which I before ventured to deliver.

ON THE
THEORY AND TREATMENT
OF
GRAVEL.

FROM p. 98 to 101, and at p. 160 to 166, &c. I have considered the subject of gravel with reference to the external characters and chemical composition of the different kinds of sediment; and it now remains that I should offer some practical observations on the rationale and treatment of this distressing complaint.

With respect to a large proportion of gouty persons, as I have before mentioned, it is a curious fact, that, in the history which they give of their constitution, they state that before the invasion of the gout, they were remarkably subject to the gravel; but that since the gout has supervened, they have been only slightly affected with that disorder. Many of my gouty patients relate that they suffered in a very severe manner at an early period of life from the gravel. The gout becoming their disease, the urine for the most part ceases to furnish the crystallized sediment which we denominate the gravel; but abundantly supplies that which is pulverulent in its form, and of the pink or brick-dust colour in appearance. The explanation of this phenomenon is, in my opinion, to be sought for in the altered state of the sanguiferous system. I have already stated at p. 160, that the two sediments agree very much in chemical composition, and have suggested the idea that the excess of animal matter and mucus, which

accompanies the pink and lateritious deposit, prevents the uric acid from assuming a crystalline form; for in however abundant a quantity this kind of sediment may be found, it is nevertheless completely soluble in the urine at its natural temperature in the bladder; even though a much larger portion of uric acid may in this manner be secreted, than in the urine which deposits crystals. Dr. Prout remarks on this point; "This appears to be a wise provision of Nature; for if the uric acid were secreted in a pure state so abundantly, as the animal œconomy under certain circumstances seems to require, it would not be held in solution in the urine." This comparison of the change of constitution in the gouty patient is exactly in point, in considering the pathology of gravel.

That the one disease should apparently pass into the other at different periods of life, agrees with the general idea that gout and gravel have a very close connexion; and most authors who have written upon the one disease, have also treated of the other. The analogy has been further supported by the well-known fact, that the uric acid is invariably the chief constituent of the gouty calculi, and most commonly of the urinary; and because it enters almost entirely into the composition of the red gravel, as it is called, and in a great degree of the coloured sediments already mentioned. Notwithstanding these close links of connexion, however, we must see an important difference in the doctrine which is due to the two diseases. The prevailing theory on the subject represents that gravel is a disease arising from an excessive secretion of uric acid; but, as I have mentioned at p. 117, the appearance of the crystals is not to be received as a proof of such a fact; although this conclusion is certainly just, in the case of a considerable deposit of the pink or lateritious sediment. In the example of the crystallized

deposit, I conceive that a new combination of the uric acid takes place, rather than an increase of its proportions; and my grounds for this conclusion I have already stated. The influence of an excess of acid matter in the alimentary canal, appears to be the true exciting cause of this change of action in the kidney; or in other words, of that new arrangement of the elements of the urine, out of which the crystals called the gravel are produced. The class of persons therefore most liable to this disease are such as live upon inferior diet, and take hard malt liquor, and have that species of indigestion produced which is accompanied with the acetous fermentation. Those, on the contrary, who habitually take a great excess of animal food, together with stimulating liquors, of the vinous kind particularly, produce not only acid matter in the alimentary canal as a consequence of the faulty digestion, but also gradually induce an excess of supply to the circulating system beyond the power of healthy assimilation, and beyond the wants of healthy nutrition. In the former case, nutrition is deficient, and the body assumes the appearance of a want of adequate supply; but the gouty patient is exactly under the opposite circumstances, and bears, in his external characters, the stamp of repletion more or less strongly marked.

In the case of gout, the vessels, being surcharged with chyle, impose upon the kidney an increased exertion to excrete a part of that excess which cannot be properly assimilated; and thus, as it seems to me, not only is that disturbance in the natural balance of the elements of this secretion produced, which causes a separation of the uric acid from its usual combinations, but a considerable quantity of animal matter is also excreted, constituting the united product; namely, the pink or lateritious sediment; and hence my theory will appear to

explain why gouty persons cease to have so determined a form of gravel as they had at an earlier period.

Nature, ever wise in her intentions, in all these morbid affections of the digestive organs, whether coming under the denomination of gravel, or gout, or indigestion, or bilious complaint, appears to employ the kidney for a beneficial purpose,—to counteract the bad effects of the imperfect digestive process in the alimentary canal. That in the case of gravel, the salutary effort of the kidney to carry off unassimilated matter should give rise to another kind of disease, does but add one to many other instances in the actions of the animal œconomy under disorder, in which we see that the effort of restoration is sometimes itself productive of other form of complaint.

Of all the individual causes which have appeared to me to operate most strongly in giving rise to gravel, is the use of hard ale. The class of persons making this liquor their favourite beverage, also very commonly take spirits either raw or mixed. I have in numerous instances seen the patient obtain almost a permanent cure from this disorder by discontinuing such a baneful habit, and adopting a careful regimen.

In the ingenious Essay on Gravel, lately published by Magendie, the following causes are enumerated as chiefly productive of the uric acid species of this complaint.—“1st, An increase of the quantity of uric acid, the quantity of urine continuing the same, or not being augmented in a relative proportion to the acid. 2dly, A diminution of the urine, that of the uric acid remaining the same, or not diminishing in proportion to the quantity of urine. 3dly, A diminution of the temperature of the urine, the quantity and nature of it remaining the same as natural, or undergoing the modifications above described.”

I have stated at p. 117 the error of concluding that the uric acid is of necessity in excess in the instance of crystals being found in the urine. Magendie speaks of the uric acid as being merely solidified; and other authors have previously considered the red gravel to be a concrete state of the uric acid, arising from its great excess; but in addition to the fact just stated, I have repeatedly proved that it enters into new combinations. Also phosphate of lime is blended with the uric acid compound. I must therefore consider the formation of these crystals as derived from a species of indigestion which influences the action of the kidney, so as to alter the natural arrangement of the elements of the urine. In reference to the second position of Magendie, I must remark that the deposition of uric acid does not happen as a consequence of a deficient secretion of urine, which is of a deep colour and high specific gravity. Nor can I agree with the author in believing that a low temperature of the urine has any share in causing the crystallisation in question. These crystals are insoluble in boiling urine. I am led to entertain an opinion that this crystallisation takes place before the urine arrives into the bladder.

Further, it does not accord with my observation, that an excess of the use of animal food, producing a too great abundance of urea in the system, is a sufficient cause of gravel. Hence arises that kind of indigestion which is attended with a deposition of the pink or brick-dust sediment in the urine, but not of gravel, unless the influence of fermented liquors be added; of which, acid wines, perry, cider, but above all, hard strong ale, have the most noxious effect.

Of the very marked distinction which is due to the pathology of the two diseases, gravel and gout, I shall subjoin the following interesting illustration.

My friend, Mr. Travers, one of the surgeons of St. Thomas's Hospital, having noticed that several patients suffering from stone had been lately admitted into the hospital, from the district of country between Tunbridge Wells and Lewes in Sussex ; and hearing that others had been the subject of operation in the country, or were still labouring under the malady ; availed himself of a favourable opportunity of making some local inquiry into the facts ; and I had the pleasure to accompany him. We learned from an intelligent surgeon of Uckfield, that the gravel was prevalent among the poor in his neighbourhood, and that stone was of no uncommon occurrence. The class of persons, to whom the disease was said to be confined, lived almost wholly on vegetable food, and upon the stimulus of gin and hard beer. They were of course the labourers of the country, and were represented to be meagre in appearance, sallow looking, and by no means strong men. I examined a specimen of the domestic water of the village, which was common to the springs, and found it to be of remarkable purity ; its specific gravity at 60° was only 1.0020.

From another medical gentleman in the neighbourhood of Tunbridge Wells, I learn that gravel is a very prevalent complaint amongst the poor country people, affecting men, women, and children. He has mostly noticed the uric acid species. He describes their diet to consist of vegetables, milk, hard puddings, pickled pork, cider, hard porter, and spirits.

I contrast therefore these accounts of the habits and appearances of persons suffering from gravel, with the representation which I have given of the gouty patient at p. 136 and afterwards ; and an obvious inference follows from the statement.

Authors * upon gravel have confidently stated, that the urine which deposits the crystallised, or the pulverulent and coloured sediments (pink or lateritious), is preternaturally acid. So strongly indeed does Mr. Forbes entertain this opinion, that he speaks of an influx of acid matter† from the alimentary canal; and both he and Dr. Egan evidently consider, that the crystallised deposition from the urine is exactly analogous to the effect produced by the direct addition of any acid to healthy urine. Their experiments in proof of this opinion of increased acidity of the urine are however very fallacious, and do not prove more than is offered by the usual specimens of urine of high specific gravity, obtained from

* See, particularly, Forbes upon Gravel and upon Gout; and an Experimental Inquiry into the Nature of Gouty and Gravelly Concretions, by Thomas Egan, M.D. F.R.S.; Nicholson's Journal, vol. xvi.

† He also carries on the idea to the belief that such acid, taken into the circulation from the primæ viæ, detaches the uric acid from the circulating blood, and causes the gout by depositing the uric precipitate on tendons and ligaments, thereby exciting that inflammation (see p. 98, 99, &c. of his Treatise, 2nd edition) which constitutes the gout. Hence one false hypothesis is built upon another; for no proof is offered of the presence of uric acid in the circulating blood. Afterwards this author observes (p. 93), "The acid that occasions the precipitation may sometimes be phosphoric acid, of which the quantity is preternaturally increased; but more frequently it appears to be a strange acid introduced into the alimentary canal."

I formerly stated, that, in conjunction with Dr. Prout, I examined a flannel vest, which in different parts, from being long worn, was deeply reddened by animal matter, exhibiting appearances similar both to the pink and lateritious sediment; expect-

persons in health, not depositing the gravelly crystals. These observations are also to be taken in connexion with what I have said at p. 117. Whether the urine of persons who suffer gravel, be, *cæteris paribus*, more acid than that of persons in full health bearing a high specific gravity, cannot be determined without a series of careful comparative experiments. In the mean time, while I admit that the generation of acid matter in the primæ viæ is really the exciting cause of this peculiar action of the kidney, I deny the conclusion, that (as in the artificial experiment with urine and an acid) the crystallised precipitate in the disease called the gravel, is simply a deposit of the uric acid.

The same improvements in chemistry, which, a few years ago, led to the accurate knowledge of the composition of the urinary calculi, also became the foundation of a new method of practice in the treatment both of the stone and the gravel. Previously to this*, although the

ing that if the skin did secrete uric acid, it would be found under these circumstances. The coloured parts of the flannel were steeped in a dilute solution of pure potash, the liquor was filtered, and muriatic acid was added to a concentrated portion; but no uric precipitate followed. Respecting the question of a predominance of the phosphoric acid, I have already offered my details.

A second and very favourable opportunity occurred to me of experimenting upon this point. I examined, in the way I have stated, the flannel vest which was worn by a person throughout a severe paroxysm of gout. His urine deposited bright pink sediment very copiously. I could not discover in the matter of the cutaneous secretion the least trace of uric acid.

* Dr. Murray, in the fourth volume of his *System of Chemistry*, third edition, p. 590, has the following interesting observation:—"It is but justice to remark, that Dr. Wollaston's disser-

urinary concretions had long been an object of notice with the chemists, and Scheele had discovered the existence of uric acid in the urine, the treatment of them was conducted upon one uniform principle of belief, that alkaline remedies would be capable of acting as solvents for calculi in general. The analysis which chemistry has within the last twenty years afforded, became the natural step to new and improved views in practice.

Mr. Brande, in "A Letter on the Differences of the Structure of the Calculi, which arises from their being formed in different Parts of the Urinary Passages; and on the Effects that are produced in them by the internal Use of solvent Medicines*," has added several interesting facts to our knowledge; although I cannot help suspecting that the particular situation of the calculous concretion is dependant on accidental causes, and cannot be chosen as the proper ground of classification, either for the arrangement of the chemist, or for the guidance of the practical physician.

In the different situations of the urinary organs we find every species of calculus, and so indiscriminately blended, that the distinction in question seems to me wholly unsupported, and can only be well introduced as auxiliary matter of information.

tation was published in the Philosophical Transactions, two years before the memoir of Fourcroy and Vauquelin was read before the French National Institute; and although the experiments of Pearson, published in the Philosophical Transactions the year after Wollaston's, are referred to in that memoir, no notice is taken of those of the latter chemist; yet they anticipate nearly every thing which the French chemists have announced as their discoveries on this subject."

* See Phil. Trans. for 1808, Part II.

Mr. Brande, in speaking of the earthy phosphates, remarks, "I am induced to believe that the last-mentioned substances, although the production of the kidneys, and held in solution*, are never met with in a separate state till the urine has been at rest, and therefore calculi from the kidneys are never composed of the phosphates."

Whatever may be the particular situation in which the calculus is found, it is clear that we must make reference first to the action of the kidney, and further, in the series of the causes, to the state of the alimentary canal. Every patient who suffers from gravel, complains also, more or less, of indigestion. The immediate cause of the calculous sediment or concretion is the altered action of the kidney, by which the elements of its secretion become disturbed from the healthy balance, and undergo new and confused arrangements.

The bladder is evidently the mere receptacle of the urinous precipitates, and cannot be in any other manner concerned in the present theory of the disease; although I am quite aware, that it does take the chief share in producing that concretion of the respective matters, which we call the stone.

The comparative rarity of calculi composed of the uric acid and animal matter, as ascertained by Mr. Brande†, is quite in agreement with the fact which I

* I have pointed out, at p. 165, that the suspended phosphates are not soluble in the urine, even at its natural temperature.

† "The following is a statement of the composition of the different calculi found in the bladder which I have examined;—

have stated, of the complete solubility of the pink and lateritious sediments in the urine at its natural temperature (see p. 97); and we may conceive, that without a very favourable nucleus a concretion of this kind would never take place*. Hence again the reason of the unfrequent occurrence of the stone among gouty persons.

The distinctions in the composition of calculi, afforded by the researches of the chemists, were very naturally hailed with satisfaction by the physician, who was taught to believe that the solvents, which acted in an evident manner upon the calculi out of the bladder, would most probably dissolve them, even in the bladder of the living subject. Such was the view at first conceived; but the impracticability of using solvents as internal medicines of sufficient strength soon appeared; and an attempt, which was more particularly followed up by Fourcroy and Vauquelin, was made, of injecting solvents into the bladder. This, however, for the most obvious reasons, was a practice that could not be pursued without producing effects from irritation, more than sufficient to counterbalance the advantages of any chemical result.

16 were composed of uric acid.

45 ————— uric acid, with a small relative proportion of the phosphates.

66 ————— the phosphates, with a relatively small proportion of uric acid.

12 ————— the phosphates entirely.

5 ————— uric acid, with the phosphates and nuclei of oxalate of lime.

6 ————— chiefly of oxalate of lime."

* See account of the calculus, p. 164.

It was next the adopted theory, that although the power of medicine could not be confided in to the extent of procuring the solution of the stone in the bladder of the living subject, the principles of chemistry would still allow of very important application, for the purpose of counteracting the further formation of a calculus; and alkalies and acids were, as before, administered; although upon a different principle, according to the nature of the urinous sediment; and from which, the judgment of the composition of the calculus should be deduced.

Mr. Brande, in a second paper*, has stated some interesting experiments and instructive facts on the medicinal effect of magnesia as contrasted with that of the alkalies; the preference of the former medicine, as a remedy, having been suggested by Sir Everard Home, on the grounds that a neutralising substance, which was possessing the least solubility, would remain the longest in the stomach, and counteract "the formation of uric acid;" and that this was a more eligible practice than the fruitless attempts towards its solution when in a state of concretion.

The ingenuity of the idea was confirmed by experiment, as to the superior success of the remedy. A gouty person was also much relieved by its use, and hence, from that period, the employment of magnesia by persons subject either to gout or the red gravel has been in very general estimation.

Alkalies and acids, however, have still been cherished as remedies for the respective indications of a calculus consisting of the uric acid, and the earthy phosphates; and, indeed, Mr. Brande, in each paper, has, in concurrence with the general doctrine, advised the use of acids in cases of the last description.

* Phil. Trans. 1810. Part I.

It is now important to examine, what are the just pretensions of chemistry in relation to the practice of medicine, in complaints so important to humanity as the stone and the gravel; and first I shall discuss the treatment of the uric acid calculus.

I have already stated very fully the idea which I have been led to entertain of the cause of the formation of gravel; and consistently with my experiments and conclusions, I cannot admit the propriety of the hypothesis which I have quoted, that by the chemical effects of magnesia the formation of uric acid is prevented; and I should rather state, that by the influence of remedies on the alimentary canal itself, the kidney is favourably influenced in its secreting functions, so that the separation of the uric acid from its ordinary state of combination no longer takes place. This appears to me a distinction of some importance, as I shall presently endeavour to shew; but even if no distinctions in practice did follow from this difference of theory, the views of science demand that we should be accurate in our principles.

Further than this, I do not entirely admit the claims which are allowed to magnesia as a remedy in the morbid affections in question, and contend,—

1st. That although some advantage is afforded to the alimentary canal, by its power of neutralising acid matter, yet that its chief superiority over the alkalies depends on its purgative qualities; so much more easy is it to arrest the morbid process of indigestion, by a medicine which removes the cause, than by one which merely has the effect of temporary correction.

2dly. I condemn any unlimited confidence in this medicine, as being in most cases of complaint a very inadequate remedy for the disease which is existing. The gravel and the gout are, as it were, but the symptoms of

the morbid actions of other parts; and such primary disease, and true cause, is to be found in the digestive organs. But the gravel has a deeper foundation than the mere production of acid matter in the alimentary canal. In these cases we must look with vigilance to the state of the stomach, of the liver, and of the bowels, and of all the stages of digestive assimilation, in connexion with the wrong function of the kidney, in a manner less superficial than the rule of placing all our dependance on magnesia, or upon any alkaline medicine whatever, seems to imply. We shall learn that these remedies may be useful and even important as auxiliary parts of treatment; but that they do not deserve any higher character, or stronger dependance.

A lady of delicate constitution consulted me on account of the uric acid gravel, which had distressed her frequently for several years. She complained much of indigestion; but her appetite was seldom deficient. She suffered pain in the loins and region of the kidneys almost constantly; and sometimes was severely affected in a transient manner, with a pricking pain darting down from the seat of the left kidney to the bladder, obviously in the course of the ureter. This patient was put under a course of magnesia and carbonate of soda, without deriving any advantage: but in one week after the alterative use of *pilul. hydrarg.* and a draught containing *magnes. sulphat. infus. rosæ.* with *tinct. cardamom.* and four drops of *tinctura opii* as a dose, repeated once or twice daily, she derived the most sensible benefit.

I merely relate this case as one example amongst a series of cases which I could quote, of the superiority of the treatment by alteratives and aperients over the chemical remedies. I should observe, that, of her symptoms of indigestion, acidity of stomach was the most re-

markable. The bowels were usually relaxed; but if accidentally in the opposite state, the nephritic symptoms were invariably much aggravated.

Dr. Prout, in his paper "On the Chemical Properties and Composition of some of the proximate Principles of the Urine*," has added his valuable testimony on the preference which is due to the medical treatment, rather than the chemico-medical treatment of gravel. I take the liberty of extracting from this paper the following observations, which are exactly in point:—"Vitiating secretions of every description must be the result of general or local causes, or of both united. But when we reflect how little liable the secreting organs are to be affected, and how seldom, in point of fact, they are affected, except through the medium of the general health, we are naturally led to look here for the primary cause of their derangement. The inference is obvious. Remedies, no matter of what description, that have a tendency to restore the general health, must have a tendency to insure the due performance of all the bodily functions, and secretion among the rest. I need not enlarge here upon principles which are well understood, and the elucidation and application of which are justly ranked amongst the greatest discoveries of modern medicine; but shall merely observe, that by paying proper attention to the general health, and especially to the functions of the stomach and bowels, I have in numerous instances witnessed the speedy removal of urinary deposits, and the complete restoration of this secretion to its natural appearance and properties. In adults, as is well known, both the phosphoric and lithic diatheses prevail, and often alternate in the same person. I have, however, generally, seen both equally

* Medico-Chirurgical Transactions, vol. viii.

yield to the same principles of treatment, and sometimes even to the same remedy, and am disposed to think, therefore, that they are more intimately connected than commonly imagined.

“ When we reflect that all urine (except perhaps in extreme cases of diabetes) contains both lithic and phosphoric acid, although only one of the diatheses generally prevails at the same time, the conclusion is probable, reasoning merely *chemically* upon the subject, that the exhibition either of acid or alkaline remedies may produce harm as well as good : and if we take into account also the capricious nature of secretion, and the frequent alternation of these two diatheses in the same person from unknown causes, it becomes an exceedingly difficult task to adjust the remedy to the disease ; and the *chemical* probability will be, that the disease will ultimately be increased, instead of diminished. Lastly, the *object* of the chemical practitioner is at best but of a secondary description, namely, to prevent the effects of disease rather than to remove it. From these and other circumstances, therefore, which might be mentioned, I have been induced to consider chemical remedies as palliatives only, and to explain their acknowledged good effects even in this way, rather upon their general than their chemical operation.”

Magendie, in his late essay, has chiefly confined his practical considerations to the uric acid species of gravel, and in conformity to his theory, that this principle is derived from an excess of azote in the system, directs a restriction to those articles of diet which contain the least azote. He relates two cases in which the patients almost confined themselves to the use of sugar for their support, eating frequently more than a pound daily. The gravel disappeared, but after some weeks the plan was laid

aside, from its causing debility of the stomach and disorder. The following diet is recommended by the author :—" Without doubt, bread, particularly that made of rye-flour, pastry, the farinaceous legumens, Italian paste, rice, potatoes, the green legumens, sugar, &c. may be advantageously employed as food, particularly when simply prepared ; with these the patient may without danger satisfy his appetite. He must not, however, indulge in the use of these substances without regulation, as some of them, such as wheaten bread and pastry, contain a considerable portion of azote. When patients adopt this regimen, they must avoid spirituous liquors, and wine in an undiluted state, and they ought to drink copiously of aqueous fluids : such liquids rendering the urine more abundant without increasing the quantity of uric acid, will necessarily diminish the proportion of the latter to the former."—In prescribing the free use of any agreeable aqueous liquid, so as to increase the quantity of urine and prevent the solidification of the uric acid, he expresses that ten or twelve pints daily will form a proper quantity if the disorder be severe. He next considers the means of saturating the uric acid ; and enters into the common theory and practice of employing the alkalies, concluding with this observation. " But whichever of them may be employed, its evident effect should be an alkaline state of the urine*, without which nothing can

* I have just examined the urine of a gouty gentleman, who suffers from a constant secretion of uric acid and soda (*chalk stones*) in his hands and feet, in a far greater degree than any patient who has come under my observation. It reddened litmus in the usual manner, after becoming cold ; and yet, from the addition of muriatic acid, not the smallest precipitation of uric acid was produced. Hence would it not appear that the acidity

be expected from them in the cure of gravel; at least, no result which can be theoretically accounted for will be produced."

A case is related of a merchant who was alternately prosperous and in adversity, and lived accordingly; suffering both gravel and gout, when he indulged in the pleasures of the table; and obtaining an exemption from both complaints, as the necessity of abstinence occurred.

I have offered this brief abstract of the opinions of this author, because his celebrity is calculated to give new weight to the chemical method of treatment, with the addition of a severer mode of regimen than has hitherto been laid down. As I am led to entertain very different views on the subject, I wish to place the question before the profession as distinctly as my limits will admit, for their adjudication. I should explain the advantages of a plan of abstinence described by Magendie rather upon common principles, of giving repose to the digestive organs, and thus altering the action of the kidney, than upon the theory of merely lessening azote. I learn by a recent communication from Uckfield that the indigent patients, whose looks bear but little marks of repletion and of any excess of azote, suffer for the most part from the uric acid gravel. I would state that the patient should carefully pursue a regulated regimen, but that this should consist of a mixture of animal and vegetable food, every indigestible article being avoided; and that the drink at the dinner meal should be confined to water rendered conveniently agreeable, or soda water; and a limited quantity of sound wine. I strongly object to

of the urine is not to be referred to the uric acid? And it appears to me to give considerable support to the hypothesis which I have offered at p. 145.

large quantities of even aqueous drink, as tending to weaken the stomach and bowels by the distention which it produces. A continuance of the low regimen recommended by the present author, would, in my opinion, tend to produce much more evil to the constitution, than the inconvenience even of a complaint so troublesome as gravel.

On the effects of the chemical treatment contrasted with the use of alteratives and aperients, I have already offered my sentiments.

The administration of acids, and especially the muriatic, in the case of the earthy phosphates appearing in the sediment of the urine, is still recommended by recent authors in general, on the chemical principle of specifically counteracting such a secretion. Mr. Brande has further detailed his opinion on this subject in a late ingenious paper in the *Journal of Sciences and the Arts*, vol. vi. As it would appear from the ordinary theory, that the agent which holds the earthy phosphates in solution is an excess of phosphoric acid, it seemed to follow that in the event of a solid deposition consisting of the phosphates, the free acid was lost, and the urine must consequently become alkaline. The reasoning of Berzelius which I have quoted at p. 119, demonstrates very clearly the fallacy of this doctrine; and I have now arrived at the end of an extensive examination of specimens of urine depositing the phosphates, *not crystallised*, and also holding them in solution, yet still having the power of reddening litmus in the usual manner. I have also examined the urine of patients pursuing a course of the liquor potassæ in the quantity of two or three drachms a day, and of others under a long course of magnesia, or the pure or carbonated alkalies, and have not found a single exception to this fact. I suspect, therefore, that the opposite statements have often been made from

hypothesis only ; and for the sake of correspondence with the supposed theory to which I have just now referred.

I have here, however, to mention a curious fact, which I have several times witnessed, that if the vegetable alkali be given in union with lemon juice, in free doses, and at short intervals, and especially, I think, if in a state of effervescence, the urine soon becomes alkaline, and exhibits on its surface a great abundance of the ammoniaco-magnesianphosphate crystals*. The knowledge of this result has of late particularly inclined me to prefer the saline form of the draught mentioned at p. 186, usually also directing it to be taken in the state of effervescence, during the symptoms of acute gout, when the uric acid is deposited in great excess, as evidenced in the pink or lateritious sediment. While this diathesis prevails, the effect upon the urine just now described is not easily produced. I should add that the crystals are so infinitely fine that no sensible irritation is produced by them. They will disappear also very quickly after the discontinuance of the medicine, or from the administration of an aperient.

Returning from this digression, I must refer to the prevailing opinion of authors upon the skilful delicacy which is required to administer the alkalies and acids in calculous complaints, lest the happy balance of affinities should be destroyed by such inaccuracy ; and that from the want of a necessary knowledge or attention, we should add to the one or other species of disease, accordingly as our corresponding medicine should be misapplied.

That the mineral and vegetable acids will influence the state of the digestive organs very differently from the alkalies or alkaline earths, is a truism which I cannot

* I wish to point out this distinction. The urine is alkaline, when the crystals are formed ; acid, when the sediment is pulverulent.

wish to dispute; but the propriety of their administration on the chemical principles already stated I must entirely deny. Also, if we had it in our power to controul the action of the kidney with such facility, our judicious intention would be frustrated, from the circumstance, that the urinous sediments, and calculous concretions, are for the most part of a mixed nature in their composition.

It certainly happens, according to my experience, that when the gravelly deposit consists of the earthy phosphates, and which may be safely presumed when colouring matter is absent, that instead of the plethora of the habit and the state of vessels bordering upon inflammatory action, which usually accompany the deposition of the uric pulverulent sediments, more or less of debility, joined with a morbid sensibility of the nervous system, commonly prevails; and hence, the mineral acids which act as tonics to the stomach, will often be found useful. To point out once more the error of expecting to produce a chemical change in the state of the urine by medicines either alkaline or acid, as they are usually administered in the modern chemical treatment of gravel, I may first refer to the experiments of Mr. Brande, in which he made trial of very large doses of the alkalies, and was led to judge in that experiment which is most in point, that the effect of the alkali in becoming predominant in the urine "was at its maximum, probably in less than a quarter of an hour after it had been taken into the stomach; and in less than two hours, the whole of the alkali had passed off." I have already related the results of my own experience in general practice with alkaline medicine, in regard to the present question.

Berzelius says, "it is often impossible to diminish the acid of the urine by the use of alkali, in those who suffer from an excess of uric acid; and I have myself tried in

vain the effect of acids in neutralising or acidulating an alkaline urine. A middle aged man was laid up with the gout: his urine was foul and alkaline, holding the earthy phosphates suspended in an undissolved state. I gave him the sulphuric acid, without any change; and afterwards the phosphoric, without any effect, until its dose was so much increased that it became laxative: the urine then became acid, and deposited uric acid as long as the laxative effect continued, but no longer, although the dose of the acid remained unaltered.

“Lastly. I tried the acetic acid, and with as little success*.”

Having now pursued this theoretical discussion to a sufficient length, I arrive at a conclusion, which I have verified in numerous cases, that in proportion as our means of treatment, in the calculous affections of the kidneys, are most favorably adapted to restore the healthy state of the digestive organs, they are, in the same ratio of effect, calculated to correct and remove any errors in the secreting action of the kidney itself.

A careful scientific observation of the sediments of the urine is important in a two-fold point of view. It is the surest ground which we can take for the anticipation and prevention of that calamitous disease the stone; and it likewise offers a valuable source of information, by enabling us, according to my experience, to discriminate the particular morbid states of the digestive organs, and the relative condition of the sanguiferous and nervous systems.

When the urinous precipitate is of the uric acid description, we find in association with it such a prevalence of acid matter in the alimentary canal, that the use of the

* View of Animal Chemistry, p. 107.

alkalics, and the alkaline earth magnesia, occasionally becomes of important service as auxiliary medicine; but when we reflect on the fundamental truth, that the gravel in its determined forms, no less than the gout, is a disease created by long continued errors of diet, it must at once appear demonstrated, that a more radical method of treatment is required, than such medicines alone can accomplish. The cure is to be found in the judicious employment of purgatives, alteratives, and corrective tonics; and, for permanent security, in the just regulation of every part of the general regimen.

In addition to the description which I have already given of the class of persons most liable to gravel, I may observe, that, in adults, the uric crystals (red gravel) prevail with the greatest frequency; and in children, the sediment in which the phosphates abound or alone exist, commonly called white sand, is the more usual occurrence: but, both in adults and in children, we meet with the two species of crystallized precipitate mixed; and this still more frequently happens, when the sediment is pulverulent.

The question of selecting the use of the mineral acids, when the crystallized sediment is consisting distinctly of the phosphates, is now briefly to be considered.

Mr. Brande, in the paper to which I have last alluded, has fully entered into the subject of the use of the mineral and vegetable acids, in cases of the present description; advocating their employment upon chemical principles, with the same force that he recommends the alkalies, when the crystallised sediment consists of the uric acid compounds.

In several cases the most exactly adapted to my purpose, in which the sediment consisted of the mixed phosphates without any portion of uric acid, I made trial first

of alkalies and acids, separately, and precisely according to the chemical theory; but with no permanent good effect; and in some instances with the disadvantage of causing painful irritation by means of the acids. Certainly I did not disturb the uniformity of the chemical state, which is so much talked of, and substitute the deposition of the urates for that of the earthy phosphates. Nature herself is often capricious in producing this change, but it is very little amenable to the influence of art. I next prescribed occasional purgatives, and, as a daily plan of medicine, a combination of the carbonates of soda and magnesia*, with sulphate of magnesia, to be taken in any simple vehicle, with the addition of lemon juice, so as to make it an effervescent draught. Together with this draught, I directed a mild mercurial oxide, in the alterative dose, with rhubarb and soap, and also enjoined a strict plan of diet and regimen. I thus perfectly succeeded in effecting a permanent change in the action of the kidney; and in due time, directing the use of stomachic tonics, the health became entirely restored.

Magendie, who has argued in favour of the use of the alkalies when uric acid, as he believes, is in excess, so much, that he speaks of the use of magnesia in a quantity of ten grains to an ounce or more in the twenty-four hours! expresses himself in the following discouraging terms upon the employment of acids, when the urinary sediment consists of the phosphates:—"I have several times endeavoured to attack the phosphate deposits from the urine, by the mineral and vegetable acids, and have never

* When the urinous sediment consists of the phosphates, we find reason to believe that the acid matter produced by indigestion equally abounds in the alimentary canal, as when the urates are deposited. This is at least true as a general statement.

obtained any satisfactory result. It has indeed appeared to me that the deposit has been increased by these means."

When inflammation has been excited in the kidneys, the ureters, or the bladder, by means of the calculous crystals acting as irritating foreign bodies, the usual means of treatment as for the phlegmasiæ in general, are for the most part to be employed. If the inflammatory diathesis prevail, bleeding must not be neglected. The warm bath is a remedy of unequivocal propriety, and often affords very prompt and material relief. Purgatives are of indispensable importance; and the use of calomel and antimonial powder, with a portion of the colocynth extract, jalap, or rhubarb, in conjunction with a saline purgative, usually insures the best effects. Respecting opiates, I think it necessary to make a few observations. The spasmodic nature of the pain which attends the nephritic attack, would seem, on the principles on which I have before advocated the use of opium, to demand and justify its free administration; but some exceptions in the present case of suffering are to be pointed out. It is our object to promote the secreting action of the kidney; to dilute the urine, and so to increase its actual quantity by the influence of the mildest soothing drinks, very freely taken, that it will at the same time be less stimulating to the passages under inflammatory irritation, and by actual distention of them produce the effect of dislodging and carrying off the minute calculous concretions. Experience shews us that the effect of opium very usually tends to keep the secreting function of the kidney, when it is under inflammatory irritation, in further restraint, even although it be used in combination with purgatives. It is therefore a medicine which, upon this general principle, is to be made questionable in its internal use, in this disease; and we shall

sometimes find it more advantageous to administer it in the way of injection into the bowels. This practice should be pursued with very great freedom in the dose. In conclusion, however, I must be permitted to observe, that if such intense and sudden pain, as sometimes afflicts the patient, call for instant relief, we shall find that we cannot withhold the prompt and ample exhibition of opium ; and which, administered on the principles I have stated under the head of Narcotics, even with all the general objections just now stated, will hold its sure dominion over the present most immediate evil, the pain of the attack.

When the nephritic symptoms assume the chronic form, similar modes of treatment are to be pursued, with that moderation, however, which the milder disease invariably allows.

I shall conclude this brief sketch of the present subject, by relating the happy effects of a warm bath in a case of acute nephritis, which occurred to a medical gentleman, fifty years of age, of full habit and of the sanguineous temperament. His habits of living were indulgent, but not excessive. He stated that most commonly his urine deposited either pink or lateritious sediment. He had never been affected with gout. The bowels were habitually inactive, and he experienced occasional indigestion. One morning, when thinking himself as well as usual, he was suddenly seized with an acute pain in the situation of the left kidney. In half an hour urgent vomiting took place. The pain passed down in the course of the ureter to the left testis, which was strongly retracted. The efforts to pass urine were very constant and agonising, and it was voided only by teaspoonfuls. He was free from fever, and the pulse was not affected. He lost sixteen ounces of blood by cupping

on the loins, with a little relief. He rejected from the stomach fluid purgative medicines, retaining however some calomel. Two hours elapsed without any material ease being procured. He then used the warm bath, and almost instantly obtained from it delightful relief. He went into bed freed from acute pain, complaining only of a dull heavy sense of aching across the loins. The stomach soon retained purgative medicine, and by means of which the cure was in a short time effected. He had reason to suspect that he passed some gravel when in the bath. By careful regulation of the bowels, and proper attention to regimen, this gentleman has passed over twelve months without any return of complaint.

OF PROPHYLACTIC REGIMEN.

I HAVE to consider the present very interesting subject in the following several points of view; namely, as it relates to the gout; to gravel; and to the morbid states of the digestive organs in general.

The means of preventing the return of gout, are in general more easily pointed out than practised; so great is the force of established habits, and so irresistible is the love of indulgence. In truth, however, it does not appear to me that gouty persons should be bound down to any rules of living which can be considered painful, provided they exercise a little philosophy; or that they should be

obliged to use any care, which a slight perseverance may not render agreeable. Indeed the real comfort, as well as benefit, which follows the observance of such wholesome restraints in diet, as the particular constitution of each individual patient may require, very largely exceeds the transient pleasure of any sensual gratification afforded to the palate.

The discussion of the present important consideration may be divided into what relates to the general management; the diet; and the occasional use of medicine.

I may first observe that the views of *prophylaxis* are both prospective and retrospective; and those who may be led to expect the gout by inheritance, cannot too early be taught to pay the most careful regard to their constitution; nor too surely confirm the best habits by long practice*.

When the disease has made its invasion, the permanent cure rests indeed with the patient alone: and if he be wisely his own physician, before its tyranny be established, future prevention will be in a great measure at his command.

I know no rule more important to be mentioned, than that the patient, who finds out by sad experience his weakest points of attack, and the peculiar influence of particular remote causes, should guard himself against them with a nurse's eye, and with corresponding care.

* The little Treatise of Lewis Cornaro, on the sure methods to attain a long and healthful life, is well deserving the reader's best attention. He emphatically exclaims, "Oh, what a difference there is between a regular and irregular life! One gives longevity and health, the other produces diseases and untimely death."

Thus some, from exposure to wet and cold, incur either certain gout or rheumatism ; while others, although very gouty subjects, escape such trial with impunity. I have met with two gouty persons in particular, who can actually remain in wet clothes without any consequent injury ; but if they commit any remarkable excess at the table, an attack of gout almost with certainty follows. Such extreme instances are rare ; but there is good ground for the general assertion, that every gouty patient is rather more susceptible to some one remote cause than another : and although he cannot be too much a practical philosopher in his whole conduct, he should be scrupulously vigilant against the enemy in his most vulnerable quarter.

When the residence is a matter of convenient choice, a gravelly soil on a middling level, and protected from the north and the east wind, should be selected for the purpose. A situation which is damp and cold, or even damp and warm, is much to be condemned. For those who are not yet rendered delicate and enervated by the long influence of the disease, I would be disposed to prefer the bracing quality of a cold dry climate, to the relaxing powers of one that is hot. It must at the same time be stated, that those who are tender feel enjoyment only in summer months, and some even obtain exemption from their gout by removing to a hot climate. Van Swieten, Haller, and other authors relate some instances of extraordinary cure produced by such emigration. I have met with some gouty persons, who found themselves free from the disease during their residence in the East Indies, where business had taken them, although their habits were not more moderate than before. This seems to admit of explanation, from the profuse perspirations which regularly occur in such situations, and which ob-

viously counteract the plethoric state of habit*. For this reason chiefly, I presume it is, that the gout does not often make its first invasion in a hot climate, even with those who live freely. We also find that some robust persons of a sanguineous temperament, who are not subject to gout, in returning from a hot climate in which they enjoyed their health, to the variable one of this country, acquire an inflammatory disposition of vessels; and the explanation of the fact has already been offered.

Further, I may mention, that the benefit of equal temperature is most sensibly felt by gouty persons. Thus I have known of some who have quitted this country for India, with a gouty and rheumatic state of pain and infirmity, afflicting them upon their embarkation and afterwards; but no sooner have they arrived in a warm latitude, than ease has returned; and gradually their natural recovery has taken place. I must, however, observe, that no climate nor place of residence can give protection against the effects of improper habits of living. I know some gentlemen who experienced their first attack of gout when in India.

When we consider that vicissitude of temperature, or, in other words, wet and cold, is the most frequent of all the exciting causes of gout, it is incumbent on us to consider the best means of lessening the susceptibility of the body to the impressions produced by such means. During eight months of the year in this climate, the utility of wearing flannel next to the skin is too well known to require comment; and whether it should be discontinued

* I do not find that gouty persons are particularly liable to deficiency of perspiration. The contrary indeed happens with the majority, as they come into the class of the *corpulent*; with whom, exercise produces a very free action of the skin.

in the warmer four months, must always be a matter of careful consideration. Even a leather waistcoat, over a thin one of flannel, is advisable in a few instances, when we see the susceptibility to the influence of damp and cold so remarkable, that more or less of rheumatism or gouty sensation follows from every slight accidental exposure.

Every gouty patient should keep in mind, the infinite importance of preserving the feet *always* dry, and comfortably warm: but while this caution is observed, the debilitating influence of *hot covering* should be studiously avoided. Upon the employment of cold sea bathing as a remedy to strengthen the frame, I can only recommend a general rule of caution, it being a very doubtful measure for a gouty subject; and the cold fresh water bath should be considered as almost inadmissible. The temperate bath of the Buxton temperature (82°) will be found most advantageous, both to strengthen the constitution and the limbs.

From well established experience, I can confidently advise, as equally safe and useful, the following daily practice; on which, indeed, although I have already spoken, I shall now dwell more at length. Sponge every morning the whole of the feet, between the toes, all around the ankle joints (and the knee joints also, if they have been the seat of the complaint), with salt water, or water in which salt is dissolved to the point of saturation*; care being taken, that the *chill* of the fluid be

* The most convenient method which I can recommend, is, that a large covered vessel should be kept in the chamber, containing bay salt and water; and it is only necessary to notice, that the salt is always in excess; which may be concluded when any appears at the bottom undissolved. A gallon of water requires about three pounds of salt. Each morning a clear portion is to be taken out, as much as is wanted for use; and to

always just removed by the addition of a sufficient proportion of boiling or very hot water.

The skin being wiped perfectly dry, diligent hand rubbing (the best kind of flesh brush) should be employed for as long a time as is convenient; and should invariably be continued until a sensible glow of the skin is produced. In the whole process, one part should be finished before another is begun, lest evaporation should take place from the moistened surface in an unfavourable degree. It will sometimes be objected that this daily attention is too troublesome and tedious, and that the benefit, which it is likely to produce, is not worth the purchase. Such observations rest on the same weak basis, as those which apply to general care in regimen; and spring only from want of exertion, and the misapplication of time. The comfort of this cleanly custom is very great, and the assistance which it affords to the weakened parts, and to the proper functions of the skin, is of such importance, that I am persuaded its auxiliary power in the *prophylaxis* of gout is not inconsiderable. The numerous communications which I have received on this point are quite decisive in its favour. The subsequent friction* of

be rendered of the necessary tepid temperature by the addition of hot water. By the thermometer the range may be from 80° to 90° of Faht. The patient should rest the heel on the shelf of the tin japanned foot tub, when sponging the parts, so that the water passes into it as it is used.

* Sir William Temple, in speaking of the benefit of friction, strongly observes, "No man need have the gout who can keep a slave."

"Dessault cite un exemple connu à Bordeaux, d'un viellard centenaire, qui trente ans avant sa mort s'étoit garanti et guéri de la goutte, à laquelle il étoit fort sujet auparavant, en se faisant brosser et frotter chaque jour, soir et matin, avec une main garnie d'une mitaine de laine."—Barthez, tom. i. 194.

the parts possesses the obvious advantages of increasing the superficial circulation, and exciting the absorption of such deposits as previous inflammation has produced, either in the moving textures, or in the cellular membrane. In proportion as the energy and strength of the limbs are preserved, the patient not only obtains the power of using regular exercise, but also fortifies the parts very usefully against future disease. By familiarising the surface to the moderately cool temperature which I have mentioned, the susceptibility even of the whole body to atmospherical changes is in a considerable degree lessened.

With reference further to this latter important object, I recommend the patient to wash the whole head every morning, by means of a coarse towel dipped in water perfectly cold; using afterwards dry friction with the towel for a sufficient time. I have for a long time witnessed the advantages of this custom, without knowing a single instance of inconvenience resulting from it.

The powerful effect of *indolence*, in assisting the first invasion and all the returns of gout, brings us to reflect on the great value of regular exercise among the remedies of prevention. It is the injudicious practice of some persons, to mix indolence with exertion, by being inactive at home during the greater part of the week, and taking excessive exercise on occasional days. I know some gouty sportsmen who exemplify this statement very strongly, in performing even feats of riding at particular periods, but are sedentary in their general way of occupation. In this manner a state of exhaustion rather than wholesome fatigue is produced; and when by accidental excess in walking, more particularly, the weakened joints are thus *over-exerted*, the intended benefit is sometimes converted into an exciting cause of a paroxysm.

Horse and foot exercise should each be used as answering different intentions. These means of health, so useful to all persons, are of particular importance to the gouty, with whom repletion is so great a part of their disease.

Sydenham expresses himself very forcibly in these words, after speaking of horse exercise:—"And indeed I have often thought if a person was possessed of as effectual a remedy as exercise is, in this and most *chronic* diseases, and had the art likewise of concealing it, he might easily raise a considerable fortune."

The shaking exercise of riding on horseback appears well calculated to quicken the mesenteric circulation; to increase the gravitation of the contents of the alimentary canal; and to urge the peristaltic motion. It is a familiar fact, that active exercise of this kind is a speedy remedy for a fit of the *piles*, when unattended with inflammation.

Those persons, and especially the elderly, who have a weak and languid circulation in the extremities, should be attentive, when using horse exercise in the cold season of the year, to clothe their limbs with great care.

The additional advantages of walking exercise are made apparent, in the consequent improvement of strength and motion afforded to the lower limbs.

Such as are crippled, and disqualified from the modes of exercise already mentioned, should daily take an airing in a carriage; but it is only some real necessity of this kind*, or the obstacle of bad weather, that should

* On the useful influence of various modes of gestation, and the reasons for which carriage exercise is sometimes to be preferred, see Dr. Wollaston's interesting Essay, Phil. Trans. 1810, Part I.

give it more than occasional preference. Those who court indulgence must not expect exemption from the gout. In some instances of martyrdom from the disease*, a sudden adversity has proved a piece of good fortune! Even the influence of bad habits of living is exceedingly counteracted by active exercise and labour. The gout very rarely visits the poor man's cottage.

Many examples might be quoted, illustrating how effectually the predisposition to gout has been restrained, by taking a timely warning, and adopting an active course of life, with every observance of careful habits.

The superior advantage of the fresh air of the country over the confined atmosphere of the metropolis, is on all occasions sensibly felt by the invalid; and is no less calculated to preserve than to restore health. The gouty citizens of London should be studious to unite daily exercise in the country, with their less salutary pursuits in town.

The hours of rest and of rising deserve an attention of which the gouty class of society are too often unmindful; but their error consists rather in taking too much repose than too little. Neither more nor less than eight hours in bed, I should offer as the best general rule; and that the pillow should be sought before twelve, and quitted before eight.

In how great a degree health of mind belongs to health of body, I need not argue. The union of these blessings was the poet's warmest prayer:

“ *Orandum est, ut sit mens sana in corpore sano.*”

The physical rules of health belong also to the moral

* Van Swieten's Commentaries, § 1255, and Hoffman's Rat. Syst. Med. vol. v. 518.

code; and serenity and cheerfulness come as uninvited guests, where health presides.

As far as concerns the intellectual faculties, it must be stated that very sedentary habits of application, and long-continued intense thinking, are highly unfavourable with relation to the gout. I am acquainted with a gouty gentleman, who was formerly a London accomptant; and he relates that his severest fits were always induced by occasional excess of application to his business. Some who have retired from the occupations of the town to a country life, have lost their gout with a total change of situation and habits; and all receive benefit by such a removal. The rational and active exercise of the faculties is as much to be recommended as that of the body; and the proper rule consists only in avoiding excess of study. Sydenham, in the prefatory epistle to his Treatise, declares, "that his immoderate application to his work occasioned the severest fit of gout which he ever had;" and apologises for having confined himself at the time to the subjects of gout and dropsy; adding, "that the gout constantly returned, as often as he attempted to go on with his work."

I come now to consider the most favourable system of diet.

As, in the acquired disposition to gout, excess in diet is the indispensable remote cause, and, even in the hereditary disposition, is more or less the necessary antecedent to its developement, it follows that this part of my subject is of the highest importance to be considered. It appears to me, that a general rule of abstinence has been too much insisted upon, for the gouty; and that little is often performed, because too much is required. I am disposed to contend, that the security of the gouty does not, except in particular cases, require their being re-

strained from such a portion of the good things of the table as affords the truest enjoyment. With regard to the solid articles of food, it may be said, in general terms, that the choice should be of such as agree best with the individual. *Quod sapit, nutrit*, is an axiom of much truth; yet its use requires limitation. I can assert, from much observation on what happens with gouty persons especially, that the most usual injury which improper diet produces, takes place in the intestinal canal; and, as I have before observed, in the duodenum especially. The mind is unconscious of the disordered functions of secretion which thus arise in the advanced process of digestion; and the patient not finding any sense of oppression or indigestion in the stomach itself, feels satisfied and convinced that his food must have been wholesome, and suitable to his constitution.

An exact rule of diet for general application can scarcely be offered; for in some idiosyncrasies, the most wholesome food proves the most injurious. Dr. Spurzheim informs me, that his coadjutor, Dr. Gall, cannot ever partake of *mutton*, in whatever manner it may be dressed, without immediately suffering irritation in his stomach in a most severe degree; and the same effect has with certainty followed, when the dish has been purposely disguised by his friends.

I might mention many other extraordinary instances of individual peculiarity, with respect to those simple articles which agree so well with persons in general. The succeeding observations have a general application to the health of invalids, for whom attentive regimen is necessary, whatever their chronic disease may be.

When the *kind* of diet is wisely chosen, according to the lessons of individual experience, the next and most material rule of caution, is the *quantity* of food which is

habitually taken. An accidental indigestion from some particular article of food may often cure itself; but the consequences of habitual error in quantity are of a much more permanent nature.

The advantage of dining at an hour not later than four in the day, is very well known; but with this salutary custom, the rules of fashionable life are wholly at variance. When the hour of dining is early, any intermediate nourishment from breakfast may be avoided; but in the opposite circumstances some refreshment should be taken; rather, however, upon the principle of preventing that exhaustion of the stomach, which arises from a long-continued absence of stimulus, than as affording a greater supply of nourishment. To the majority, some plain seed-cake, to which carraway seeds and a little ginger may be usefully added, will prove very sufficient. In the summer season an orange, also, will not be improper. When thirst is troublesome, well-made rennet-whey is more to be recommended than soda water, which, if taken freely, causes distension, and is sometimes unfavourably stimulating. I have seen it agree better, when the excess of carbonic acid has been allowed to escape; and its pleasantness is much increased if some sugar, and a very small portion of lemon-juice, be added. Thus mixed, it may be taken in effervescence.

One, whose avocations are fatiguing, with feeble powers of constitution, must be permitted a more supporting method. An egg lightly boiled, with bread, and wine and water; or calves' feet jellies, with bread or biscuit; or a little well-seasoned plain soup, may in this case be recommended to the patient's choice. It may be laid down as a general rule, that meat should be eaten only once a day. It may be optional whether the

breakfast shall consist of tea, or bread and milk, or light cocoa or chocolate; but coffee at this meal has appeared to me objectionable, as it is rather a heating stimulant, and not so good a diluent as tea. Butter in moderation, good in quality, and spread upon the bread without exposure to the fire, is wholesome and quite allowable.

A few observations in detail upon the dinner meal may not be unacceptable to the reader, in connexion with this part of my subject.

By the gouty patient, and also by those not gouty, whose digestive functions are improperly performed *below the stomach* (that is, chiefly in the duodenum, where the assimilating process may be considered to be most actively carried on), the principal error is committed, as before observed, in the *quantity* of food at dinner; and one great incitement to this excess, is the variety of articles which cookery presents to the palate, in its most tempting forms. Under these circumstances, not only is food taken in quantities much greater than correspond to the natural appetite, but an excessive excitement is given to the stomach by the varied nature of the stimuli; and which must, sooner or later, lead to debility of that organ. With respect to gouty persons, I believe there are few examples in which we do not hear of what is called “a good stomach,” in the early part of life. Simplicity of diet, indeed, consists principally in the small *number* of the articles to be eaten at the same meal. By the virtuous observance of this rule, the appetite will seldom urge the taking of more than what is useful; and the stomach will be sufficiently spared, both in its secreting and muscular functions. Also the subsequent process of assimilation will be more easily and perfectly performed in all its stages; and unhealthy corpulency, and a radical disposition to the true plethoric state, will

be obviated. I would offer as the best rule, that only one kind of meat should be partaken of at the same meal; but I would not prohibit fish as a preliminary part of that meal, if desired by the patient as a gratification. A little soup, also, may without impropriety be occasionally first taken, in addition to what I have mentioned; but I must observe, that such as are remarkably plethoric, should generally make a point of avoiding nutritive soups; and every one who has any regard for his stomach, and considers its consequent distention from indulgence in this bulky nourishment, should be sparing in its use. A proportion of some well-boiled vegetable is as useful as it is agreeable. Most kinds of pudding may be eaten with propriety; but baked pastry, to the butter of which too much heat has been applied; and also confectionary, in most of its forms, should be entirely avoided. A little old cheese, good in quality, seems to assist digestion. It is a material point that the dinner meal should be deliberately eaten, and the food well masticated.

Thus far, then, as to the quantity of the chief meal of the day; in addition to which, the other meals should only be considered as accessory refreshments. How much is the picture, which I here offer, reversed by the practice of the majority; and more especially by the indulgences of the votaries of fashionable life! With those, whose breakfast, and luncheon, and dinner, and sometimes even supper, consist of animal food, with corresponding liquors having more or less of heating stimulus, must it not follow, that the gout or some other disease will be induced? It will occur indeed as an effort of Nature to interrupt the destructive process, and accomplish the reduction of the system by means of pain, and the effects consequent on pain.

Those who disregard all rules of temperance, can neither receive nor deserve a cure from the physician.

With respect to the particular kinds of food, and modes of cookery, which may be considered as the most wholesome, a few further observations will, I hope, be found sufficient. To expatiate at length on this subject would require a separate treatise.

Experience seems to shew, that of the flesh of all the larger animals used as food, *mutton* is the most easy of digestion; and particularly when roasted. Veal, however, appears to have its texture more favourably prepared for the action of the stomach by boiling*. It may be stated to be the least digestible of the meats in general; and this fact is referable to a principle which I think may be laid down, that animals, such as sheep and game, which are allowed to range in fields, acquire much muscle and little fat in proportion, while the reverse of this takes place in stall-fed cattle, which are much covered with cellular texture and fat, instead of the more useful qualities of muscular fibre. When the animal is not too aged, it is the muscular fibre that affords the best stimulus to the stomach, and the most favourable material for digestion. Pork, in most of its forms, appears, with the invalid, to be a meat of difficult or uncertain digestion; and, whenever eaten, its skin and fat

* It appears to me, that the skin and cellular part of meat are more favourably prepared for the agency of the gastric juice (to express myself familiarly, are rendered more fit for digestion) by the influence of the boiling process; but, on the contrary, that muscular parts are rendered comparatively more loose in their texture by means of roasting. As illustrations of the probability of this reasoning, I may mention the articles, calf's foot, and veal, to exemplify the first position; and the second is instanced by the superior tenderness of beef moderately roasted, over that which has long been submitted to the boiling temperature.

should be avoided. I learn, however, from good authority, that pork broth (quite freed from fat) agrees remarkably well with very weak stomachs.

Among fish*, salmon, from its fatty nature, must be considered as unwholesome, in whatever way it may be dressed; and the most objectionable, next to salmon, are the eel and mackarel. Shell fish, although agreeing well with some persons whose stomach is not healthy, should rather be placed in the list of things forbidden†. But it must be said of oysters, and especially when boiled (the beard always being removed), that they suit a weak digestion in general remarkably well; and of their very nutritious quality, I need not speak. With regard to vegetables, all such as are in season, and well

* The herring would appear to deserve particular praise, as a meal for the gouty. Dr. Clerk (Edinb. Med. and Phys. Ess. vol. iii. p. 462) relates a case, in which the eating of salted herrings, every kind of drink to relieve the thirst being carefully avoided, very readily cured the paroxysm. If we reason upon this case, we must apparently refer the success to the stimulant or other properties of the muriate of soda; but as a remedy in gout, it appears to me as little worthy of recommendation, as it is certainly disagreeable in its sensible effects.

† It must be kept in view, that the present very critical observations relate only to the management of the digestive organs, when in a weakened and disordered state of function: and it is important also to observe, that the conclusions which I have now ventured to offer upon diet, have been derived from the reports of dyspeptic persons; among which class of invalids, I have sought every convenient opportunity of obtaining information on these points. In connexion with the present subject, the experiments which were performed by Mr. Astley Cooper, and are detailed at the end of this section, will be found interesting; and particularly deserving the attention of the physiologist.

boiled, may be esteemed as useful as they are agreeable; but I should recommend that when the stomach is weak, only one kind should be taken at the same meal; and whatever vegetable produces flatulency should be considered unsuitable.

In some instances of great debility of the stomach, so that a very moderate meal produces an oppressive state of distension, apparently caused by fermentation, it proves useful to the patient to abstain wholly from vegetables. The nature of the animal food should also be attentively considered. There are certain grounds of observation, upon which a medical judgment may be formed, to determine whether the diet, in particular cases, shall consist most of animal or vegetable food. These grounds are, the nature of the various secretions.

To pursue my subject, I may remark that dressed onions usually agree well with gouty stomachs. Of raw vegetables, those which are stimulant, as the horse-radish, mustard, &c. are alone proper; and I may here take occasion to notice, that a moderate portion of the stimulus of pepper and mustard, when relished with the article eaten, is advisable, under the circumstances of languid digestion; but certainly, the habitual free use of condiments should be guarded against: they are not necessary to a healthy stomach. Salt is a stimulus which we may be said naturally to require in union with food; yet it should be taken in moderation on account of the thirst which it excites, and the distension of the stomach and bowels consequent upon an excess of fluid. Vinegar should never be taken but sparingly. New pickles are much to be condemned; but such as are old and well-seasoned, partaken of in moderation, sometimes agree, even with a faulty stomach. Yet, in the list of things that are proper, they are not the articles which deserve

to be particularly chosen; and, together with raw vegetables, the one or two above-mentioned excepted, are to be excluded from the regimen of gouty persons.

Before concluding, I must observe of bread, that those persons who are subject to habitual costiveness, will find great advantage in shunning the astringent white bread of the baker, and in taking brown bread made with flour from which the coarsest part of the bran only has been removed. I am acquainted with many examples in proof of the utility of attending to this apparently trifling change in the plan of diet.

I proceed now to treat of the *fluids* of the table, with relation to the prophylactic management.

I have already dwelt (p. 69 and 101) at some length on the comparative influence of different fermented liquors; and return now to the position, that, with respect to the gout, the quality and quantity of what is drunk is of still more importance than of what is eaten: but on this head, also, I have no restrictions to propose which are inconsistent with the true enjoyment of social life. As a general statement of the question, I would contend, that a small portion of wine after dinner is as useful and innocent for a gouty person as for any other. Some few, who are of a sanguineous temperament, and are much disposed to the inflammatory plethoric state, would be wise in shunning all fermented liquor, and adhering to the safe beverage of water; provided that such a plan proves favourable to the nervous system; but on the other hand, I am convinced that I offer a concession on the side of truth, in stating, that the gouty, for the most part, acquire, by that course of living which has brought on the disease, such a state of stomach as concurs with the influence of their temperament, in rendering the moderate

use of wine both useful and necessary*. It produces a comfortable feeling; and its beneficial effects, when it is used within the strict rules of propriety, are felt throughout the whole frame. I am disposed to consider, that taking less than three glasses is an unnecessary forbearance; and that taking more than six is the beginning of excess and injury. In determining the exact quantity most favourable to the individual patient, I may observe, that a truly careful attention to his own feelings would, for the most part, be a sufficient guide; the object being this, that the wine should produce a feeling of comfort, without any sensible heating excitement.

* I feel it important to explain my sentiments on this point without ambiguity. On reviewing the above doctrine, I am convinced of its correctness. It should be considered, that the particulars with regard to diet, which constitute a very material part of medical treatment, must be modified in relation to particular individuals, in the same way as medicine; exercise; or any other class of remedies. Every general rule of regimen must, in each separate instance, be varied in some of its circumstances. Such gouty persons, for instance, who are yet young and full of vigour, would wisely fulfil the object of prevention, by confining their beverage to water. The second class, whose strength has been impaired by the disease, may with benefit employ wine, under the limitations which I have expressed. The third class, whose constitution is become infirm and nervous from long and repeated attacks, *require* the cordial power of a little good wine; and with all other correct management united with such cautious indulgence, will, I am persuaded, improve the tone of their stomach and the energies of their general health, without aggravating the predisposition to gout.

The direction of a glass of wine as a measure is obviously indefinite, and I know not how to make my meaning clear, otherwise than by stating a glass to be equivalent in quantity to an ounce and a half.

Some very interesting cases have come under my observation, in which, not only an abstinence from wine and every kind of fermented liquor has been steadily pursued for a great length of time, but even animal food has been entirely avoided; with a view to prevent the returns of gout. The results of three experiments of this description I have related at p. 321, 400, and 441, and I shall here offer a few additional facts.

C. R., aged 48, of vigorous constitution and rather plethoric, much subject to *hereditary* gout, discontinued the use of wine and all fermented liquors for eighteen months. He did not procure a longer interval from gout than usual, nor was the succeeding attack slighter. He felt his stomach so sensibly weakened, and his nerves so much depressed, in consequence of the plan, that he returned to the moderate use of wine; and with equal comfort and advantage.

H. A., aged 54, slight in form, and not plethoric, subject to *hereditary* gout from the age of 24, abstained wholly from wine and all fermented liquors for fifteen months, also being very sparing in the use of meat. At the end of this period he was attacked with gout more severely than on any former occasion; in a greater number of parts; and the fit was longer than any which he had ever experienced.

J. K., aged 52, tall and slight, and not plethoric, subject to severe attacks of *acquired* gout since the age of 32, strictly discontinued the use of animal food, wine, and all fermented liquors, for the long period of four or five years. When two years of this probationary regimen had expired, he was attacked with gout most severely; the fit proved of more than usual duration; nor was there less inflammation of parts than on former occasions.

These facts appear to me sufficient to shew, that the

entire disuse of wine by a gouty patient does not carry with it the certain advantage of counteracting the returns of the disease, which some contend for; and I consider that in most instances it is not a safe measure to the powers of the constitution: but I wish it to be clearly understood, that while I advocate this doctrine, I only recommend the very moderate use of wine; under different degrees of restriction with different persons; and desire of all, without exception, an attention to every other point of regimen, as indispensable.

I must add, also, my entire acquiescence, that a suspension, for a time, of nutritive diet and of the use of wine is capable, in a few instances, of affording material benefit to the constitution. Indeed, in some cases of long-established visceral obstruction and vitiated state of the alimentary secretions, both in gouty persons, and in others, I have found the happiest consequences to follow from the temporary adoption of diet, consisting chiefly of milk, vegetables, and farinaceous food, in conjunction with a course of medicine; but such a plan has been provisional only, and with reference to existing disease; and is entirely to be distinguished from the regimen of a severe abstinence for life.

To resume my subject, I may state, that, in the choice of the particular kind of wine, a little latitude may be allowed, provided it be *old and of the best quality*. I have met with some gouty persons, who feel both comfort and advantage from good claret, and are heated and inconvenienced by a much smaller quantity of good port. Setting aside the idiosyncrasy of particular individuals*,

* I know a lady who is invariably rendered immediately ill by the smallest quantity of port wine, although she drinks white wine with benefit. Her symptoms are an hysteric difficulty of breathing, extreme general nervousness, and head-ache.

it appears to me that good old port is very well suited to the stomach of a gouty man; and that sherry, when old and genuine, is more favourable to him than Madeira; it being, as I think, the least acescent. I am induced to form this conclusion, from a few trials which I have made with good specimens of each wine, as to the comparative proportion of an alkali which they have required, in order to be rendered neutral. The most injurious wine which the gouty patient can drink, is, Burgundy; and fortunately it is the most difficult to be procured. The general rule of avoiding all light acescent wines, and also such as are *new*, whatever the vintage may be; and sub-acid liquors, as cider and perry; is a caution so familiar to the gouty, that I need only advert to it in this general manner.

During dinner, *water*, for the purpose of dilution, is the true beverage of health. If any wine be taken at the same time with the food, it should be in very sparing quantity. Of the useful beverage, water, however, some persons drink an injudicious quantity, thereby distending the stomach so as to weaken its muscular powers. In most cases, I conceive that half-a-pint with the dinner meal is sufficient.

Against strong *malt liquors*, the objections are very numerous. Ale and porter, from their superior nutritive properties, being much more conducive to plethora than wine, must have a strong influence in inducing and increasing the predisposition to gout; and in the dietetic plan of those who take wine, the use of these liquors should not be permitted. Small beer of good quality, and free from all acidity; or, in the technical expression, not in any degree *hard*, is a wholesome beverage; and, although less salutary than water, it need not be altogether denied, except to the corpulent and plethoric; and provided it be taken only in careful moderation, and that dyspepsia is not present.

*Spirits**, however diluted, should be shunned as a certain enemy; and therefore never taken, except under the prescription of the physician, in order to answer some particular intention.

In directing my consideration to the most proper regimen for persons afflicted with gravel, I may make useful reference to my narrative at p. 576. The patient should be instructed to avoid all those causes which tend in any degree to produce acid fermentation in the stomach and alimentary canal. Thus, *hard* malt liquors, and especially of the stronger kind; cider; perry; spirits in every form; a diet too much vegetable to allow to the powers of the stomach and the general system the natural support which is required; are the causes of injury chiefly to be avoided. The labourer, upon whom the disease most frequently falls, must assist his strength with such malt liquor only as is entirely free from acidity; and good porter appears to be the most useful. The affluent patient should confine himself to the moderate use of pure wine; shunning all the kinds in which acid predominates. When medical treatment is required, the use of ass's milk is much to be recommended.

Another point, in particular connexion with the present subject, now presents itself.

* Although good wine contains so large a proportion of alcohol, as to amount generally to one-fourth, according to the experiments of Mr. Brande (Phil. Trans. 1811, Part II.); yet the spirit is in such a state of combination with several peculiar principles, that its influence on the stomach, when thus combined, is very different from that of raw spirits, mixed or unmixed; and more especially, in proportion as the wine has received its improvements from age, are its stimulating properties favourable, and less likely to be injurious.

In every age of the world, and in every medical writing upon regimen and diet, the qualities of water have been a favourite and just object of consideration. Nature does, indeed, present an astonishing diversity in the composition of her springs, derived from the strata through which they flow. The beautiful and important chemistry which is thus going on in the inmost recesses of the earth, and which furnishes to man both for his uses in health, and for his occasion in sickness, the most salutary beverage, and often the most valuable medicine, is one of the innumerable wonders of the creation, which must raise our admiration and our gratitude to the supreme Author of every good!

The kinds of water which are chosen for our domestic use, are such as are the most grateful to the palate, and are divided into the *hard* and *soft*. This distinction has been deduced from the difficulty or facility with which the respective kind forms an admixture with soap. If difficult, the inference follows that much saline matter is contained; which, attracting the alkali from the soap, leaves the oil detached, forming flakes or curds with the water. If, on the contrary, an easy *lather*, as it is called, take place, it indicates that the saline matter can be present only in small proportion; and the water, being free from other substances, is then denominated pure. The kind of water denominated hard has always been considered as unfriendly to health, and as specifically injurious to persons afflicted with gravel and stone. Many probably imagine, that the earthy salts which it contains assist in making up the mass of the calculous concretion. If we view the composition of water according to the general doctrine, it is curious to consider, that the only ingredients which it is found to contain of difficult solubility, are such as are seldom or never found in

urinous concretions; namely, the sulphate and the carbonate of lime*. Unless, therefore, they undergo decomposition in the alimentary canal, or in the circulation, this cannot be a mode in which they would become injurious. In the more rational theory, however, of the natural composition of water, as Dr. Murray has with so much originality and useful ingenuity suggested, the arrangement of the constituent elements appears, in the majority of cases, to be wholly different from that which is commonly conceived. According to the views of this chemist, the almost insoluble salts in question, the carbonate and the sulphate of lime, do not actually exist in the proportions which are usually stated, if at all; but on the contrary, their acids arrange themselves with other bases, so as to produce the salts which are soluble; and hence the hardest waters cannot be supposed to possess the unwholesome qualities so freely assigned to them. It is not indeed probable that Nature would present to the hand of man, a fluid, which should be grateful to his palate, necessary to his wants, and at the same time injurious to his health.

I disbelieve the general conclusions which are entertained of the hardness of water, as the cause of gravel and stone; and doubt not, that the true and principal

* Mr. Howship, in his "Practical Observations on the Diseases of the Urinary Organs," gives the following account of a very large renal calculus, which was in structure "compact and earthy, in consistence resembling bird-lime, and of a pale yellowish grey colour." By analysis, Mr. Brande found this matter to consist of carbonate of lime, mingled with an extremely tenacious animal matter. He observed, "that it was the first instance he had ever known of the kidney secreting the carbonate of lime."

explanation will always be found in the patient's unfavourable mode of living as to regimen and diet; and in the use of liquors much less innocent than the water of the district, which so commonly receives the whole reproach.

Notwithstanding these remarks, I do, however, freely admit that a water which is the most free from foreign ingredients, approaching to the state of *distilled*, is the most favourable diluent with our food; the best solvent in the digestive process; and better calculated to counteract indigestion, than a water strongly impregnated with earthy salts, whatever their state of combination may be. In this manner, then, I accommodate my reasoning to the facts which occasionally appear, that persons, who suffer much from gravel, are extremely sensible to the influence of particular waters which they drink; and invariably declare that they suffer inconvenience from the use of those which are hard. If this be true, the explanation, as I conceive, must be founded entirely on the sympathy before explained between the digestive organs and the kidneys; and hence, whatever most conduces to the healthy process of digestion, harmonises with the general feelings of the patient, and promotes every healthy function.

With respect to the use of medicine in the prophylactic treatment of gout, we are to be directed chiefly by the state of the alimentary canal and of the kidneys. The daily action of the bowels is of essential importance to every gouty person; and the degree in which this may be required, will have reference to the particular constitution of the individual. For the purpose in question, as well as on account of some useful influence it may have in exciting secretion both in the alimentary

canal and in the kidneys, I can, from experience, recommend the pill prepared according to the following formula:

R Gum. gambog. gr. j.
Pilul. hydrarg. gr. v.
Pulv. alöes compos. gr. v. ad x.
Saponis duri gr. ij.
Decoct. alöes compos. q. s. fiant pilulæ iii. vel iv.
hora somni sumendæ, alvo astricta, vel pro re nata.

The following is a pill of more simple composition, and adapted to familiar use in a still greater degree; viz. a dose of five or ten grains of the pulvis alöes compositus, half a grain or a grain of pulvis jacobi, with a grain or two of soap (to obviate the hardness of the pill, when kept in quantity for use), formed into two or three pills with the decoctum alöes compositum. For those who are corpulent, and readily disposed to become plethoric, I should add to the dose half a grain or a grain of gum. gambog. that a greater diuretic effect may be produced.

It was the theory of the humoral pathologists, that in gout, as in several other diseases, the blood should occasionally be subjected to a regular course of purification, by means of vegetable medicines, consisting of various and opposite qualities, intended to modify each other, so as to harmonise into one general purpose. Thus, Sydenham advised an electuary composed of thirty-one ingredients. He adds to his observations for its use; "But care must be had to make choice of such simples as are most agreeable to the patient's palate, because it must be continued a long time, namely, for the most part of his life." He also recommended a dietetic decoction, consisting chiefly of sarsaparilla, which he desired to be begun immediately on the going off of the fit, and to be

continued in the intervals, “ during the remainder of life.”

The occasional employment of the corrective vegetable diluent, the simple or compound decoction of sarsaparilla of our Pharmacopœia, in conjunction with the alterative use of *Plummer's* pill, may be useful in that state of stomach which is accompanied with much itching of the skin, and occasional irritable eruption ; but either as a daily beverage in the manner here mentioned, or as an important remedy, it does not seem entitled to much confidence in relation to the gouty diathesis. I think that the merit of causing the mercurial alterative to agree better with the stomach and the constitution, is due to this medicine. I have seen instances in which the compound decoction has proved heating in its operation, and the simple has afterwards agreed perfectly well. In general indisposition of a chronic nature, arising from, or connected with dyspepsia, this decoction combined with potass. carbon. appears sometimes to have a very useful influence on the stomach. In order to save the patient the necessity of drinking a large quantity of fluid, it may be desirable to condense the medicine, by joining the extract of sarsaparilla with the decoction.

It is with much satisfaction I state, that many of my patients, accustomed to be attacked with gout severely at particular periods of the year, have pursued a moderate course of corrective and aperient medicine in union with regulated regimen, with such success, that they have entirely prevented the gout from taking place ; and have also derived very sensible benefit to their constitution.

In regard to the best means which we can employ to prevent the return of gout, it should fairly be considered that one cannot say to a patient, “ You never shall have the gout again.”—This would be tantamount to an assu-

rance to any other person, "You never shall be ill again." As a general statement, it is true, that any remote cause of injury, such as indiscretion in living, exposure to wet and cold, violent passions of the mind, bodily injury, excessive fatigue, and so on, which in another individual not disposed to gout, will usually produce some form of disease, accordingly as the predisposition may be, does, when applied to the gouty patient, bring into action the gout, and most frequently the gout alone.

Alkaline medicine has been often recommended in the gout upon chemical principles, in reference to the theory of uric acid being the proximate cause of the disease. Dr. Wollaston, in describing the composition of gouty concretions, concludes his paper* as follows: "The knowledge of this compound may lead to further trial of the alkalies, which have been observed by Dr. Cullen to be apparently efficacious in preventing the returns of this disease (First Lines, par. 558); and may induce us, when correcting the acidity to which gouty persons are frequently subject, to employ the fixed alkalies, which are either of them capable of dissolving gouty matter, in preference to the earths (termed absorbent) which can have no such beneficial effect."

At p. 582, I have mentioned the grounds on which the use of magnesia has been preferred to the alkalies, as a remedy in *gravel* when the uric acid is supposed to be secreted in excess. The practice has been extended also to the treatment of gout. Mr. Brande, in the paper before quoted (Phil. Trans. 1810), relates the case of a gentleman who had attempted the use of the alkalies with much disagreement to the stomach, and next resorted to magnesia. Having pursued the plan for some time,

Phil. Trans. 1797. p. 337.

he was considerably relieved from the appearances of uric acid and mucus in the urine; and the author concludes with stating; "It is also deserving of remark, that there has not been the slightest symptom of gout from the time of the last attack, which is more than a year back; a longer interval of ease than this patient has experienced for the last six years."

Doubtless, this favourable report of the effects of magnesia has had a very chief influence in encouraging its habitual employment, by persons subject either to gout or gravel.

On the theory of the operation of alkaline medicine, in cases of a supposed prevalence of uric acid in gravel, I have already very fully offered my sentiments; and it remains that I should briefly discuss the practical question of the prophylactic powers of this class of remedies with respect to the gout.

A gentleman severely afflicted with gout, informed me, that in the space of two years he took thirteen pounds of subcarbonate of soda, with the hope of successfully preventing the accession of the disease. He was led to the use of this medicine, in part from the urgent feeling of heart-burn and other symptoms of dyspepsia; and in part from a strong confidence in its prophylactic powers. The result was, that he found a slight palliation of his dyspeptic symptoms; but his fits of gout were scarcely influenced by this treatment, either as to frequency or degree. When he came under my care, it was evident that he had long suffered from an unhealthy state of liver; and being treated accordingly, he received decided benefit, and procured an exemption from gout for an unusual period of time.

In Case X. at p. 312, I have mentioned the temporary advantage obtained from the continued use of mag-

nesia; and I am in possession of a copious list of cases, in which, gouty persons have given to this medicine a full trial, and with a zealous confidence in its powers. So long as it has acted as a purgative, it has proved more or less useful to every patient; and many, in the advantage which they have for a time derived, have begun to hail it as a panacea for all their ills. Others have questioned its good effects, from the uncertainty of its operation on the bowels; some have found it both inert and to disagree; and in no case, in which I have investigated its effects, has it appeared to produce any very permanent improvement, so as to afford the desired exemption from gouty attacks.

From much observation of the effects of this remedy in gout, I have been led to the decided conviction, that its only useful agency has been in its conjoint action as an antacid and a purgative; and that in no other way, than by such influence on the stomach and alimentary canal, can I consider that it has any claims to our regard.

It is at the best, therefore, in most cases, only a palliative treatment; and sometimes carries with it this serious evil, that the practitioner and the patient may be proceeding with an inefficacious or even hurtful remedy, in a case which is demanding some radical means of medical treatment, and the best regulations of diet and regimen.

The habitual employment of magnesia is further liable to an objection of considerable importance, which has lately been brought to light. In the *Journal of Science and the Arts*, No. 11, Mr. Everard Brande has related two very striking and instructive cases, in which, the long continued use of magnesia was productive of a concretion of this earth with the mucus of the bowels, in an immense quantity; and which, in the case wherein the details are stated, gave rise to many of the worst symp-

toms attendant on obstruction of the intestinal canal. They were removed only by active purgatives, which expelled the concremented matter.

This patient has since come under my own care; and I have every reason to believe that an excessive quantity of mucus in the intestinal canal was the occasion of the concremented accumulation in question. It is rather remarkable that the magnesia almost invariably acted as an aperient.

In the second case it is stated, that "not only large quantities of a concretion of a similar description were voided, but, upon examination after death, which took place perhaps six months after any magnesia had been taken, a collection, supposed to be from four to six pounds, was found embedded in the head of the colon, which was of course much distended."

The popular use of magnesia, and the rarity of such consequences as are here described, would scarcely seem a sufficient ground of objection to its continued employment as a domestic remedy; but these cases ought to be kept in view by medical practitioners, and the administration of magnesia should be laid aside, when any remarkable obstruction of the intestines is manifested.

Dr. Whytt relates an instance of the remarkable effects of *lime water* in procuring entire exemption from the returns of gout, which before had been frequent and severe*.

The patient took it in large quantity, so that its purgative operation was considerable. It is evident from the narration, that its chief effect depended upon its operating in this manner. Sir Gilbert Blane, in making remarks on the effects of the pure fixed alkalies and of

* Edinburgh Med. and Phys. Ess. vol. iii. p. 459.

lime water, in several complaints, states, in reference to the *gout*, "that in some cases they seemed to protract the intervals of this disease, while in others my expectations were entirely disappointed." The occasional use of soda water, to an extent that does not produce uneasy distention of the stomach, will be attended with good effects to the gouty, as well as to any other dyspeptic invalid.

In final conclusion of this part of my subject, I have to state, that, in my own experience, I prefer, in the class of alkaline medicines, magnesia either pure or carbonated, and the neutral carbonates of soda and ammonia; but, in the use of the former medicine, I never prescribe it as a separate remedy; and most commonly direct either of the latter articles as one only among other ingredients. Also I join with any of these substances which I may employ, the use of some purgative and alterative medicine, on which I can place a certain dependance; regarding the alkaline agents as useful chiefly from their neutralising power, and in that manner as being auxiliary to the plan of cure.

It may be useful to observe, that when acidity and flatulence prevail, the calcined magnesia is much to be preferred to the carbonate. Its superior strength I have mentioned at p. 186.

With a view to the prevention of a plethoric state of the vessels, and of the consequent introduction of a paroxysm, some have resorted to general bleeding at particular seasons. I have met with instances in which the taking away blood for another disorder, in a gouty person, has received the credit, and perhaps deservedly, of remarkably postponing the subsequent attacks. It should be considered, however, that what is good as a remedy, is bad as a habit. The prevention of an excess of

blood, by suitable abstinence and exercise, is the only legitimate method of counteracting the plethoric state. On the idea of lessening the quantity of circulating blood in some degree, and of establishing a drain for supposed morbid matter, issues have had their advocates. This treatment appears much too local to act upon the constitution for the prevention of gout. It is a positive and disagreeable inconvenience; and without a clear indication for its employment, and with different views from those which I have now stated, may justly be superseded by other and less objectionable means.

Upon principles of practice founded on the doctrine of debility, *tonics* have been often employed as a remedy for the prevention of gout.

Of the *Portland powder*, once a favourite, but now an obsolete remedy, little need be said, except in reference to some general reasoning which it suggests. Its composition* resembled the *annalia medicamina* of Cælius Aurelianus, and other gout antidotes of the ancients. Dr. Herberden spoke of its effects with some approbation. Dr. Cadogan censured it severely; remarking, that "he had observed between fifty and sixty of its advocates, some his patients, some his acquaintance or neighbours, who were apparently cured by it for a little while; but in less than six years time, *omnes ad internecionem cæsi*, they all died to a man†." A stimulant bitter, as this was,

* For an interesting essay on the subject, see "An Inquiry into the Origin of the Gout Powder." By John Clephane, M.D. Med. Obs. and Inqui. vol. i.

† Cadogan on the Gout, &c. p. 79.

taken for many months in succession in daily doses, would have the effect of exciting the appetite beyond the powers of digestion and healthy assimilation; and hence it might indirectly lead to occasional apoplexy, which it was accused of doing. In the employment of tonics during the interval of the paroxysms, with a view to prevention, we should bear in mind, that the gouty require correction rather than excitement of the digestive organs: and this is to be effected by a regulation of the secretions, and of the action of the bowels, by means of occasional medicine; and by attentive regimen.

I have already spoken of chalybeate tonics under some circumstances of debility; but I am not disposed to admit of their propriety as *prophylactic* in the time of health. A chalybeate water, so useful to the constitution in many instances, is seldom admissible in the gouty habit; in which the vessels are easily stimulated to unhealthy action: and indeed I am well acquainted with some instances, in which, this agreeable remedy, resorted to by the patient on the general principles of giving strength to the constitution, has in a short time brought on a gouty attack.

Upon a review of the whole of the present subject, we arrive at the conclusion, that the prevention of gout consists rather in doing little, than much, by the employment of medicine; except as may regard that important object, the due regulation of the bowels; and unless, from the faulty state of the constitution, the use of medicine be really required; in which case, as I have described at p. 624, we may interpose our treatment with such success, that we anticipate and prevent the accustomed return of the paroxysm. I mean therefore to signify by the above remark, that in regard to the gout, the common principle holds good which applies towards all other diseases, namely, that the cardinal rules of preserving health are founded on temper-

ance and exercise; on the choice of all those means which are found by the individual to invigorate the system; and the shunning of whatever tends in any way to injure and enfeeble the constitution.

THE EXPERIMENTS ON DIGESTION,

Referred to at page 612.

By the obliging friendship of Mr. Astley Cooper, I am enabled to offer his statement of a series of experiments which he performed upon dogs, with a view to ascertain the comparative solvent power of the gastric juice upon different articles of food, and to derive any useful conclusions which they might afford, for the dietetic management of the human stomach, when in a weak state of digestive power. These experiments were related in the lectures which Mr. Cooper delivered at the Royal College of Surgeons, five years ago; but they have been published only in this Treatise.

In the conducting of these experiments, every practicable uniformity of method was observed. The substances were cut to a determinate form, and weighed. They were then forced into the animal's throat. A given time having elapsed, the dog was killed; and the substances not yet dissolved by the action of the gastric juice, being again weighed, their loss, and consequently their degree of *digestibility* as food, under the healthy action of the stomach of the dog*, was estimated. Raw food, and the lean parts only of meat, were given, except when the contrary is expressed.

* Approaching nearly in structure to the stomach of the human subject.

EXPERIMENT I.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Pork	long and narrow	100 parts	1 hour	10
Mutton	_____	_____	_____	9
Veal	_____	_____	_____	4
Beef	_____	_____	_____	0

EXPERIMENT II.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Mutton	long and narrow	100 parts	2 hours	46
Beef	_____	_____	_____	34
Veal	_____	_____	_____	31
Pork	_____	_____	_____	20

(EXPERIMENT III.

Kind of food.	Form	Quantity.	Animal killed.	Loss by digestion.
Pork	long and narrow	100 parts	3 hours	98
Mutton	_____	_____	_____	87
Beef	_____	_____	_____	37
Veal	_____	_____	_____	46

EXPERIMENT IV.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Pork	long and narrow	100 parts	4 hours	100
Mutton	_____	_____	_____	94
Beef	_____	_____	_____	75
Veal	_____	_____	_____	69

It is probable that the digestion of the dog, with regard to pork, differs from that of the human subject; as,

when the human stomach is at all weakened, the order of digestion in these meats appears to be,

1 Mutton

3 Veal

2 Beef

4 Pork

Something also may be attributed to the absence of *fat* in the above experiments; and more especially the fat of pork.

EXPERIMENT V.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Cheese	square	100 parts	4 hours	76
Mutton	_____	_____	_____	65
Pork	_____	_____	_____	36
Veal	_____	_____	_____	15
Beef	_____	_____	_____	11

EXPERIMENT VI.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Beef	long and narrow	100 parts	2 hours	0
Rabbit	_____	_____	_____	0
Cod fish	_____	_____	_____	74

Hence it appears that fish is easily digested*.

* In the Edinburgh Medical and Surgical Journal, No. 49, the Reviewer (by whose flattering approbation of my Treatise, I cannot be otherwise than much gratified) observes, in commenting on these experiments, that "it may admit of a question, whether quick solution in the stomach be a proof of *easy* digestion." This objection is ingenious, but I conceive not valid; and I am disposed to contend, that, *cæteris paribus*, we

EXPERIMENT VII.

Kind of food.	Form.	Quantity.	Loss by digestion.
Cheese	long and narrow	100 parts	29
Fat	_____	_____	70

EXPERIMENT VIII.

To the same dog, beef and a portion of raw potatoe were given, of each, 100 parts.

			Loss by digestion.
Beef	-	-	100
Potatoe	-	-	43

The skin in contact with the potatoe was not acted upon. Under the skin the potatoe was dissolved, but the gastric juice had not reached its centre. Where the skin was separated, it was dissolved.

In the dog, according to the following experiments, roast veal is more difficult of digestion than boiled.

EXPERIMENT IX.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Roast veal	long and narrow	100 parts	—	7
Boiled ditto	_____	_____	—	30

EXPERIMENT X.

Kind of food.	Form.	Quantity.	Animal killed.	Loss by digestion.
Roast veal	long and narrow	100 parts	—	2
Boiled ditto	_____	_____	—	31

are entitled to the inference, that solid articles of food are digestible in the stomach in proportion to their degree of ready solubility. It is true that these experiments relate only to the digestive powers of the dog, and of that animal in health; but they are nevertheless instructive.

The following materials were next the subject of trial.

EXPERIMENT XI.

Material.	Quantity.	Animal killed.	Loss by digestion.
Muscle	100 parts.	4 hours	36
Skin	————	————	22
Cartilage	————	————	21
Tendon	————	————	6
Bone	————	————	5
Fat	————	————	100

The appearances produced were these :—In the muscle, a separation of the fibres by gradual solution of the connecting media first took place; and afterwards, the fibres themselves became broken down into very minute portions.

The skin, upon its under surface, was broken down; but upon its upper surface was not altered.

The cartilage was apparently worm-eaten.

The tendon was reduced to a pulpy gelatinous substance.

Further experiment on the digestion of bone.

EXPERIMENT XII.

		Animal killed.	Loss by digestion.
Thigh bone	100 parts	3 hours	8
Ditto	————	6½ hours	30
Scapula	————	6 hours	100

In the human subject the stomach is capable of acting upon bone; of which fact, the following case is an example.

On Monday the 28th of March, a little girl, nearly four years of age, accidentally swallowed a *domino*, which passed the bowels in rather more than three days. The medical attendant (Mr. Maiden, of Stratford) observing that it was much smaller in size than the other dominos of which it had been one in the set, was induced to weigh it, and found that its weight was only 34 gr. while that of the others was 56 gr.; so that it had lost by digestion 22 gr. Those surfaces of the domino, which, when swallowed, were hollow and blackened as usual, were found prominent like buttons.

ON

RHEUMATISM.

PREVIOUSLY to the short Treatise which I am about to offer on the present important and interesting subject, I must beg the indulgent attention of my readers to a few explanatory observations.

In the concluding part of my preface to the first edition of this work, I professedly attempted to give a general outline only of the nature and cure of Rheumatism; and reserved the practical consideration of the chronic species of the disease, “entirely as matter for a future volume.” Less than a year having elapsed, when the favourable reception of my book demanded another edition, I reflected that I should discharge my duty better to the public, and, as I conceived, also more effectually sustain the little credit which I had already acquired, by applying myself diligently to improve and enlarge my first volume, rather than to hurry to the press prematurely, a *second*, which I could not, in so short a space of time, have the opportunity to execute in any degree to my own satisfaction.

Two years having now passed away since the second edition of this work left the press, it may reasonably be expected that I should have redeemed my early pledge of extending very considerably my observations on rheumatism; and indeed this was my intention when I began the present volume. But so copious have been my

materials on the preceding subjects, and so irresistibly and insensibly have I yielded to the usual zeal of authorship, that already I have anticipated nearly all the space which a single volume will conveniently allow; and must therefore be now contented to circumscribe my new details within very narrow limits, while, at the same time, I beg yet to retain the privilege of offering a more complete treatise on the subject at some future opportunity, should my present Essay receive sufficient approbation to encourage me in such an undertaking.

Of the general application of the term *Arthritis*, which was in use with the ancients to express every kind of disease affecting the joints, I have before made mention; and, at the same time that this want of nosological distinction forbids any precise confidence in their practical authority on the subjects of gout and rheumatism, it proves the slow advances of medical science. The confounding together of gout and rheumatism, as one disease, demonstrates to my mind a remarkable inaccuracy of observation: for, although they produce occasionally some local symptoms in common, we see that in a genuine example of each disease, both constitutionally and locally, the most obvious difference of characters is presented. Sydenham himself observes of rheumatism, "This disease, when unattended with fever, is frequently mistaken for the gout, although it differs essentially therefrom, as will easily appear to those who are thoroughly acquainted with both diseases; and hence it is, perhaps, that physical authors have not mentioned it; unless indeed we esteem it a new disease." Ballonius in a treatise "*De Rheumatismo et Pleuritide dorsali**,

* Paris, 1642.

appears to have had the merit of first treating of rheumatism as a separate disease. He relates that it had previously been confounded with catarrhus and arthritis. He bestowed upon it the name of rheumatismus, deriving its etymology and expression from *ρευμα*, a defluxion; and still preserving a full belief in the doctrines of humoral pathology, as the following account will shew.

He considered the pain of rheumatism to be produced by the acrimonious qualities of the impure blood, passing off to the skin through the muscles and nerves as a *epuratory* process. Gout, he remarks, is a disease of a certain part, and periodical; rheumatism, of the whole body, and uncertain in its time of coming. He thought, however, that without much care in the habits of life, rheumatism would certainly lead to the gout. Attributing the disease to a corrupted state of the blood, he considered venesection as the only fit method of evacuation according to the usual principles of humoral practice, which require the offending humour to be evacuated by its appropriate and respective channel. He affirmed that all medicines taken by the mouth were hurtful, and served to increase both the pain and inflammation.

Dr. Cullen has defined rheumatism according to the following translation:—*A disease from an external and often an evident cause; pyrexia; pain about the joints following the course of the muscles, fixing upon the knees and larger joints in preference to those of the feet or hands, increased by external heat.* He next establishes the division into two species, the *acute* and *chronic*; the presence of pyrexia serving to mark the former state of the disease; and its absence, the latter. I shall speak of these forms, both together and separately.

SEAT OF RHEUMATISM.

IN defining the seat of rheumatism, a less general mode of description than that adopted by Dr. Cullen appears to me required.

The fibrous textures of the body may be stated as the true seat of rheumatism; and most commonly the tendinous structure is the part affected*. The general expression, "of the joints," is very indefinite. It comprises some parts of structure, which may fairly be questioned as being the legitimate seat of this disease; for example, capsular ligament, and cartilage. It is probable that these parts become more frequently affected with inflammation in persons who are subject to rheumatism; but it appears to me a more fixed and deep-seated affection, than constitutes the peculiar character of rheumatic action. The ligaments which are most susceptible to rheumatism, are rather those which are subservient to the joint, as accessory, than as immediate connexions; and those which belong to muscles and tendons. The ligaments of the vertebræ, which are most external, are sometimes affected, and occasionally in a separate and distinct manner. The sheaths of tendons, the aponeuroses or fasciæ (tendons expanded on a wide surface), and the bursæ mucosæ, are parts of structure much more com-

* In the review of my observations, with which I was honoured in the *Annals of Medicine and Surgery*, Sept. 1816, I find an interesting quotation from a Thesis upon inflammation, which is stated to have been published by Dr. Elliotson, some years ago. In this I have the pleasure to see a remarkable coincidence with my own opinions, as to the usual seat of rheumatism.

monly affected than ligament. Whether or not, in a muscle, each constituent part be primarily affected with this inflammation; or, whether it be confined to the tendinous portions, is a question not easy to be determined. The latter conclusion seems the most probable. The pain which in some states of rheumatism is felt only upon the action of a muscle, may be explained on the presumption that it is propagated from the tendinous insertions along the course of the fibres; or, that the aponeurosis being affected, it is put upon the stretch, and causes the pain and tenderness which are felt upon motion; giving to the patient's sensations the idea that the fleshy part of the muscle is the seat of complaint.

Dr. Carmichael Smith, in his useful and very original paper* on inflammation, defines the acute rheumatism, "an inflammation of the muscular fibres." This is evidently too limited a view of the subject, if not erroneous. From frequent careful examination, by pressure, of the whole course of a muscle, when its functions have been affected by severe rheumatism, the inference has in general clearly presented itself to me, that the internal fibrous structure has not been the seat of complaint. In reference also to the fugitive character of rheumatic inflammation, the immediate subsequent recovery of muscular action, and the permanent nature of the symptoms which appear to mark a diseased state of the fibres of a muscle, in such distinct cases as have come under my observation, I am principally led to the conclusion, that any in-

* Of the different kinds or species of inflammation, and of the causes to which these differences may be ascribed.—*Medical Communications*, vol. ii.

inflammatory process from rheumatism, which may occasionally take place in the muscular fibre, is a remote consequence only, and not a primary character of the disease. If the muscular fibres were really the common seat of inflammation in rheumatism, there would surely be the attendant phenomena of swelling and tenderness in the substance of the muscle, much more distinctly than we find to take place; and might we not expect, as a common consequence, more or less of thickening of the fibres, in correspondence with the effects of increased determination of the blood, which are invariably manifested in other parts? The reverse of this very commonly happens in rheumatism with respect to the muscles; but we may always discover an increase of bulk in the tendinous and bursal structure.

When the muscles remain permanently weak after a rheumatic attack, and often waste in size, the symptoms will admit of sufficient explanation from the morbid and impaired state of the associated tendinous and bursal structure, and the consequent disuse of the muscles.

It would seem, from the soreness which the patient sometimes describes as if fixed in the bone itself, that the *periosteum* is occasionally the seat of rheumatism.

The nerves themselves, we know, are sometimes separately affected. Of this form of the disease, the *ischias nervosa*, as it has been called, is the clearest example. To determine whether the filaments of the nerve itself, or whether its covering texture, is the primary and legitimate seat of the disease, is a problem of some difficulty; but the former conclusion appears the most probable. I am strengthened in this opinion, from considering the sudden and *electrical*-like shootings of pain, which, I know from my own experience, often affect the

branches of nerves when under rheumatism*, even in the incipient state of the complaint.

Serous-membranes are liable to disordered action as a consequence of rheumatic inflammation affecting the contiguous fibrous texture. As a rare occurrence, the dura mater appears to become the seat of inflammation under acute rheumatism, and its contiguous serous membrane, the tunica arachnoides, becomes excited to increased action. I remember an instance of this kind, in which the symptoms were those of effusion, and the event was quickly fatal. It was under circumstances of previous quick transference of the inflammation of the limbs from one part to another; but the inflammation of the limbs did not cease with the new action of the brain. The patient was a young lady of delicate constitution, aged fifteen.

The pericardium also sometimes becomes the seat of similar morbid action, which I conceive is secondary, and induced by the inflammation which is affecting some part of the tendinous structure of the heart. Happily such an occurrence is very rare in acute rheumatism. I have seen one very clear example of this occurrence, in a man aged twenty-four. In this case also, the inflammation in the limbs had been quickly wandering, but did not become suspended by the internal affection. After many distressing symptoms, which had the duration of about fourteen days, the patient died. On dissection,

* In respect to my present observations on the textures which may be the seat of rheumatism, it must be allowed, as an apology for the conjectural manner in which I have treated the subject, that opportunities of ascertaining the morbid anatomy of the disease cannot easily be obtained.

recent layers of coagulated lymph were found lining the greater part of the pericardium, and also appeared partially on the surface of the heart. The pericardium was thickened, and contained $\frac{3}{4}$ of muddy serum*.

The diaphragm is occasionally affected in acute rheumatism, and is a severe modification of the disease. It appears to occur in proportion to the acuteness and violence of the attack, united with its disposition of quick transference from one situation to another.

THE SYMPTOMS.

RHEUMATISM, in its acute form, is distinguished by a great uniformity of symptoms. As attendant upon an inflammatory attack, and constituting what is commonly denominated the rheumatic fever, they are those which usher in the phlegmasiæ in general; and the most remarkable I shall now enumerate.

The attack is first announced by cold chills, succeeded by flushings of the face and partial heats; a sudden repugnance to food; general lassitude; and depression of

* This case is to be distinguished from the more chronic disease of the heart itself, which has been described, and illustrated with many interesting examples, by Sir David Dundas, in the first volume of the Medical and Chirurgical Transactions. In all these cases, the disease is represented as having succeeded one or more attacks of rheumatic fever. Of seven cases which proved fatal, six were inspected; and in all, the heart was uniformly found to be enlarged. I should rather be inclined to consider the general rheumatism of the constitution to be a predisposing cause of this disease of the heart, than to pronounce it rheumatism of the organ.

spirits. There is more or less of universal soreness and aching of the body; and the local evidences, which characterise the nature of the disease, appear in the course of twenty-four hours, more or less unequivocally marked. At first the lower limbs are usually chosen by the disease as the situation of the attack; and in general, the ankles and the knees become at once, or in very quick succession, affected. Upon examination of the limbs, we find, at the immediate onset of the attack, what is commonly called a swollen state of the parts. But this description is too indefinite. The increase of bulk does not, as in gout, and as in common inflammation, arise from sudden increased effusion into the cellular membrane; but from an inflammatory distension of the bursæ mucosæ, and of the sheaths of the tendons. Soon, in most situations, the blood becomes more determined to the surface, and a vivid redness of skin follows, arising from the presence of arterial blood in the extreme capillary vessels. Sometimes, however, the redness appears only in very small patches; or, is even altogether absent. This last circumstance is most to be observed when the bursal structure is alone affected. The patient is fixed almost immoveably by pain. Except in cases of the utmost urgency, he feels occasional ease when he is perfectly at rest; and describes a sense of heat and aching in the parts, rather than of severe pain; but on those attempts to move, which an excessive state of restlessness makes involuntary, or which may be required, such agony is produced in the effort from the propagation of pain along the whole course of the muscles, that the intense sufferings of the disease are at once portrayed.

A sympathetic fever of the inflammatory type quickly becomes established, and is indicated by a full and quickened pulse; a burning heat of skin; a fur upon the

tongue, at first white, but which quickly becomes a thick and brownish coat; excessive thirst; a costive state of the bowels; a scanty secretion of urine, which appears very deep in colour, and cloudy without a distinct sediment. With these symptoms, also pain of the head, and a ready tendency to delirium, occur.

It is the very character of rheumatic inflammation quickly to change its seat, and the abatement of inflammation and pain in one part does but prepare the way to the same symptoms in some new situation. This transference or alternation sometimes takes place with surprising rapidity. The smaller toes and fingers are the parts the least liable to this inflammation; but in a severe attack, scarcely any part of the tendinous and ligamentous structure is spared. The upper limbs become next affected; although, I should add, the order of the attack is chiefly to be referred to the particular circumstances of exposure, in which the body has been accidentally placed.

The pain produced by acute rheumatism is described as that, which, it is conceived, would arise from the gnawing of the part by a dog. A sense of burning heat, some throbbing, and pricking and shooting, are also felt; and if the diaphragm, or the tendinous portions of the intercostal muscles, or of any of the muscles of the chest, be affected, great difficulty of breathing takes place, and this is often severely spasmodic.

Irregular pyrexia is joined with irregular profuse perspirations, which take place as an apparent effort of Nature to give relief. These discharges from the skin are seldom productive of advantage, and very commonly serve only to increase debility. The anxious physiognomy of a patient under a fit of rheumatism marks the severity of the disease; and, together with this language of the features, there may be observed a peculiar relaxation of the

skin of the face, which is pale and flushed alternately, and for the most part bedewed with a greasy kind of moisture. In the same manner as in the gout, the pain is often alleviated in proportion as the inflammation appears external with redness on the surface. The most intense state of suffering is in the affection of the deep-seated parts, and before the blood is propelled into the cutaneous vessels.

The duration of an attack depends chiefly on the medical management which is adopted; according to which also, chiefly, it is favourably removed, or degenerates into the chronic form. Its critical termination is very commonly accompanied by a deposition of the lateritious sediment in the urine; or by a gentle diarrhœa; or by a general moderate perspiration.—These indications, and actions of crisis, also happen either separately or in conjunction.

SEQUELÆ.

THE sequelæ, or occasional consequences of acute rheumatism, are most remarkably seen in a permanent change of structure and function in the textures which have been affected with active inflammation, and are left in the state of chronic rheumatism. The bursæ mucosæ and the sheaths of the tendons are in this case the most common seat of complaint, appearing distended and much thickened, in the manner which has been before mentioned (p. 44) as consequential also to gouty inflammation.

It sometimes happens that a very troublesome disease of the bursæ remains, proving of difficult treatment. The appearance is chiefly that of exceeding distension, so that it seems like a bag of fluid. The part is occasionally tender

and painful, sometimes also joined with slight redness of the skin; but in general it does not, even in this state of tenderness, give sensible distress to the patient, except upon sudden motion, or more than ordinary exertion of the parts.

Of inflammation of the pericardium, and of the dura mater, as occasional occurrences in acute rheumatism, I have already spoken.

The debility which ensues after a long course of acute rheumatism, occasionally introduces some other constitutional disease, according to the particular tendency of the individual. Thus, consumption may follow; chorea sancti viti; intermittent fever; &c. The consequences which I now mention, I have myself witnessed.

PREDISPOSING CAUSES.

THE existence of *hereditary structure* may be assumed as an occasional predisposing cause of rheumatism; but it is obvious that the inference can be drawn only from general reasoning, and not from demonstration. We certainly see instances in which the disposition to rheumatism is strongly prevalent in the same family. Although a similarity of habits in regard to clothing, temperature of apartments, exposure, &c. may very reasonably be viewed as the chief source of the agreement in question, we should not, I conceive, exclude similarity of structure from its share in the explanation of the fact.

Age, Bodily Structure, &c.—The early years of infancy are exempted both from the acute and chronic rheumatism, and old age from the acute; but from ten years of age to fifty, it appears that persons of all structure, temperament, and habits, become almost indis-

criminally affected. A period earlier than the twelfth year, or later than the fiftieth, although not exempted from acute rheumatism, rarely affords examples of it. From each form of the disease, very fat persons are the most protected; and, vice versa, thin persons, and such as have a delicate thin skin, are to each the most liable. After the age of fifty, rheumatism most commonly appears in its chronic form.

Dr. Cullen has stated that persons of the sanguineous temperament are the most disposed to acute rheumatism; and of the propriety of this opinion I am well convinced: but, nevertheless, the observation just now mentioned, as a general position, is founded in truth.

Whatever causes induce debility, either general or partial, in the tendinous, ligamentous, or nervous structure, become predisposing to chronic rheumatism. Thus, a continued fever, in the general weakness which it produces, is sometimes an introductory source: and to this may be added the relaxation of the system and of the surface of the body, occasioned by the free employment of mercury. A strain or contusion occasionally lays the foundation of partial rheumatism, which takes place either in the acute or chronic form, but most commonly in the latter.

An unhealthy state of the digestive functions, deranging the nervous system, and producing irritation and debility, becomes the ground-work on which the disease sometimes establishes itself.

Either sex is indiscriminately liable to rheumatism; but in an extensive comparison, it has appeared to me, that men constitute the largest proportion; and, as it

would seem, on account of their being most exposed to wet and cold.

The *season of the year* is a predisposing cause, more or less active according to the degree of moisture and variable temperature which is prevailing. Authors have all concurred in fixing upon autumn as the most frequent season of rheumatism; for then, the apparel of summer is still worn rather carelessly, or with doubt as to the change required; and the weather of summer and winter constantly intermixes. In connexion with this remote cause, it is manifest, that habits of effeminacy very powerfully predispose the body to rheumatism. Warm apartments, and great warmth of clothing both by day and night, come under this head.

Excess, or irregularity in diet, is not to be accused as a remote cause, otherwise than in this general manner; —that if the individual have a constitutional tendency to rheumatism, it will be brought into action by a disturbance of the digestive functions; a diseased irritability of the system being thus induced.

Excessive perspiration, however produced, rendering the body more liable to the injurious effects of exposure to a damp and cold atmosphere, may be added as a very active predisposing cause.

EXCITING CAUSE.

THE influence of *variable temperature*, either generally or partially applied; and either through the medium of

wet, of cold air alone, or of both conjoined, appears to me the only *exciting* cause of rheumatism. In proportion to the previous debility of the body wholly, or of particular textures in part, will its effect be produced. In every climate the disease is found; but it occurs chiefly according to the degree of effeminacy in the customs of the inhabitants, and the variableness of temperature which prevails.

PROXIMATE CAUSE.

AN inquiry concerning the predisposition to rheumatism which exists in the constitution of some individuals, rather than to any other disease, involves the intricate question of *proximate* cause. Upon this point I shall be contented to say a few words. In reference to any of the phlegmasiæ, it may reasonably be assumed, that the particular texture, or part of texture, which becomes the seat of disease, under a general and common cause of injury, such as cold, is less strong in its organisation, and, consequently, less equal to maintain its healthy functions than other parts. In this local difference of powers, the predisposition to one or other of the phlegmasiæ may be stated to exist. In the most healthy person we cannot suppose that each separate texture is 'equally strong with the rest; for however this might be considered to be the case at birth, the various circumstances to which we are exposed, both of a partial and general nature, will forbid its continuance. The sensible difference of structure in a part possessing strength sufficient for the performance of healthy functions, and in one that falls short of this condition, cannot be demonstrated *à priori*. Even in parts which have been deranged in function, the

evidence does not always appear to us in the inspection of structure after death; and yet the general conclusion in question is not the less to be derived. We constantly see, in the ill consequences of exposure to wet and cold, that the weakest part of the body, unless it should have been protected by provident clothing, is the most visited by the disease which follows.

It appears to me that rheumatism in its *primary* character is rather a local than a constitutional disease. The fact, that some persons are constitutionally predisposed to it, is not at variance with the present position. The internal functions of the body are not necessarily first affected. The pyrexia which arises, as connected with the inflammation of the affected textures, is truly sympathetic. I would say that it is a disease of certain textures not demanding a specific state of constitution. The invasion of the disease has not, apparently, like gout, any necessary and essential dependance on the previous condition of the chylopoetic viscera, or on the healthy equal balance of the circulation. The predisposition which may exist in any individual is certainly increased by a derangement of digestive functions; or by any other error of constitution. It is in consequence of being involved with various states of constitution, that rheumatism is to be considered as a constitutional disease, rather than on account of its own legitimate and original character. In a true comprehensive pathology, no disease can be pronounced so local as to be entirely independant of the constitution. It is only in an inanimate machine, that a part of the structure may be injured without affecting the use and convenience of the whole. The intimate connexion which all local diseases possess with the constitution, is an important part of pathology, and has been ably illustrated by Mr. Abernethy. It will be seen that it is in the analytical view of

the subject only, that I consider rheumatism, in its primary character, a local disease. In gout, the affection of the constitution is the antecedent, and that of the external parts is the consequence; and this is the essential part of the pathology of gout: but in rheumatism, this order of the disease is reversed; as relates to the irritation and derangement which belong to the constitution.

The influence which is again reflected to the affected parts from the constitution, according to the particular condition in which it may be, is equally seen in inflammation produced by mechanical injury, as in rheumatism.

Rheumatism, like gout, affects those textures which are scarcely susceptible of the suppurative process from inflammation; but when the diseased textures are of the synovial kind, as the sheaths of tendons and the bursæ mucosæ, an increased secretion is produced, which, together with the preternatural fulness of the surrounding blood vessels, causes more or less of distension and swelling. Whether the inflammation of rheumatism be truly common inflammation, but differing in the symptoms which arise, both on account of the nature of the textures which it affects, and the general mode of operation in the agent, *cold*, in some way applied, producing the disease; or whether it be one distinctly *sui generis*; is a question which may be thought to admit of some argument. The remarkable manner in which rheumatic inflammation quickly transfers itself from one part to another, is an obvious ground of distinction from the phenomena of inflammation of similar textures, produced by mechanical injury.

From a careful review of the whole subject, I am therefore induced to define rheumatism, *A peculiar species of inflammation, affecting parts which have a fibrous texture, and most frequently the synovial membranes, producing much sympathetic irritation in the constitution, and fever of the inflammatory type.*

RATIO SYMPTOMATUM.

THE explanation of the peculiar phenomena of rheumatic inflammation is chiefly derived from the nature of the textures which it affects; and any difference of external characters, which it assumes in its different modifications, is to be traced chiefly to the particular texture affected by the disease. The locomotive organisation of the body comprehends so many distinct parts which are exercising similar kinds of function, that under the general inflammatory diathesis of the system, one part will scarcely be affected by rheumatism without other parts participating. Hence the shifting and continuous nature of acute rheumatism; its spreading suddenly from one texture to another in the same limb; or transferring itself with equal rapidity to another part of the body; observing usually the general law of affecting some branch of the muscular or articular structure. This propagation of pain and inflammation is found to take place most remarkably in tendinous parts; and in the ligamentous texture next in degree.

The most fixed locality of symptoms appears in the bursæ. This order of the phenomena also prevails in chronic rheumatism; with the exception, that in this latter species of the disease, some distinct nerve, or branches of nerves, is frequently the texture which becomes affected.

DIAGNOSIS.

THE discrimination of rheumatism from gout has already been pointed out at length; and as this part of the

subject relates to *acute* rheumatism, it appears to me all that need be stated.

OF THE TREATMENT.

I shall take a cursory view of the principal remedies which are in use, in the treatment of acute rheumatism.

General Bleeding is a remedy of great importance in this disease, but requiring very careful management. A degeneracy into chronic symptoms is in no way so readily introduced as by an injudicious employment of the lancet. If the diaphragm or intercostal muscles be affected in the course of the disease, so that the respiration is performed with extreme difficulty and pain; or if any internal viscus be seized with inflammation, while the system is under the influence of acute rheumatism; a prompt detraction of blood is indispensable; and its repetition also will be suggested by the usual principles of practice. If one of strong muscular fibre and of sanguineous temperament be seized with acute rheumatism in full health, bleeding at the commencement of the attack is a measure of the utmost necessity and value; and the propriety of its repetition will be clearly indicated, both by the good effects which it may have produced, and by the urgency of the subsequent symptoms. When its employment is proper, its controul over the violence of the disease is more immediate and effectual, than that of any other remedy. It should be exercised with great circumspection in persons of languid constitution, in whom the circulation is excited rather by pain and general irritation, than by the true inflammatory diathesis.

Sydenham, in his first Essay upon Rheumatism, ad-

vocates the propriety of bleeding as the chief remedy which should be employed in the cure. He advised it on the grounds of believing the disease to be inflammatory, "as indicated by the resemblance of the blood which is taken away in pleurisy;" and from his fondness for the humoral doctrines. He speaks of febrile and morbid matter derived to the limbs and causing the symptoms. In a subsequent short treatise "of the epidemic diseases, from the year 1675 to 1680," he regrets the practice of taking away blood so freely as he had before recommended; and in a case in which he made a comparison of his modes of treatment, concludes with stating his change of opinion, according to the following translation:

"I ordered the patient to live upon whey only for four days, after which I allowed him white bread, besides the whey, for his common food, namely, once a day, instead of a dinner, till he was recovered. Contented with this slender diet, he persisted in it eighteen days; only towards the latter end I permitted him to eat white bread also at supper. He drank a gallon of whey every day, which afforded him sufficient nourishment. At the end of this term, when the symptoms were gone off, and he got abroad, I allowed him to eat flesh, as boiled chicken and other food of easy digestion; but every third day he lived upon whey, only, till at length he recovered; and by this method escaped the inconveniences I mentioned above, which had been very troublesome ten years before, when frequently repeated bleeding was used by my order for his cure." On a subsequent occasion, however, Sydenham still appears to shew his partiality to the practice of bleeding.

Sir John Pringle, in his account of the diseases of the army in several campaigns, represents the frequency of acute rheumatism, and his successful treatment by re-

peated bleedings. He says, "that when the rheumatism was attended with an inflammatory swelling of the joints, sweating was improper, and the cure was only to be obtained by repeated and almost daily bleedings, till the patient was without fever, and the pains were either entirely removed, or became easier. And in this course we may proceed the more boldly, as those that are subject to the distemper are generally in the vigour of life, and are either plethoric, or at least able to bear great evacuations. Add, that frequent bleedings weaken the body perhaps less in this disease than in any other*."

Dr. Cullen considered blood-letting "to be the chief remedy of acute rheumatism" (par. 463). He remarks, "The blood ought to be drawn in large quantity, and the bleeding is to be repeated in proportion to the frequency, fulness, and hardness of the pulse, and to the violence of the pain. For the most part, large and repeated bleedings during the first days of the disease, seem to be necessary, and accordingly have been very much employed: but to this some bounds are to be set, for very profuse bleedings occasion a slow recovery; and if not absolutely effectual, are ready to produce a chronic rheumatism."

Even Dr. Haygarth, who wrote on the subject of rheumatism, as he states, "for the principal purpose of recommending the Peruvian bark in preference to all other remedies," mentions bleeding as one of the evacuating remedies occasionally necessary to be premised.

The best practical authorities therefore, we see, have sanctioned the use of general bleeding in this disease; but we observe at the same time the doubts which they have

* See page 182, 8vo. edition.

thrown on the practice, as forming a general principle of treatment.

I have already concisely stated my own view of the question; and shall conclude with expressing, that if the case require this remedy, it is very important to use it early in the disease; and that even under very urgent symptoms, unless indeed affecting some internal parts of structure, we should scrupulously hesitate in its repetition, whenever we see that relief is not in any degree afforded by the first copious depletion.

Emetics.—Dr. Haygarth states, that it was his usual practice in acute rheumatism, as a preliminary to the use of the bark, “to give either the antimonial powder or tartarised antimony, generally the former, till the stomach and bowels are sufficiently cleansed.” Emetics administered at the commencement of the attack are useful on the same principle, that in the first formation of a febrile disease of every description, the action of vomiting, by its influence on the circulating system, and by the complete relaxation of the skin which it produces, moderates the force of the symptoms which are forming; and it further renders material service as an evacuant. If the patient be seized in consequence of exposure, shortly after some convivial occasion, on which he has indulged in improper diet, the present remedy should not, on any account, be neglected. The use of an emetic, if at all employed, should obviously, for the reasons I have stated, be the first remedy of our choice, when bleeding is not required; and otherwise, the second.

Cathartics.—The advantage of making a detraction from the general circulation, by the channel of the alimentary canal, is no less remarkable in rheumatism, than

in every other inflammatory disease. In proportion as we pursue this practice upon a continued principle from day to day, do we obtain its good effects in acute rheumatism; the circulation becomes moderated; the inflammatory diathesis subdued; and the absorbent system is excited to increased action. Hence, we powerfully promote the removal of those excessive secretions of the synovial membranes, which have been already described as causing the distension and impeding the motion of the affected parts. A saline purgative, administered in small doses, and at repeated intervals, is the most advantageous. The action of the kidneys becomes excited, and the cure by this effect is assisted. Calomel in occasional doses, for the purpose of purgative operation, is unquestionably a valuable remedy in the acute rheumatism; but if employed with such frequency as to produce mercurial fever, its effects appear to me always sooner or later injurious. In very numerous instances in which I have witnessed the treatment to consist in the use of calomel, antimony, and opium, given in repeated doses at short intervals, however favourable their immediate operation may have been, the final result has appeared to be an increased susceptibility of the patient to relapse.

To fulfil the views just stated, we may choose, according to circumstances, the use of a saline purgative in conjunction with our sedative, at short and regular intervals; or, having prescribed calomel and the antimonial powder, with or without the colocynth extract, at bed time, we may administer a liquid purgative in the morning early, and equally observe due and regular intervals in the use of other medicines both by day and night.

Sudorifics.—The sudorific plan of treatment not un-

frequently disappoints our expectations, so as to aggravate rather than relieve the symptoms. Even when most successful, it is attended with the ill effects of producing much debility, and increasing the sensibility of the surface; so that, for a considerable time, almost any degree of exposure is hazardous. A moderate relaxation of the skin is free from this objection; and the employment of antimony in conjunction with calomel and purgatives, or of antimony or ipecacuanha with opium in moderate doses, so as to produce a favourable determination to the surface, is entitled to our fullest regard.

In the distinct division of the purgative and fever medicines, as just now suggested, a draught containing the nitrate of potash, camphor julep, or water simply, a portion of either the vegetable or volatile alkali neutralised, accordingly as a sudorific of mild or stimulating properties may be required, and some tartarised antimonial wine, usually proves very useful; but whether the juice of fresh lemons, or acetous acid be employed, it is sometimes expedient to prescribe its use in the state of effervescence with the alkali, and especially when the stomach is in a weak and irritable state.

Sedatives.—From the free administration of opium, when at the same time correctly fulfilling all other indications, I have invariably derived the most satisfactory results. In the recommendation of this important medicine, I think it necessary to dwell upon the necessity of a due attention to the action of the bowels, the kidneys, and the skin, as essential to the advantages which it is capable of affording. Also, I must not lose sight of the objection with regard to the inflammatory diathesis, which sometimes prevails so strongly, that its previous reduction or abatement is indispensable to the employment of opium in any form.

When, however, the inflammatory diathesis is slight, and more especially when the increased action of the heart and arteries arises chiefly from the irritation of pain, we should consider that the use of opium, guarded, I again say, by the influence of other medicines, is our most sovereign remedy.

The pulvis ipecacuanhæ compos. is a preparation very much deserving our choice; but on comparing the results of my experience, I am most induced to recommend the use of crude opium with the pulvis jacobi, in the proportions expressed at p. 224. In directing so small a dose of the antimonial powder, it is to be considered that the attention to the action of the skin, for which the use of this medicine is chiefly intended, is more particularly to be studied in the prescription of the fluid formulæ.

Finally, I have to observe of the use of opium, that, under correct indications for its employment, no further guidance is required for the freedom of its dose, than the urgency of pain and high constitutional irritation; such being the symptoms for which we have recourse to the powers of this friendly agent.

Peruvian Bark.—I have allotted a separate discussion for the virtues of this medicine as a remedy in acute rheumatism, from the remarkable credit which it has acquired with some physicians. It is a curious circumstance, however, that Sydenham considers the free use of the Peruvian bark as one of the causes which predispose to that species of rheumatism, which he designates the *Scorbutic*. But on reading his account of its symptoms, I am impressed with the idea, that the pains which he describes as attendant on the disease, were sympathetic only from a morbid state of the digestive organs; in the production of which, a continued

administration of the Peruvian bark might certainly be concerned, as tending to induce or increase visceral obstruction.

Dr. Haygarth, to whom I just now referred, after stating that he first made trial of the Peruvian bark on the authority of Dr. John Fothergill, and next taking notice that Dr. Saunders had for many years recommended it both in his lectures and publications, proceeds with the following recital. "To sum up the whole in a few words. After the stomach and bowels have been sufficiently cleansed by antimony, I have, for many years, begun to order the powder of the Peruvian bark in doses of gr. v, x, or xv, every two, three, or four hours; and if this quantity has a salutary effect, it was gradually increased to gr. xx, xxx, or xl, with sedulous attention never to add more than what perfectly agrees. It has generally been taken in milk, mint water, or the decoction of bark." He mentions a few exceptions to his success, which otherwise was very uniform. He declares that "the pains, swellings, sweats, and other symptoms of inflammatory fever, manifestly and speedily abate; and gradually cease, till health is perfectly restored." He further observes, "Another circumstance merits great attention. When the rheumatic fever has been treated by bleeding, leeches, sudorifics, &c. it is well known that pains of the diseased joints and muscles often afflict the patient for many months, or even years. In my Clinical Reports I find no instance of this kind, and have reason to think that bark entirely prevents this cause of the chronic rheumatism, as a consequence of the inflammatory fever."

His concluding panegyric is very strong: "Except mercury in the syphilis, there are few or perhaps no ex-

amples, where a remedy can produce such speedy relief and perfect recovery in so formidable a disease. For many years I have been thoroughly convinced, that the Peruvian bark has a much more powerful effect in the rheumatic than in any other fever; and that it does not even cure an ague so certainly and so quickly*.”

On reading this statement, who would not imagine that a specific cure for the acute rheumatism was here discovered? Experience however shews, in this as in many other instances, the extreme difficulty of collecting such sure evidences in medicine, that certain rules of practice may be laid down. I have repeatedly followed the authority which I have quoted, in administering the bark in the first stage of the acute rheumatism, its use having been premised by due evacuations; but I do not remember more than one case, in which I found it successful. *A priori*, we could not expect it to be a medicine otherwise than injurious, when a sympathetic inflammatory fever is present; and all the secretions, except that of the skin, which is irregular, are more or less impeded.

I should venture to state, from such experience as I have had, that the Peruvian bark, either administered as Dr. Haygarth directs, or in decoction with tincture and sulphuric acid, is often a very suitable and valuable medicine to be employed when the convalescence begins; namely, when the fur of the tongue is fast contracting its edges, and has an appearance of detaching itself; when the urine is of light specific gravity; the state of the bowels is natural; the skin is relaxed; and simple debility prevails; but that under the opposite circumstances it will, I fear, entirely disappoint our hopes of its asserted specific powers.

* A Clinical History of Diseases, &c. p. 89.

Regimen and Diet.—It is the usual practice in the management of this disease to place the patient between blankets, or to cover him with the warmest clothing. It is true that it forms a part of the treatment by sudorifics, and may appear essential to it; but also, whatever medicines may be administered, this heating regimen is not only too much used by the patient and the nurse, but even by the prescription of the physician. There is no conclusion of which I am more convinced, than that this is a very erroneous proceeding. I have on several occasions directed the patient to be gradually removed from this *hot-bed* of perspiration, and to be placed between well-aired sheets, using only the lightest covering with which he could feel comfortable. I have also kept the chamber freely but cautiously ventilated; adding at the same time the comfort of a good fire, as the season of the year the most frequently giving rise to this disease certainly requires. I have been delighted with the expressions of relief which the patient has always acknowledged from such change of treatment; and I am well convinced, that the method of evaporation by the skin, as arising from forced perspiration, is in most instances only calculated to aggravate the inflammation and pain; to prolong the disease; increase debility; and, above all, to induce the continuance of the disease in its chronic form.

The diet should, invariably, be of the lightest kind, and chiefly consist of light diluents. The whey, so justly praised by Sydenham as all the support required in the most acute stage, with tea, thin gruel, and the grateful addition of subacid fruits, as oranges and grapes when in season, include all that need be directed. Boerhaave relates, that when he laboured under the most exquisite tortures of the rheumatism, he supported himself entirely on milk whey for twelve days. When the symptoms

subside, it is of infinite importance that the return to solid animal food should be very gradual; and it ought not to be permitted while any degree of inflammatory tendency remains. I have seen all the symptoms reproduced by a premature meal, consisting even of chicken: but a light broth, fever being entirely absent, does not carry with it this hazard.

OF LOCAL TREATMENT.

IN co-operation with these constitutional remedies, I have now to consider the question of local external treatment. Upon similar principles of practice to those which we exercise in treating common inflammation, when it affects the textures which are not fibrous, as glands, cellular membrane, skin, &c. I contend that the abatement of local inflammation in rheumatism, by judicious means of evaporation, is an object of great importance, and of rational intention.

The easy spontaneous transference of rheumatic inflammation from one part to another; and the fact, that sometimes this transference suddenly takes place to internal parts (the diaphragm more especially), entirely forbids the application of direct cold, as a mode of lowering the inflammatory action by its sedative powers. Against the practice of evaporation by means of excessive perspiration, I have already advanced my objection. The middle line of treatment offers itself to our choice, as a method free from the obvious disadvantages which are attendant on the employment of either extreme heat or extreme cold. A moderately cool air of the apartment, and an equality of its temperature as much as can be produced; cool or tepid drink; as light covering of the

bed as is consistent with the comfortable feelings of the patient in regard to warmth; are means which comprehend a great part of the treatment now in view. Lastly, I have to mention, with earnest recommendation, the constant employment of the tepid evaporating lotion (p. 254) to the inflamed parts, precisely according to the method before described. The good effects of this remedy, judiciously conducted, are often truly surprising. But it is to be understood, that, as with respect to gout, so in rheumatism, local evaporation is to be considered only as an auxiliary to the general treatment. In this character, I repeat, it is a valuable remedy; and when employed as a subordinate agent to the constitutional remedies already described, it is, according to my experience, equally safe and useful. In proportion as rheumatic inflammation is local, we are made sensible of the beneficial activity of the evaporating treatment. When the inflammation wanders quickly from one part to another, almost eluding our pursuit, we ought probably to rely distinctly on constitutional means; and certainly to make our local treatment both more subservient and considerate.

In some cases of severe acute rheumatism in which the inflammation has affected equally the upper and lower limbs; and in other cases, in which the acute inflammation has been only partial; I have seen the best effects produced by means of this lotion. One lady, who had found any attempt to move her limbs attended with the greatest agony, in a few hours after its free application, was enabled to walk a little about the room.

Another patient, a very delicate female, suffering much torture from the inflammation affecting her knee, which had been produced by a partial exposure to cold, so much extolled the effects of the lotion, that she thought it had sufficient power to accomplish her cure.

The Convalescence.—The use of a bark tonic, conjoined with any of the mineral acids, but usually the sulphuric as the most favourable, is at this period, in most cases, of unquestionable propriety, and is usually attended with the best effects.

The stiffness, aching, and debility of the limbs, which follow the acute state of pain and inflammation, yield only to active exercise and friction. I have often directed the patient, when convalescent from acute rheumatism, to overcome the seeming incapacity by strenuous exertion, and to walk several miles in the day; beginning of course with moderate attempts. The best results have followed; and, when the season of the year or weather allow this remedial method, it cannot be too strictly enjoined; so that particular contradictions do not forbid its practice.

The Prophylaxis.—When we reflect on the cause which alone excites this disease, namely, the influence of vicissitude of temperature, and that some accidental or constitutional relaxation of a part or the whole of the body is the most frequent state of predisposition, on which it makes its invasion, we must at once be convinced how much the prophylactic management consists in the use of those means which may invigorate the frame, and lessen its susceptibility to the impressions of variable atmosphere. With a view to this important point, I advise that every morning the patient should wash his head and neck, by means of a coarse towel, with cold water; and sponge the feet, also, every morning, with water just freed from unpleasant *chill* by the addition of warm water. I have seen abundant proof of the preservative influence of this method, steadily pursued. One gentleman, who a few years ago was affected with

rheumatism on every exposure to a moist atmosphere, or a cold east wind, assures me, that since an attention to this plan, which he adopted at my persuasion, he can bear very free exposure with impunity. I could enumerate many other instances illustrative of the value of the practice. Sea bathing, or, for some individuals, even the cold bath, during the summer season, is a prophylactic remedy of great value. In the general plan of effecting a hardihood of constitution, small heated rooms, the habit of a *warm bed*, and similar indulgences, must be avoided. In this precarious climate, the useful rule of wearing flannel next to the skin during the greater part of the year, and for persons who are delicate and very susceptible *unremittingly* through the year, is too well known to require particular injunction.

An occasional exception, however, occurs to the propriety of this prudential method, except in the very severe season of the year. It now and then happens, both with gouty and rheumatic patients, that the wearing of flannel next the skin, and of warm stockings, produces so much perspiration upon ordinary slight exertion, that the consequent evaporation causes a sense of chilliness, and really renders the whole body more susceptible to a change of atmosphere, than when light clothing is used.

OF CHRONIC RHEUMATISM.

THE symptoms of chronic rheumatism have a much less definite and evident character than those of the acute. It is only in this species of the disease, that we find the nerves separately affected. This distinction is most frequently exemplified in the rheumatism of the sciatic nerve. Any other principal nerve and its branches may be affected in the same manner. In correspondence with this state of complaint, the spasmodic and painful action of the muscles, to which the branches of the affected nerve are distributed, is to be remarked. It usually happens indeed that pain is felt only upon motion. This circumstance is most remarkable when the nerves of the lower limbs are affected. The patient very frequently will be able to walk for a short distance without inconvenience; but either gradually or suddenly he is seized with pain and lameness, sometimes with sudden cramp, and anxiously seeks to terminate his difficult exertion by rest. In a recent attack of this kind, it often happens that the nerve is affected with inflammatory action; in which case the pain continues with little intermission, and is of the piercing, shooting kind. This is usually attended with a soreness of the parts to pressure, which is chiefly to be distinguished in the track of the nerve which is principally affected. In a more chronic or passive form of the complaint, notwithstanding the intensity of occasional pain, and various mixed sensations of pricking, burning, and numbness, the muscles and integuments very commonly allow of the strongest pres-

sure without inconvenience, unless indeed it be made upon the immediately affected branch or branches of the nerve. In cases of long standing, the symptoms are such as indicate all absence of inflammation, and put on only the marks of relaxation. In this example, the limb is readily sensible to the variations of atmosphere; is occasionally affected with much coldness; invariably suffers an accession of pain from over-exertion; the muscles are deficient in their natural firmness and size; and the limb in this condition exercises an active sympathy with any disorder of the digestive organs, whether accidental or continued; and is even powerfully influenced by causes which affect the mind alone.

If chronic rheumatism affect the bursal, tendinous, or ligamentous textures, we find that two distinct states of inflammation may exist, although general sympathetic fever may not be present; namely, the active and passive: the locality and slighter degree of the disease still distinguishing it from the constitutional attack which we designate the rheumatic fever, already considered. The external characters which appear, are very characteristic of the particular texture which is affected. In some instances, we see in the same individual case, the bursal, ligamentous, and tendinous textures, all under the influence of the disease; but in others, the affection of one part only is remarkably distinct. When the deep-seated ligaments are the seat of complaint, no external morbid character is apparent; but the indications of the disease when the superficial ligaments are affected, may usually be detected by manual examination; and those which appertain to the bursæ and tendons, are rendered clear by the evident increased distension, and by the tenderness of the parts produced on pressure. The muscular aponeurosis and periosteum, also, when affected with

rheumatism, become more or less sore to the touch. In that common form of complaint, familiarly termed the *stiff-neck*, we have a clear example of tendinous and aponeurotic rheumatism; and also of the active state of inflammation appearing distinctly as a local complaint.

The synovial membranes of the joints appear to be liable to rheumatic inflammation; or, probably, I should give a more correct statement by saying, that certain individuals possessing the rheumatic diathesis are susceptible to an inflammation of the synovial membranes of the joints, from exposure to wet and cold. It is not a very common occurrence, and, I believe, happens only to those persons who have a specific predisposition in these textures to be so affected. I conceive, that in most of the cases of this description, the constitution may be possessing a scrophulous tendency. Mr. Brodie, in his very instructive "*Pathological Researches, respecting the diseases of the joints**," has referred to rheumatism as one of the causes which induce inflammation of the synovial membrane. He speaks of the disease as sometimes acute, but more frequently as assuming the chronic form; "that it takes place from various causes, but in most instances from the application of cold, which explains why it is more liable to occur in the superficial joints, such as the knee and ankle, than in the hip and shoulder, which are defended by a thick mass of soft substance, from the influence of the external temperature."

In connexion with acute rheumatism, or rather with that continued form of the disease which partakes of the active and passive character of symptoms in an irregular manner, we meet with cases of painful inflammation of

* Medical and Chirurgical Transactions, vol. iv.; and his extended work lately published.

the eye, exhibiting much peculiarity in the nature of the symptoms. My friend, Mr. Wardrop, in an ingenious paper on the subject, which he has just contributed to the Medical and Chirurgical Society, has defined this disease, "rheumatic inflammation of the eye." The most leading feature of the complaint, is, that the pain and inflammation of the eye very much alternate with the same symptoms affecting the limbs; and require similar constitutional means of treatment: but I must add my conviction, that local remedies are equally necessary in this as in any other species of inflammation of the eye. In such cases as have come under my observation, together with the use of leeches, the free employment of poppy fomentation has afforded much relief, and has been far more successful than any cold applications; which I have seen to disagree. Mr. Wardrop considers this particular species of ophthalmia "to be chiefly seated in the sclerotic coat;" and offers his grounds of diagnosis.

SEQUELÆ.

I HAVE already stated that chronic rheumatism itself is a frequent consequence of the acute, and have defined the usual characters of this state of the complaint; but we have also to consider that chronic rheumatism often originates as a distinct form of the disease, and is not only, as its name (from χρόνος time) implies, most tedious in the duration of the sufferings which it produces, but further leads to very important changes of structure in the particular textures which it affects. I shall, for the present, attempt only a brief sketch of the morbid anatomy of chronic rheumatism; but it is undoubtedly a subject very much demanding further and accurate investigation.

The consequences of the disease, which are most obvious in the living subject, appear in the bursæ, tendons, and muscles. In very debilitated persons, the bursæ are often enlarged and soft, appearing like bags of thick jelly. For the most part, however, they are hard and unyielding; and the smaller ones, especially, become much indurated. The tendons are thickened and knotty, as before described. The muscles are wasted, and are either flaccid and very evidently weak; or appear hardened and rigid, with a thickened state of the aponeurotic covering. The ligaments undergo remarkable changes, becoming thickened and rigid; and from their consequent loss of elasticity, produce great lameness. The occasional diseased action of synovial membranes has already been mentioned; but I have to add, that, without the distinct marks of preceding inflammation, a jelly-like effusion into the joint is occasionally found as a sequel of the rheumatic action. It is highly probable that the nerves suffer, from rheumatism, some change of structure, either in the sheaths which inclose their filaments, or in the filaments themselves; but the opportunity of proving this conjecture is equally rare and difficult. The conclusion may be presumed from the loss of nervous power almost amounting to paralysis, which occasionally affects rheumatic limbs.

Mr. Stanley, of St. Bartholomew's Hospital, has favoured me with an interesting account of a dissection of the joints of a man apparently between forty and fifty years of age, in which the following very remarkable appearances were found. He was unacquainted with the history of the case; and therefore I cannot offer the statement as an example of the consequences of rheumatism; but the narrative is too curious to be omitted.

“The articular cartilages in almost every situation ex-

hibited a perfectly white surface, such as would be produced by a very thin layer of plaster of Paris spread over them. In some of the joints, a small quantity of the white substance was found in a fluid state within the capsules, which circumstance rendered it probable, that the white matter had been in every joint originally deposited loosely within its cavity, and that, mixing with the synovia, it had become smeared over the cartilages, giving to them a white covering. Almost all the joints were thus affected; those of the extremities in the greatest degree. Even the articular cartilages belonging to the extreme joints of the fingers and toes were perfectly white. Around some of the joints of the toes, the same kind of deposition had taken place into the cellular substance, externally to the joint. By chemical analysis, the white matter was shewn to be carbonate of lime. Specimens of the joints are preserved in the museum of the hospital."

What has already been observed on the subject of the *remote causes*, as connected with acute rheumatism, may be applied to the chronic.

THE DIAGNOSIS.

THIS question is to be considered separately and at some length.

Of the discrimination of chronic rheumatism from chronic gout; and of the distinguishing characters of that disease, which has been called by Dr. Haygarth, nodosity of the joints, I have formerly spoken.

Rheumatism affecting the lumbar muscles, called lumbago, is distinguished from nephritis, by the evident increase of pain in the loins, in the motions of the body, and more especially in moving from the horizontal pos-

ture; and further, by the absence of the distinct nephritic symptoms, which are well known. The rheumatic affection of the tendinous fibres and aponeurotic covering of the muscles immediately contiguous to the kidneys, is less easily discriminated; because the local uneasiness produced is in a great measure common to each complaint. I have always found that this species of rheumatism is increased by the heat of the bed. The locality of the affection; and the absence of those symptoms which point out distinct irritation of the kidneys, as pain in the course of the crural nerve, with some attendant sympathies; and of the calculous sediment in the urine; will guide us in this diagnosis.

When the mucous membrane of the kidney is affected with inflammation, either chronic or acute, a pain at the upper part of the loins is felt more or less severely, and there is difficulty in raising the body from the horizontal posture; there is pain also in progressive motion, which arises chiefly from the action of the *psoæ* muscles. This state of complaint is distinguished from lumbago by the attendant symptoms; as sickness and much irritation of stomach; irritability in the functions of the urinary organs; and a deposition of mucus in the urine, usually in a copious quantity.

A pain between the scapulæ, or of the shoulder at the head of the joint, is sometimes confounded with rheumatism, when really proceeding from the influence of disordered digestive functions. The ambiguity will be removed by careful investigation of the state of the several secretions; by the appearances of the tongue; and by the character of such dyspeptic symptoms as may be present.

As depending on a morbid state of the digestive organs, and consequent general irritation, we see that wandering

muscular pains, or even transient pains affecting other textures, are occasionally described as rheumatic, which should rather be considered as the pains of sympathy; and in the investigation of such cases, the observations just now made are also applicable.

Rheumatic pains, which are produced by the employment of mercury without sufficient caution with regard to exposure, are of very common occurrence, and are readily traced; but a perplexity in these cases often arises in determining whether a *syphilitic* cause be not also present. It usually happens, however, that some characteristic symptoms, which become very clearly instructive upon investigating the history of the case, are associated with chronic pains of the present description. The periosteum of certain bones, as the os frontis, or tibia, or ulna, is in these cases almost with certainty affected with extreme tenderness on pressure. Very commonly, also, there is more or less of periosteal thickening, and especially at the middle of the anterior angle of the tibia, called the *shin*. I have already, at p. 344, mentioned some other points of discrimination.

Those irregular pains and spasms, which attend the commencement of a disease in some part of the vertebral column, claim a very careful distinction from chronic rheumatism; and I may also add, from those pains which receive the more *convenient* denomination of nervous.

I have once seen an aneurism of the aorta mistaken for lumbago. The patient was a young man of healthy appearance, between thirty and forty years of age. The symptoms came on very gradually. He first complained of pain in the upper part of the loins, which caused him to walk as a person affected with lumbago. It was a constant pain, but was always increased upon an action of the bowels taking place. After a time, the appetite became

affected, and there was loss of flesh and strength. The symptoms of pain and difficulty of taking exercise increased. The disorder was treated empirically as for lumbago. At length, a strong pulsation was accidentally discovered in the abdomen, just above the umbilicus; and a consultation being held, the case was pronounced to be an aneurism. Need I add the details of the certain melancholy result! Upon examination, an aneurismal tumour was found upon the aorta, between the superior and inferior mesenteric arteries. It had ruptured, and the hæmorrhage had taken place partly into the cavity of the abdomen, and partly into the duodenum.

Some painful affections of the nerves and the connected muscles occur, which, without attention, may be confounded with rheumatism. This difficulty will only occur, when the nerves belonging to muscles, either in the upper or lower extremities, are the parts affected. In such cases, it commonly happens that the influence of cold has not been the exciting cause; which, as I have stated, is, invariably, the parent of rheumatism. In the present form of complaint, we shall also find that the functions of some one or more of the principal constitutional organs have been previously much deranged; and that the painful state of nerves and muscles is sympathetic.

TREATMENT.

In my theoretical and practical *analysis* of rheumatism, I shall lay down the following principles of classification for the investigation of each individual case; proposing this as a method to ascertain the various sources of a modification of the symptoms, and of the appropriate treatment consequently required.

1st. The age; general structure; original temperament and constitution.

2dly. The acquired state of constitution from habits of living, in regard to diet; clothing; hardihood, or effeminacy; and modes of exercise.

3dly. The mode in which the exciting cause has been applied; the nature of any injurious remote causes; the accidental state of the constitution from any other disease; and especially the influence of a morbid state of the digestive organs.

4thly. The particular part of structure affected; whether as being ligament; aponeurosis; tendon; bursa; nerve; periosteum; or other texture; and the morbid state of its organisation.

5thly. Climate; season of the year; locality of residence.

To detail the diversity of treatment, which has been proposed by the regular part of the profession for the relief of chronic rheumatism; and to enumerate the very names of the boasted remedies of the empirics, would itself form no inconsiderable history of medicine. This disease we may believe to have been almost co-eval with man; and to have been the occasional consequence of exposure to atmospherical changes, in the rudest states of society. If, therefore, the experience of ages cannot offer us any certain means of remedy for the evil, it is at least a proof how much remains to be accomplished in this inquiry. We find, it is true, in authors, various medicines recommended as specifically suited for rheumatism; and the patient often undergoes such routine of treatment on true empirical principles.

It will, perhaps, be said, that chronic rheumatism is a specific disease, affecting all individuals alike in a general manner; and subservient therefore to plain and uniform

principles of treatment. Now it appears to me, that no two cases can admit of the same treatment; and that, on the contrary, according to various causes of modification, it must differ more or less in some important point.

The first, second, and fifth heads of remote influence are the most simple grounds of observation. The third is in part attended with much obscurity; but embraces such important points, that without giving them a full consideration in every case, we are not entitled to the expectation of a successful result.

The fourth head comprehends causes of peculiar symptoms and of relative treatment, which, I conceive, deserve much more attention than they usually receive.

In the brief consideration of my subject, which I purpose now to offer, I shall first discuss the treatment of the rheumatism of the nerves.

If the patient be seized in a sudden manner after exposure to cold, and symptoms arise which indicate that the nerve is in a state of inflammatory action, those means which are suited to the reduction of inflammation are to be employed with more or less activity. But it is usually required that the abstraction of blood should be made as contiguously as possible to the part affected. For example, in a newly formed sciatica, attended with local indications of inflammatory action and with sympathetic fever, cupping at the hip, and blisters afterwards applied, will, in conjunction with purgatives and other suitable medicines, produce the best effects.

If, however, the disease have been gradually produced; or from the peculiar temperament of the individual and his state of constitution, passive symptoms only appear, and the healthy functions of the nerve seem disturbed without inflammatory action, we are to consider whether we shall select the use of medicines which sti-

mulate, or of those which act as sedatives. For this guidance, we must investigate the state of the constitution; and seeing that the affected nerve (for I select the sciatic as the most remarkable and by way of example) is a part of the brain and nervous system, we must observe its state of function, not merely as the seat of disease, but as a material index of general information. What are the causes which most produce the paroxysm of pain, and in what degree do they act? Is the action of the nerve much influenced by the mind? by diet? by the state of the atmosphere? or by bodily exertion? What is the state of the digestive organs, as indicated by the various symptoms which I have so often mentioned? Is general debility, or universal muscular relaxation, present? Do these causes exist more or less in conjunction? Or, do all the functions of health proceed regularly, leaving the rheumatic nerve alone in fault? Of what duration has the complaint been?

It is of great importance that effective treatment should be adopted in the commencement of a case of this kind, and steadily pursued; for a neglected rheumatism of the nerves constitutes one of the most difficult chronic cases in medical practice.

The stimulant method of treatment comprehends the use of electricity; rubefacient embrocations; irritating plasters; the application of heat through the medium of dry substances, as salt inclosed in flannel; local vapour; the use of the flesh-brush, or strong hand-rubbing; and violent exercise: with the internal employment of gum guaiacum; bark; æther, the sulphuric or acetic; the volatile alkali; turpentine; and a few other exciting medicines.

The sedative method includes the local abstraction of blood; the use of the warm bath; the Buxton waters;

soothing fomentations; local moderate warmth, as by flannel; opiate liniments and plasters; bodily rest; and, internally, narcotic medicines; gentle sudorifics; agreeable employment of the mind.

Blisters and issues constitute a mixed treatment; as, on the one hand, raising irritation on the surface; and, on the other, diminishing the circulation of the part by the discharge which is produced. Heat, also, in whatever way applied, so as to produce perspiration, becomes a remedy of two-fold character; exciting, by its peculiar stimulus to the vessels; and relaxing and reducing action, by the active process of evaporation which belongs to this forced action on the skin. When stimulant treatment has been required, I have often found advantage from electricity; and the following case is an interesting example of the success of this powerful agent. The patient was a young man of twenty-two years of age, and of a strong and healthy appearance. I shall offer the narrative of the case in his own words:

“In the spring of 1816, I experienced an attack of rheumatism in the left arm, which had been affected the preceding autumn. It was then cured by the application of brown paper, which in one night removed it. I made the same experiment on this occasion, but without success. My next remedy was a very stimulating embrocation, which I applied by means of strong friction night and morning. At first I derived from it some slight benefit; but, in a short time, the pain returned with increased severity. A large blister was then laid upon the parts, but without success. The pain had now become excessive, and prevented my sleep. The affected arm became flabby and smaller than the other, yet it was scarcely sore to the touch; differing in this respect from a former attack, when I could hardly bear the smallest

pressure. The pain was a constant aching. At last it began to extend to the knees, and came on in spasms; so violent, that, when in bed, I was obliged to put on my dressing gown and lie on the outside. After a little exposure, it would then cease. At this time I took several internal medicines, and confined myself to the house; but without any apparent good effect. The attack lasted about three months; during which period I had but little sleep. I usually went to bed at eleven or twelve, and awoke in great pain about two, and could not afterwards procure any sleep. I passed the time in reading upon the bed, as it was impossible to endure the violent pain which the warmth of the clothes occasioned.

“As the disease appeared to be confined to the nerves, electricity was next advised. This operation produced great pain and violent perspiration. After the first application, the pain on that day increased; on the following one it was moderated; and daily diminished, until I was perfectly restored. It must be observed, that on all occasions I had been advised to wear flannel, but had incautiously discontinued its use. I resumed, however, this covering to my skin soon after the attack which has been described. Two years have elapsed since my cure by electricity, and I have been entirely free from rheumatic sensation.”

The alterative plan of treatment embraces that course of remedies, both in medicine, and in diet and the general regimen, which has already been laid down in this Treatise; but to the medicines before described, I may add the use of the liquor arsenicalis, as being sometimes a powerful remedy in certain states of chronic rheumatism.

As a rubefacient application, I have found most advantage from a liniment according to the formula, p. 271;

increasing the proportion of *tinctura lyttæ*, and sometimes also adding a portion of *liquor ammoniæ*.

In a few instances, I have seen advantage derived from the volatile tincture of *guaiacum*; but I have for the most part found this a heating and prejudicial medicine. The effects of bark have appeared to me more favourable.

In the third volume of the *Medico-Chirurgical Transactions*, Dr. Marcet has communicated a very interesting account of the cure of *sciatica* by means of excessive exercise in walking, when the patient was warmly clothed in flannel.

The ingenious author of the case, who was himself the patient, relates, as follows, the effects of the proceeding which he adopted:

“ With the utmost difficulty I proceeded half a mile; and the pain I suffered contributed not a little to the effect of the exercise in promoting perspiration. I returned home in a profuse sweat, rubbed myself dry before a fire, and went to bed. In about an hour I got up, found myself very much fatigued, but in other respects not worse. Forty-eight hours after this I repeated the same kind of exercise, and found that I could walk a mile with as much ease, as I had walked half that distance on the first day. My general sensations were the same as before; but, as the fatigue diminished, I thought I could perceive an amendment in my rheumatic pains. Two days afterwards I took my third walk, proceeding as before; and after it I had a better night, less interrupted by pain than any I had enjoyed for eighteen months. From that moment I looked forward with confidence to a cure, and I have not been disappointed. Every succeeding walk has diminished my sufferings;

and I may safely say, that, after the sixth, I was as free from pain as I had ever been in my life."

This patient had previously made trial of a course of calomel, taking from two to three grains of it, with a certain quantity of opium, every twenty-four hours, during six weeks; seconding its efforts with frequent blistering. He adds, "I have applied as many as three blisters at a time, from my hip to my foot, and renewed them as soon as the skin was sufficiently healed to allow a fresh application. By other medical advice since that period, I have tried bathing in warm sea water, in artificial sulphureous waters, the same as those which had afforded me relief in Paris, and in the Bath waters. I have also used the dry pump there, vapour baths, and heated air, friction, cupping, leeches, and electricity. I have taken internally nitric acid, James's powders, guaiacum, nitre, cicuta, hyoscyamus, and the eau medicinale; bark; and, finally, arsenic; but all to no purpose. Were I to state in general terms the effect of all these, I should say that cicuta, hyoscyamus, and the eau medicinale, procured me temporary relief; and that bathing of whatever kind usually made me worse. In short, I had nearly exhausted the materia medica, and I had no hope left but in a more favourable climate."

Of his particular method he gives the following description: "I usually proceed to my sweating walks in the following manner. Next to my skin I wear stockings, drawers, and a shirt, all of fleecy hosiery. Over these I put one, two, or three pair of flannel drawers; one, two, or three flannel waistcoats; and round my hips and loins I gird six yards of thick flannel; making, besides the drawers and waistcoats, eight thicknesses of flannel on the chief seat of pain, and the origin of the

sciatic nerve. Over all this I wear warm pantaloons and a great coat. When I have walked one or two miles, more or less, according to the heat of the day, I am generally in a profuse perspiration. I return home, take off my wet clothes, have a couple of changes of well aired flannel, and then lie down upon a bed not warmed. I use no means to excite further perspiration after the muscular action is over; but, on the contrary, rather incline to check it as speedily as I can, taking particular care however to avoid catching cold.

“I do not perceive that the quantity I perspire, has any influence on the efficacy of this remedy. I imagine that a violent action produced in the general system is the chief cause of its salutary effect. In consequence of this opinion, I cease the exercise the moment that a very increased action is well established. This is fully produced with the above quantity of clothing in moderately warm weather, by walking from one to two miles. For patients who are very much disabled, the quantity of clothing might be increased, and the distance diminished. When the excitement is well established, I find my pulse rise to between 90 and 100, and it is full and strong.” Notwithstanding the apparent simplicity of this method of practice, and its remarkable success in this case, I must observe, that it is too active a plan of treatment to be indiscriminately employed; and, indeed, should never be resorted to without medical sanction.

When a rheumatism of the nerves prevails in a general manner over the body, the Buxton waters sometimes produce the best effects.

In this state of constitution, the use of horse exercise, freely and regularly taken, proves of the utmost service.

If this form of rheumatism; or, I may state it as a gene-

ral position, if any form of rheumatism be depending upon a morbid state of the digestive organs, every kind of treatment must be held secondary and subordinate to that which may be adapted to the removal of such fundamental cause of irritation. Among soothing medicines, I can offer very favourable testimony of the combination of stramonium and lactucarium. In some instances of recent attack, in which wandering nervous pains have been mixed with lumbago, its effects have been surprisingly successful.

In a few cases, even of long standing, I have experienced much satisfaction from this remedy.—A lady who had for two years suffered severe and almost constant pain in the nerves of one arm, producing the mixed sensations of “gnawing and shooting, heat and cold alternately, sometimes peculiar feelings of pricking as if from hot needles, and a weakened and almost disabled use of the limb,” received a cure from the last-mentioned medicines, and the free use of the stimulant liniment with tinctura lyttæ, &c. In some cases, however, both of wandering and fixed rheumatism, this treatment has entirely disappointed my expectations; and a more active plan of remedy has been required.

I was myself once the subject of chronic rheumatism, which chiefly affected the nerves of the lower limbs; producing, with the usual peculiar sensations, occasional spasms and cramps of distressing severity. I had made trial of a long course of remedies, especially of the bark in substance, with free doses of hemlock; and had used warm sea bathing for six weeks, without the smallest advantage. At last, I obtained a cure from taking a moderate dose, twice or three times a day, of the pulvis ipecac. compos. with the addition of a sufficient portion of potassæ sulph. to affect the bowels; uniting with this

medicine a generous plan of diet, and a determination, whatever consequent pain might be produced, to persevere in taking abundant exercise both on foot and on horseback.

In some cases of long standing sciatica, I have known very great benefit derived from the use of dry pumping, as it is practised at Bath.

In the worst chronic states of local rheumatism of the nerves, I have, in a few instances, seen the cure obtained, or very material relief afforded, by the successive application of blisters.

When chronic rheumatism affects the superficial ligaments, the free use of the evaporating lotion (p. 254) is very efficacious in removing the passive inflammation which prevails; and in restoring the ease and comfortable motion of the parts.

When the affection is deep seated, the treatment is often difficult and tedious. Accordingly, as the complaint has been of long or short duration, we are chiefly guided in the choice of the stimulant, the sedative, or the united method of treatment.

When the bursæ and tendons are the seat of complaint, and the disease is recent, constitutional treatment is our chief dependance; but, in this case, we shall derive much auxiliary advantage from the evaporating lotion. Leeches have been much recommended by authors; but my experience has led me to prefer this method of stimulant evaporation. If the affection of these textures, namely, the bursæ, the tendons, and ligaments, have been of long standing; and in which case, the muscles themselves have a strong secondary participation in the disease, becoming small, flaccid, and powerless, the rubbing process mentioned at p. 370, and the use of rollers, constitute the only efficient practice with which I am acquainted.

Any inflammatory state of the synovial membranes, which puts on the character of rheumatism, must be treated by the most active means which are found to remove the inflammation of deep-seated parts of important organisation. Most cases, however, of this particular description fall rather within the province of the surgeon, than of the physician.

Rheumatism, which has been produced by imprudent exposure to cold during the use of mercury, becomes, when in its worst form, a severe and tedious complaint. I have, in most cases of this kind, seen that all the ordinary methods of treatment are of no avail; or, afford only palliative and very temporary relief: but I have repeatedly had the satisfaction of witnessing the cure to be effected by resuming a well-conducted mercurial course. Dr. Bardsley, in his valuable and very instructive Reports, adds his testimony to this important practical fact. This author, in summing up his general conclusions respecting rheumatism, observes, "The cure of chronic rheumatism, even in its most severe and protracted form, is not to be despaired of, provided vigorous and active remedies be duly administered."

Except in the above-mentioned example of rheumatism; and also, in cases in which I have had reason to believe in a syphilitic cause of the pains, I have not been induced to prescribe mercurial treatment to the extent of salivation.

Lastly, I shall offer a few remarks on the treatment of lumbago; and of rheumatism affecting the aponeurotic covering of the muscles situated on the back, or in other parts. In a recent attack of lumbago, occasional purgatives with calomel, antimony, and colocynth, in conjunction with the neutral salts; the free employment of crude opium with antimony or ipecacuanha; and the use of the

warm bath; usually command the cure: and for the most part the latter remedy is not required. A large blister applied upon the loins, is a remedy at once safe, and of great efficacy. If the disease have been of long standing, it is usually more or less blended with sciatica, and presents a case of difficult management. I have seldom seen that the common warm bath, as an early remedy, has afforded any relief. Dr. Bardsley speaks in high terms of the local vapour of hot water, which he directs to be applied by means of pipes connected with a boiler; "so as only to permit the vapour to strike upon the affected part at some distance from its aperture." He also recommends a stimulating liniment to be rubbed in upon the part during the application of the vapour. He says, in speaking of the treatment, "By this its stimulating effects were heightened, pain was abated, and the cure much accelerated." This author thinks favourably of the powers of electricity in this form of the complaint; and I have myself witnessed its very excellent effects. Considerable dependance may be placed upon this remedy when under the direction of a skilful operator. I am compelled, however, to add, that on two occasions, where much benefit might have been expected, I was disappointed in the event. I am much satisfied with the auxiliary effects of the liniment described at p. 271, rendered more or less stimulating as may be required, when it is not convenient to resort to electricity; or, it may with propriety be used even in conjunction with that remedy. According to the state of the constitution, purgatives, alteratives, sedatives, and tonics, must be separately or conjunctively administered; and horse exercise, when exposure is not improper, and, when *practicable*, must not be neglected. I can, from my own personal experience, speak with confidence of its very useful effects. I must add to this

recommendation, that the affected parts must be well protected by warm clothing. A flannel bandage firmly rolled around the body; and, when sciatica is united with lumbago, carried on down the limb; often lends a very material aid to the cure.

When the aponeurosis of the muscles of the back, or in any other situation, is suddenly affected, the complaint is connected with some recent exposure to cold; and the means of treatment which restore the free action of the skin; which remove any inflammatory state of diathesis; and which soothe the nervous system (due attention being also paid to the state of the bowels); will certainly be found effectual.

The most compendious manner in which I can exemplify my opinions on the treatment of some forms of chronic rheumatism, and of painful affection of nerves connected with rheumatism, will be in the selection of a few cases, with occasional remarks.

CASE I.

A gentleman, aged 50, of the nervous temperament and of a bilious diathesis, after some hours' exposure to wet, was attacked with rheumatic inflammation in each foot in succession. The acute symptoms disappeared in ten days, but distressing weakness remained for two months. Before much time had elapsed he became affected again in the feet, and also in the knees, hands, elbows, and shoulders; in a hand and shoulder; hand and knee; hand and foot; but never in two parallel parts at the same time. The pain came on late in the evening, and continued through the greater part of the night. It was a severe gnawing pain, and seemed to be in the bone itself. It was often preceded by a sense of coldness in

the flesh, and a general chilliness. This was occasionally followed by heat, and by a swelling of the parts, which gave relief. When I saw this patient, he had laboured under his complaint for ten months, with only short intervals of amendment. It was so fugitive as to situation, that it never remained in the same part more than two or three days. He mentioned that any accidental exposure to a damp atmosphere, or to draughts of cold air, would certainly produce an active aggravation of his symptoms. The complaint also seemed to have an intimate connexion with the state of the stomach and bowels. The secretions were quite unhealthy. The urine usually deposited pink sediment. There was great deficiency of bile, as indicated by the clay-like appearance of the alvine discharges; and the action of the bowels was torpid. He had made trial of the warm bath, and with some relief to the pain; but such excessive perspiration followed, that he found himself debilitated and rendered very sensibly more tender by its use; and was therefore much more susceptible to the changes of the weather.

I prescribed the following medicines :

R Pilul. hydrarg. gr. iiss.

Pulv. alöes compos. pulv. rhei. āā gr. iv.

Sapon. dur. gr. ij.

Decoct. alöes compos. q. s.

M. intime et fiant pilulæ iii. h. s. s. et omni nocte repetendæ.

R Ammon. carbon. gr. xiv.

Succi limon. ʒss.

Mist. camph. ʒj.

Syrupi tolutan. ʒss.

Guttæ nigræ g^{tt}. viij.—M.

Fiat haustus bis terve die sumendus, prout dolores urgeant.

He was directed to apply the evaporating lotion (p. 254) to any part affected with inflammatory tenderness. His regimen, in regard to diet and clothing, was carefully regulated. By a perseverance in this method of treatment for six weeks, lessening the frequency of the doses both of the pills and draught at the latter part of the period, he obtained a cure; and has since remained perfectly free from complaint, now an interval of a year. He has used my prophylactic method of ablution of the head by means of a coarse towel dipped in cold water, and sponging the feet with salt tepid water, very regularly; and with equal comfort and advantage.

CASE II.

A gentleman of the medical profession, aged 44, subject to much daily fatigue from walking exercise, was distressed for a series of years with chronic rheumatism, and severe dyspepsia. Occasionally he suffered lumbago and a painful spasmodic state of the intercostal muscles. At one time the upper extremities were most affected; at another, the lower. At the wrist and ankle there was painful bursal distension; on the fibula of each leg, some thickening of the periosteum. The flexor tendons of the feet were usually painful. He was troubled with biles in different parts, and with psoriasis in the arms. In the thigh an ill-conditioned ulcer formed. The tongue was singularly affected with a sort of fungous ulceration, and presented a cracked appearance on its general surface. This gentleman had tried the bathing in the Bath waters, together with their internal use, without advantage; but he had derived material benefit from the warm bath at Brighton. He relapsed again and again in pursuing the duties of his profession,

It was evident that the system required more of habitual repose; and the errors of the constitution were to be treated by corrective medicine; its debility by supporting regimen. He desisted from walking exercise, and performed his usual avocations in a carriage. The black drop which had proved so useful in the preceding case, did not well agree in this, producing head-ache. The local symptoms were usefully treated by stimulating embrocations. As internal medicines, the most decided benefit was obtained from the *pilula hydrargyri*, and *decoctum sarsaparillæ*. The refraining from fatiguing exercise, appeared to be one of the most important points in the whole management of this case. He gradually recovered.

CASE III.

The following is simply an example of common rheumatic attack produced by exposure to cold; and not related either for its novelty or particular importance. A gentleman, aged 45, of the nervous temperament, perspired profusely in walking upon a day when the sun was hot, and the wind was in the east. On the night of the same afternoon his sleep was restless. On the following day he had slight rigors, and some nausea, and was affected with severe pains in the shoulders, which in the course of the next twenty-four hours extended to the knees and ankles, the calves of the legs, and the sciatic nerve on one side. He described his painful sensations as being deep seated, and as resembling a gnawing and tearing of the parts. The integuments were wholly free from redness or swelling. The abdomen was almost universally tender. The tongue was furred, the bowels were confined, the urine was deficient and

of a deep colour, the pulse was rather increased beyond the natural standard. Hydr. submur et pulv. jacob. in conjunction with a purgative of senna and magnes. sulph. were the medicines first administered; and when their operation was established, a saline draught, containing pulv. ipecac. compos. gr. v. was given every six hours. Rest and abstinence were also enjoined. In a few days the removal of all painful symptoms was accomplished.

CASE IV.

A lady, aged 34, slight, and of the nervous temperament, first suffered from symptoms of lumbago four years ago. By degrees the complaint extended to the left hip. She described the pain as passing to the groin, and all down the fore part of the thigh to the knee. The flesh was sometimes tender to pressure. Fourteen months ago, after having enjoyed a comfortable state of amendment for several weeks, she was in an unexpected moment seized with pain in the hip, so that she could suppose some burning liquid had been poured suddenly into the joint. She could not walk far without suffering severe pain, and if she missed her regular step, she would experience torture. She complained of pains of the arms, attended with a sense of extreme weariness in them. This symptom was particularly troublesome soon after retiring to bed. Blisters and leeches had been applied to the hip without producing any sensible benefit; and the use of the warm sea and common bath had been unsuccessful. The nervous system was evidently relaxed. The tongue was pale; the pulse languid; the head often painful, and the sight affected with spectra and a sense of mist.

I directed the emplastrum opii spread on leather to

cover the whole of the affected hip; the lower part of the limb, which was often affected with much coldness, to be rubbed with the *tinctura lyttæ* liniment, with the addition of *liquor ammoniæ*; warm clothing to the limb; the avoidance of walking exercise; and, internally, a draught three times a day, composed of ammon. carbon. gr. xiii. neutralised with lemon juice, decoct. cinchon. tinct. cinchon. comp. and gutt. nigr. g^{tt}. vj. Her diet was made more generous than before. From these means of treatment, this lady immediately obtained great amendment; and after a few weeks perseverance, she derived a cure, which up to the latest period of my information, a few months since, had been permanent.

CASE V.

A coachman, aged 37, from repeated exposure to wet and cold, first suffered from rheumatism eight months ago. He was distressed with flying pains almost universally, and the loins were sometimes so much affected that he could scarcely turn in bed. The lower limbs were most the seat of complaint. He described his symptoms to be severe pain and weakness in the hips and knees; and at the ankles there was remarkable bursal distension, tenderness to pressure, and occasional redness of surface. His general health was much disordered. The tongue was furred, the appetite deficient, the bowels torpid, the urine deposited lateritious sediment. Mercurial purgatives were first administered, and repeated occasionally. The alvine discharges were dark to the degree of blackness. As opium internally disagreed with this patient, I made choice of stramonium and lactucarium. He took these medicines in graduated doses, at first three times, and afterwards twice a day,

with the best effects. The passive inflammation of the ankles was removed by the use of the evaporating lotion; and afterwards much benefit was derived from the application of a roller*, and from sponging the whole of the feet

* The seasonable use of bandages in chronic rheumatism is an important point of practice; but it is evident that Dr. Balfour has greatly over-rated the efficacy of this remedy, and indeed advised its employment in too general a manner. I received the following interesting communication from a medical friend, filling the office of physician's clerk at a celebrated infirmary. I am requested to avoid the mention of names, out of delicacy.

“ Three patients were admitted, in one of the winter months, complaining of severe pain in the joints of the lower extremities, increased by pressure and by the heat of the bed. The pain was much aggravated when attempts were made to walk. To give merely a general description, I have to state that the pulse was rather full, with general symptoms of slight pyrexia. The bowels were freely acted upon; low diet was observed; and in short the patients were kept at rest and free from excitement. The febrile symptoms were thus removed, but neither was the pain of the joints removed, nor their power restored. Flannel bandages were then applied exactly upon the plan of Dr. Balfour. Immediately on the application, the patients were desired to attempt to walk, which they were unable to do, but said they felt as if supported and strengthened in the limbs, having power to stand with less pain than before. In these cases the use of the bandages was persevered in for a considerable period; as also in some other cases which were admitted about the same time. This had not the effect of removing the disease; and other treatment was then adopted. The conclusion drawn was, ‘that bandaging may aid other remedies, but alone is by no means to be depended upon.’

“ On a subsequent occasion another fair trial was given to this mode of treatment; and the reports upon its issue were decid-

and ankles every morning with tepid salt water. He recovered in the most favourable manner.

CASE VI.

A young man, of delicate constitution, had suffered occasional attacks of chronic rheumatism; the first of which was consequent to the acute form of the disease, for several years past. When he consulted me, he complained of severe lumbago, which was constant; and of flying pain in his limbs. After walking a few minutes, the pain of the back became insupportable. I directed, as internal treatment, the pills of stramonium and lactucarium; and to the seat of pain emplastrum opii spread on a large piece of leather, in conjunction with a flannel roller firmly carried around the loins. These remedies were used with such success, that he quickly recovered. Many months after, from fresh exposure, he relapsed, and was altogether in a state of debility. Now the same pills disagreed, and the stramonium produced those inconvenient effects which are not unusual in its operation, of a dryness of the throat, and a mist of sight. On the present occasion one arm also was severely rheumatic. He now derived material benefit from free doses of cinchona, the alterative aperient use of pilul. hydrarg. with extr. colocynth comp. and active fric-

edly unfavourable as to its possessing any important efficacy. Some cases in the ward not bandaged, as nearly as possible under the same circumstances as others in which bandages had been used, were the soonest relieved. In one case indeed a bandage was applied to one of two limbs equally under the influence of the disease; and the limb not bandaged was reported to be well sooner than the other."

tion night and morning with the compound tinctura lyttæ liniment as lately described. I have reason to believe that all the symptoms were gradually overcome.

CASE VII.

A lady, remarkably of the nervous temperament, suffered from chronic lumbago; and pains of the wrists and ankle joints, attended with a remarkable weakness of the ligaments; which was indeed rendered evident upon examination, from the preternatural flexibility of the joints. Walking soon produced extreme fatigue and aching; and the amusement of drawing was too great an exertion for the wrists. The use of the shower bath, graduated in the temperature and quantity of the water; a steel tonic; a supporting diet; and the air of the country, proved successful in this case.

CASE VIII.

A female servant, pursuing her occupations out of doors, upon a hot day when the wind was in the east, was seized with rheumatism in the right shoulder; which had continued eight months when she consulted me. The pain was most severe at night, when she was warm in bed. She described the usual sensation of rheumatism, a gnawing as if in the bone; sometimes it was attended with remarkable coldness, at others with burning heat. The muscular power of the whole arm was become much weakened. She had used irritating embrocations without relief. With one application she had even vesicated the skin. The volatile tincture of guaiacum had been taken

in free doses without advantage. I directed the shoulder to be covered with emplastrum opii spread on thick leather; and internally, pills composed of pulv. opii crudi with pulv. alöes. compos. to be taken twice or thrice a day, according to the degree of pain. This treatment gradually succeeded; and the cure has been permanent; a year having now elapsed.

CASE IX.

Many cases have come under my observation in which the patient has considered himself to be labouring under rheumatism, the pains having been in reality dependant upon latent syphilitic virus. I know no form of disorder more insidious, and more demanding careful diagnosis. I shall offer two examples.

A gentleman related, that, during four months previously to my seeing him, he had suffered severely and constantly from pains of the shoulders and arms, which he considered to be rheumatic; and that he had contracted the complaint from exposure, after quitting hot rooms at night, and going into the open air. These pains were not influenced by the state of the atmosphere; and the digestive functions were healthy. Most of the usual remedies for rheumatism had been tried to no purpose. My discovery of the nature of the case was directed by observing, partly on the arms, but more especially on the back and breast, a scaly eruption, which was accurately the psoriasis guttata syphilitica, as described by Willan and Bateman. The history which the patient gave of such original circumstances as might assist this explanation of the case, was too much disguised to deserve dependance. When on such occasions any concealment is practised

(upon very false principles of delicacy), we must not allow our judgment to be deceived; but, trusting to the evidence of our senses, and our own legitimate reasoning, we are to act with decision. I prescribed hydrarg. oxy-mur. formed into pills with bread, adding some muriate of ammonia; and with this alterative, joined the decoction of sarsaparilla, and the regular use of the warm bath. At the end of three or four days the pains entirely quitted, and never returned. The case required a steady treatment of three months; but the very quick relief which was afforded to the patient's sufferings, and the permanent cure which was finally obtained, gave a most satisfactory proof of the correct diagnosis of the case.

CASE X.

A gentleman had suffered various secondary symptoms, and undergone a long and apparently sufficient regular mercurial course. He had been convalescent two months, and was only recently recovered from the sensible influence of the mercury. After exposure to cold air late at night, he was attacked with pain on one side of the forehead, and in the eye. This intermitted, and in its returns extended to the upper part of the head; and so much increased in severity as to produce delirium. The eye itself became inflamed. Leeches and blisters were applied freely and repeatedly, and various medicines were given, amongst which, bark and opium, with James's powder, were the principal; but from the anodyne alone was any degree of relief obtained, and that only of a temporary nature. The intensity of the pain produced an exquisite tenderness of the parts. The lacrymal sac and duct became inflamed and painful; and the skin of the nose on the same

side now assumed a dark copper colour. This immediately induced a new and fortunate view of the disease. The medicines mentioned at the conclusion of the last case were administered with the happiest effects. After a few doses, the pain became slight, and from the expiration of a week never more returned.

In the following instance of *mercurial* rheumatism, a different treatment was required.

CASE XI.

A gentleman had undergone a long and severe mercurial course, and imprudently exposed himself to wet and cold before his recovery. The consequence shewed itself in a severe and universal attack of muscular rheumatism, which rendered walking exercise always painfully difficult, and sometimes impracticable. He was deprived of sleep at night; and such constant constitutional irritation prevailed, together with pain, that he became emaciated, exhausted, and indeed alarmingly weak. It was so well known that too much rather than too little mercury had been used in this case, that a return to that medicine did not seem reasonable. On the contrary, the vapour bath was chosen as the remedy to expel the mercury by the surface, and sarsaparilla with bark and a supporting regimen were given internally. Other treatment had been previously tried without the least advantage. The remedies now mentioned succeeded perfectly. The very happiest effects resulted from the use of the vapour bath, which was regularly continued for a considerable time.

CASE XII.

The following case exhibits very strongly the rheumatic diathesis, and the obstinacy which the chronic form of the disease, in particular constitutions, assumes under every plan of treatment which can be devised. A gentleman, aged thirty seven, of the mixed temperament, of full habit, and rather corpulent, was first affected, at the age of twenty, with a pain in the knee-joint, which continued more or less troublesome for a year. Leeches proved more useful than any other of the various remedies which were employed; but, according to the patient's belief, the disease wore itself out *with time*. Five years ago, on a cold day, when feeling in good health, he was suddenly seized with lumbago. This attack appeared to have yielded to the prompt treatment which was used; but soon after, on fresh exposure to a north-east wind in an open carriage, the lumbago returned severely, and also the pain of that knee affected on a former occasion. He remained ill four months. A course of bark, the extract and decoction in large doses, and the application of flannel bandages, appeared to be of real service, after other means of treatment had failed. He remained quite well about three years, and was capable of strong exercise both in walking and riding.

In January 1818, when exposed to the east-wind on the outside of a carriage, he felt himself suddenly seized by the cold on one side of his body; and, in two days after, the lumbago and pain of the knee came on severely. Much alarmed, and with the idea of arresting the progress of the disorder, he went immediately to Bath; and there used the hot bath, and pumping on the affected parts for

one month; also taking aperient and alterative medicines. He returned to London not benefited. Blood was taken freely from the loins by cupping, and from the knee by leeches. Electricity was next used, and various embrocations. No advantage was obtained. He went to France in June, and perspired profusely every day with exercise, the weather being intensely hot. The pain now quitted the knee and part of the loins, but took the track of the sciatic nerve, from the hip to the calf of the leg. Walking exercise invariably and almost immediately brought on pain, which he described as severely aching, and accompanied with spasmodic shootings;—very much the sensation of cramp; and this was experienced most in the calf of the leg. It was in vain that on two occasions he exercised the fortitude of walking a great distance, on the principle of treatment stated at p. 683. Rest always relieved, exercise always aggravated, his sufferings. Cupping on the hip, blisters, the vapour bath, fomentation, electricity, emplastrum opii, and flannel bandages, were amongst the external remedies employed without success. The two last afforded most comfort. The former seemed of no avail. Much friction with the hand was too exciting. Complete rest in bed for two or three weeks had a better effect. Mercurial friction to the extent of producing ptyalism, aperients, alteratives, bark, liquor arsenicalis, guaiacum, the black drop three times a day, comprised the principal constitutional remedies, which were used in proper succession. A course of Cheltenham water was taken with advantage to the general health, but without the least influence on the local symptoms. As a last resource, he went to Weymouth, to try the effect of a removal to a residence more favourable than that which he had occupied, which certainly was damp and exposed to cold winds; to try also the effect of complete repose, and, above all, a steady course of warm

sea bathing. At the end of three months, he obtained a cure. It deserves to be mentioned, that during a whole month he did not derive any sensible advantage from the bath, but latterly he became quite sensible of its happy influence. The patient has continued perfectly well. In reflecting upon all the circumstances of this interesting case, I am persuaded that the fulness of habit, and a surcharge of the vessels of the liver, were favourably removed by the use of the mercurial treatment and of Cheltenham water, so as to lay a foundation for the benefit afforded by warm sea bathing. The black drop procured temporary, but no permanent benefit. The local disease was evidently involved with a wrong state of the constitution, and time undoubtedly had a great share in adjusting the healthy balance.

CASE XIII.

A lady, slight in form and of delicate constitution, in the summer of 1817 found herself gradually becoming relaxed in her general state of health. In the month of June, after some excessively hot days, she first experienced an uneasy sensation in the legs. The following descriptive account is in the words of the patient. "The legs were burning hot and weak, and the right in particular felt asleep, and as if swollen, but it was not in any degree. I took no notice of this for about ten days, at which time I was obliged to keep always in a reclining posture, in consequence of a violent spasm in my right leg. This spasm took its rise under my knee, darting down to my heel; or, nearly, but not quite, up to my hip. It was followed by a slight convulsion of the muscles of the limb, and this was visible. I compared it to a narrow animal, about five

inches long, struggling for a passage down or up. This violent pain lasted about two months. I was never tolerably easy but when lying down entirely flat. The paroxysm of pain went off gradually, leaving my legs again sleepy and uneasy; and I felt great stiffness and weakness about my back and loins. The right leg was colder than the other, and the skin was much more pale in appearance. I may add to this account, that when the severity of the disorder abated, the sitting posture for a considerable time produced great uneasiness in the right thigh and leg, and caused in both limbs the sensation of pins and needles."

This lady could not take opium in any form without material disagreement. Stramonium produced a sense of dryness in the throat, so intolerable that it could not be continued for more than two or three days; nor did it relieve the pain. Extractum conii, however, had a manifest influence over the pain; and, although it caused giddiness, and therefore caution in its use was necessary, it proved very serviceable. So easily was the system excited during the early part of the complaint, that tonics immediately produced feverish heat. In regard to local treatment, blistering was used without advantage. A liniment, as at p. 271, with a larger proportion of tinctura lyttæ and the addition of liquor ammoniæ, was decidedly useful, producing an agreeable warmth in the limb, and causing the feeling of more energy and strength. Friction with the hand was also very useful. Warm sea bathing completed the cure; and, from its commencement, gave comfort, and very sensibly produced the best effects. Latterly, the tinctura ferri ammoniat. taken in water simply, was of much advantage as a tonic, and in restoring some functions of the constitution which had been suspended. The patient has continued perfectly free from relapse, and in all respects remains well.

CASE XIV.

A Gentleman, aged forty-two, of slight form, and appearing delicate in constitution, but yet enjoying general good health, in the year 1818, in the month of September, was ardently engaged in shooting, and walked in wet fields, regardless of any exposure. At the end of the month he became first affected with lumbago. The pain spread round to the left hip, and partly down the thigh. He was confined to his bed or chamber about a month. At one time the pain quitted the loins and hip, and went to the knee and ham. This part was fomented, and the pain again returned to the hip. He stated that cupping on the hip, and leeches fifteen in number applied to the same part, had failed to give relief. A blister had been useful. He had taken James's powder, so as to produce copious perspiration; but without any sensible advantage. He gradually recovered from this attack, and in December again returned to the sports of the field, walking from eight to twelve miles daily. A severe relapse took place; and now the sciatic nerve of the limb before affected became distinctly the part in pain. In the first instance the limb appeared to be always cramped. It was much colder than the other. He had a frequent sensation of creeping down the limb; sometimes as if an insect were crawling. The blood itself seemed to be trickling. The sensations were very various at different times. In the early part of this relapse he was under the necessity of travelling, and in this way aggravated and fixed the complaint. At length the pain became truly severe, and was a constant aching along the whole course of the sciatic nerve. The following may be added to the state-

ment of the occasional sensations before mentioned. The feeling of a remarkable bigness of the limb; coldness in one part, and burning heat in another; a heat on the surface of the foot, as if recently scalded; the intensity of the pain shifting its situation to different parts of the thigh and leg; cramps; starting of the limb, particularly on the first attempts to sleep at night. The whole limb became rather diminished in size; the muscles were much relaxed. The tenderness to pressure along the whole course of the nerve was excessive. The pulse was regular and seldom quickened. The general health appeared to be scarcely affected, except as languor and some exhaustion followed the attacks of pain, and became consequent to the necessary confinement.

The warm bath, embrocations, the emplastrum opii, in conjunction with internal medicines, failing to give relief, a caustic issue was made at the hip; and this seemed to be a source of benefit. A course of mercurial friction upon the limb was pursued for a considerable time, and the constitution was kept moderately under its influence, as shewn by the appearance of the gum: but, with regard to the state of the nerve, the mercurial action did not produce any apparent effect. *Liquor arsenicalis* was administered, with a small addition of opium; but it disagreed with the stomach and bowels. The most decided advantage invariably attended the administration of opium as an anodyne. The black drop was taken in doses of eighteen drops three times a day, joined with a saline aperient; and the patient was furnished with pills, containing in each gr. i. pulv. opii crud. gr. ss. pulv. jacob.—of which one or two were to be taken on any emergency of pain. By this treatment, a complete dominion was obtained over the severe sufferings of the disease. The returning pain was met by the regular

dose of the opiate. The recovery was slowly progressive, and, at the time that I am now writing, is not completed; but the cure promises to be certain. The dose of the anodyne was gradually lessened, and latterly was joined with a draught, as described at p. 266, cascarilla being substituted for the gentian. A course of warm sea bathing is proving highly useful. Friction for an hour daily, and the use of an embrocation, as mentioned in the last case, afford evident advantage, in assisting to restore the power of the muscles, and the healthy circulation of the limb.

CASE XV.

I am favoured with the particulars of the following case by Mr. King, of George Street.

A gentleman received a violent fall from his horse in hunting, by which the glutæi muscles were much bruised. After the abatement of swelling and inflammation, a severe and constant pain remained in the hip, which, in a short time, extended down the thigh and leg in the course of the sciatic nerve. Much medical treatment in the country was tried without success. He came to town, and placed himself under the care of Mr. Astley Cooper and Mr. King. At first the liquor arsenicalis was administered, and external stimulants were used, without the least success. The patient was then directed to confine himself entirely to bed; and the black drop was given in divided doses, three and four times a day, in a saline draught with bark. By this treatment the cure was effected in the space of three weeks.

Mr. Cooper informs me, that, in a case of sciatica which had been produced by exposure to cold, he succeeded perfectly with the treatment by anodynes and

rest in bed, after many active remedies, and some of them of a painful nature, had wholly failed.

Did the present opportunity permit, I could add largely to these examples, shewing the superiority of soothing methods of treatment in *neuralgia*, over the use of irritants and stimulants. I say of neuralgia; for when the case resolves itself into a distinct affection of nerve, it is, in most instances, no longer to be considered in the character of rheumatism. All that favours composure of the nervous system must be studied:—the medicinal plan as already described; rest in bed; the exclusion of all affairs which give anxiety to the mind. It is important also, and will prove a great æconomy of time in the event, that, in the commencement of these attacks, the patient should quietly surrender himself to the proper treatment, instead of making vain attempts of delay; by which, with certainty, the disease will become fixed, very tedious, and much more severely painful.

In cases in which the hip is severely affected, the pain being felt most near the head of the sciatic nerve, the grounds of diagnosis between any diseased action taking place in the joint itself, and the disorder of the nerve merely, are to be carefully considered. When the nerve alone is concerned, the pain and tenderness are marked along its course, both by the sensations of the patient, and by manual examination. The freedom of the action of the joint, and the preserved form of the nates, the diffused sensations of the limb, the character of the pain, the spasmodic state of the muscles, and the whole nature of the disorder, as shewn in the narratives which I have given, will, I apprehend, afford sufficient marks of distinction. When real obscurity in a case does exist, the diagnosis should be submitted to the opinion of the surgeon.

It was my intention to have entered upon the consideration of those diseases of the *bursæ mucosæ* which are distinct from the consequences of gout or rheumatism, and which are so truly constitutional, that they come chiefly under the care of the physician; but the subject is too long and too important to allow of a cursory notice. I must, therefore, postpone it entirely to a future opportunity; indulging the hope that I may be enabled to prosecute these inquiries, which I have now treated with brevity, upon an extended scale; and upon the same principles of analytical investigation which I have here attempted.

On the value of systematic arrangement, I need not expatiate. By establishing our diagnosis on the basis of anatomy and physiology, we lay the foundation of greater certainty in our inquiries; and by observing, carefully and without prejudice, each distinct appearance in diseases which is presented to our view; we gradually acquire those philosophical principles of pathology, which constitute the leading features of distinction between the art of the Physician and the pretensions of the Empiric.

FINIS.

INDEX

- Abdominal viscera, morbid states of, to be judged of cautiously by external examination 90
- Abernethy, Mr. his observation on a disordered state of the alimentary canal 83
- on the connexion which exists between the digestive organs 89
- according to his principles, unirritating doses of mercury recommended 177
- view of his peculiar practice in a morbid state of the digestive organs offered 543
- Abscess from gouty inflammation, a very rare occurrence 146
- from gouty inflammation, its peculiar nature 146
- examples of in gout 286, 325, 328, 329
- chemical examination of the matter from it 291
- Accountant, London, his worst fits induced by excessive application in his business 606
- Acetic preparation of colchicum 186
- — its sensible properties 217
- acid, its modifying influence on vegetable medicines 219
- Acid, uric, contended for by some authors as a proximate cause 115
- phosphoric, the opinion of Berthollet respecting 119, 127
- phosphoric, own conclusions respecting 128
- — table general of its proportions in the urine 120
- muriatic bath as a remedy in gout 247
- nitro-muriatic bath of Dr. Scott 327, 563
- — peculiar manner of its acting on persons under the strong influence of mercury 569, 570
- nitric, when useful in dyspepsia 548
- matter ejected from the stomach, its nature considered 106
- Acidity in the primæ viæ an exciting cause of gout 105
- Acids and alkalies, the administration of, in gravel, considered 582, 589
- Acne rosacea, strong example of, in a gouty patient 272
- Aconitum napellus, or wolf's bane, its medicinal properties 236
- Adams, Dr. his distinction between disposition and predisposition to disease 56
- Adult age as predisposing to gout 59
- Age at which gout makes its attack, tabular view of 54, 61
- or period of life of persons, as predisposing to rheumatism 650
- Air of the country very restorative 269, 605
- heated, as a remedy in gout 247
- Albuminous urine, pathological discussion respecting 298
- — the best test for its detection 300
- Alcohol, its large proportion in wine 619
- Ale, strong, the free use of, disposing to gout 105, 618
- Alimentary canal, disordered state of 36, 82, 288, 387, 389
- Alkaline medicines, as remedies in gout, considered 178, 625
- — to be used with confidence only in conjunction with other remedies 627
- Aloetic pills, useful formula of 282
- Alum, solution of, as a test with urine 281
- Ancients, their authority on gout, why doubtful 10, 640
- Anger, a fit of, compared by Sydenham to a fit of gout 114
- Animal food, how conducive to gout 68
- Anodyne, term defined 227
- Anomalous, term considered 8
- forms of illness in gouty persons, question concerning, discussed 9, 374

- Anxiety, mental, as predisposing to gout 68
- as a cause exciting the paroxysm 113, 606
- Aponcurosis of the muscles, permanent changes in the structure of, produced by gout 44
- when affected by rheumatism 675
- how to be treated 692
- Apoplexy, pathological explanation of 9, 94
- cases of, occurring to gouty persons 473
- those disposed to, should pursue a most careful regimen 475
- in connexion with gout, how to be treated 440, 472
- cases of, occurring in the paroxysm, and the treatment 458
- incidental to some gouty persons pursuing bad habits 141, 465
- may arise from injudicious treatment in gout 9, 351, 474
- Appetite, good, frequently connected with a faulty state of digestion 81
- not a test of healthy digestion 79, 82
- depraved 16
- Aretæus referred to on hot and cold gout 131
- Arsenicalis liquor as a remedy in chronic rheumatism 684
- Arthritis, the term preferred to podagra 4
- the term used by the ancients, generally, to express the diseases of the joints 10, 640
- Ascites occasioned by the eau medicinale 215
- an occasional remote sequel to gout 466
- Ass's milk much recommended by the author, 268 — See also the various cases
- Assimilation, healthy, on what it depends 79
- Asthma, as occasionally supervening on gout, or gout on it 466
- Atmosphere, effects of its variation on gouty persons 94
- Atony, the term, as employed by Cullen, considered 7
- Aurelianus Cælius referred to for ancient terms of gout 10
- Baillie, Dr. his case of a gouty patient with a curious palpitation of the heart, referred to 16
- Baillie, Dr. upon a strong pulsation of the aorta in the epigastric region 91
- Ballonius, the first author who considered rheumatism as a distinct disease from gout 640
- his theory of rheumatism briefly stated 641
- Bandages in chronic rheumatism, their propriety discussed 698
- Bardsley, Dr. his instructive reports on rheumatism referred to 690
- Bark, Peruvian, praised by certain authors as a remedy in gout 219
- its merits as a remedy in acute rheumatism discussed 663
- Barthez, as a French author on gout, referred to 131
- his pathology of gout 131
- referred to in praise of friction as prophylactic 602
- Bath, acid muriatic, as a remedy in gout, considered 247
- nitro-muriatic acid, as a general remedy, considered 563
- peculiar manner in which it acted on two persons under the use of mercury 569, 570
- a spacious tepid one recommended 366
- shower, under what circumstances recommended 367
- warm, its effects in the paroxysm 319
- sometimes excites the gout 368
- remarkably useful in a case of nephritis calculeosa 597
- sea, requires perseverance 368
- vapour 323
- to be used with discretion 369
- waters, different eminent authorities respecting 352, 363
- cases exemplifying their effects upon gouty persons 353
- their temperature stated and considered 364
- require careful preparation of the constitution 356
- Bathing, sea, cold, its propriety for gouty persons discussed 363, 601
- warm, case in which it did not agree with a gouty female 367
- sea, cold, its use as prophylactic against rheumatism 670
- cold, ditto *ibid.*
- Battley, Mr. referred to respecting opium 227
- his opinion respecting stramonium 234

- Belladonna atropa, its medicinal properties 235
 Benzoës tinct. comp. sometimes proves an useful medicine 347
 Berthollet, upon an excess of phosphoric acid in the urine of gouty persons in the paroxysm 118
 — on the acid nature of the transpiration 131
 Berzelius, his contradiction of the opinion that free phosphoric acid exists in the urine 119
 — that the matter of transpiration is always acid 131
 Bile, excess of, occasionally excites a paroxysm of gout 107, 413
 — the various states of, in disease 84
 — the most accurate test for its detection described 556
 — and bilious, terms too indiscriminately applied 549
 — experiments upon 556
 Bitters, corrective tonic, formulæ of 265, 266
 Black drop, account of the 225
 Blackhall, Dr. his opinion on albuminous urine in dropsy referred to 298
 Bladder, irritability of, in gout 18
 — — suspended by the paroxysm of gout 430
 Blane, Sir Gilbert, Bart. his abstract of diseases at St. Thomas's Hospital referred to 72
 — his remarks on the use of the alkalies in gout 628
 Bleeding, general, in gout, considered 176, 178, 321
 — used in a case of chronic gout 379
 — used in cases of chronic retrocedent gout 447, 449, 452
 — in gout, physicians even too cautious respecting 469
 — when proper for gouty persons, 178, 379, 416, 471 — See the cases of retrocedent gout
 — general, as a remedy in acute rheumatism, considered 657
 — as prophylactic in regard to gout, considered 629
 — general and local, pathological observations respecting 471
 — as a general remedy in persons of gouty constitution considered 469
 — to a great extent, used as a remedy in the paroxysm without success 321
 Bleeding, local, in sciatica 689
 Blisters condemned by Cullen as a remedy in gout 243
 — to gouty parts not approved by the author 243, 374
 — in gout advised by Musgrave and Rush 374
 — their use in local rheumatism 683, 689
 Blood, state of, in a case of gout 452
 — when drawn in disease, indications which it presents 453
 Bodily structure, as a remote cause, considered, with relation to rheumatism 650
 Boerhaave, his term of podagra referred to 4
 — did not originally distinguish rheumatism from gout 11
 — his pathology of gout 144
 Bone, example of its being digested in the human stomach 637
 Bones of the feet and hands, according to Dr. Monro, jun. changed into a chalk-like substance 45
 Bootkins and socks, their use in gout much condemned 245
 Bostock, Dr. referred to, as explaining the terms precipitation and coagulation 293
 — as having discovered urate of soda in the pink and lateritious sediment 159
 Bowels, unhealthy action of, 82, 88, 288, and particularly in the cases of chronic gout
 Brain, a morbid state of its functions as affecting the digestive organs, often not sufficiently considered 552
 — — when existing, how to be treated 553
 Brande, Mr. referred to respecting urinary calculi 579
 — referred to respecting magnesia 116, 583
 — referred to respecting the proportions of alcohol in wine 619
 — referred to respecting the use of acids in gravel 589
 Bread, brown, when recommended 614
 Brodie, Mr. his pathological researches respecting the diseases of the joints, referred to 673
 — his case of gout, the dissection 48
 Buffy coat of the blood considered 452

- Bunion, nature of, described 44
Bursæ mucosæ, enlargement of
 in gout 30, 144
 — — nodous 313
 — — in what manner occasion-
 ally affected by rheumatism 675
 Buxton waters, when useful in
 gout 362
 — useful, independently of their
 temperature 366
 — when useful in rheumatism
 682, 687
 — cases exemplifying their effects
 on gouty persons 364
 Cachexy, general, sometimes con-
 sequent to long-continued gout 334
 Cadogan, Dr. his assertion that
 gout is not an hereditary dis-
 ease, considered 54
 — his strong condemnation of the
 Portland powder 630
 Calculi, urinary, statements res-
 pecting 43, 161, 576, 580
 — analysed 161, 165
 — of the kidneys mentioned by
 Morgagni as being often joined
 with gout 43
 Calculus of the bladder, accord-
 ing to Sydenham, frequently
 associated with gout 42
 — appears to be rarely connected
 with gout 43, 467
 — urinary, from gouty persons,
 chemical analysis of 161, 165
 — uric acid, its treatment con-
 sidered 582
 — urinary, very singular one,
 containing the carbonate of lime 621
 — biliary, remarkable effects of
 chlorine upon 561
 Calomel, remarkable manner in
 which it acted in a case of gout 190
 Canal, alimentary, when the par-
 ticular seat of indigestion, how
 to be treated 547
 Carbonate of lime found in an
 urinary calculus 621
 — on the surfaces of the joints 676
 — of potash superior to the sub-
 carbonate 186
 Carditis arthritica, case of, related
 by Dr. Haygarth 469
 Cathartics as a remedy in gout 183
 — as a remedy in acute rheuma-
 tism, considered 660
 Cause, proximate, of gout 115
 — — definition of 130
 Causes exciting rheumatism 652
 Causes predisposing to gout 52
 — exciting gout 101
 Causes predisposing to rheumatism 650
 — various remote, in predispos-
 ing to, and exciting the gout 625
 Chalk-stones, various situations of 46
 — commonly give rise to pricking
 pains in the part just before the
 attack of gout 20
 — an improper term 115
 — not known except in gout 116
 — a rare occurrence among gouty
 persons 145
 — composition explained 146
 — the treatment of, considered 421
 Chalk-stone (so called) curious
 secretion of 48, 53
 Chalybeate water of Tunbridge
 wells, an efficacious remedy in
 dyspepsia 547
 Chalybeate water, its use known by
 the author to have the effect of
 bringing on the gouty pa-
 roxysm 357, 631
 Chalybeates in gout, their propri-
 ety discussed 262, 357
 Chamber horse useful 407
 Champaign, powerful effects of,
 as exciting gout 101
 Characters, external, of the gouty
 inflammation according to si-
 tuation 30
 Chelsea-pensioner's medicine 349
 Cheltenham water in certain kinds
 of dyspepsia to be used with
 caution 546
 — — useful to gouty persons 358
 Chilblain mistaken for gout 23
 Chlorine, its properties as a medi-
 cine discussed 560
 Chronic gout, its characters 14
 — — its history 331
 Chronic rheumatism, history of 671
 — diagnosis of 676
 — treatment of 679
 — as produced by the impru-
 dent use of mercury 678, 703
 Cider hurtful to gouty persons 69
 — to those affected with gravel 619
 Claret agrees remarkably well with
 some gouty persons 617
 Clerk, Dr. Edin. his diagnosis of
 chronic gout from chronic rheu-
 matism fallacious 341
 Climate, variable, predisposing to
 gout 78
 — equality of, very important to
 gouty persons 599

- Colchicum draught, the, most recommended by the author 186
 — autumnale, the preparation of, preferred by the author 188
 — tincture of, its sensible properties 217
 Colchicum autumnale, respecting 193
 — — preparation of, recommended by Sir Everard Home, 195
 — various preparations of, considered 186, 195, 198, 217, 323
 Cold, as an exciting cause of gout, considered 107
 — as an exciting cause of retrocedent gout 434
 — the only exciting cause of rheumatism 652
 — exposure to, in what manner it injures 109
 — causing retrocedent gout 439
 Coldness of the feet 93
 Cold water freely applied to the head, when useful 475, 603, 669
 Colic, severe, in gouty persons 461
 Concretions, gouty, reasoning respecting 144
 — manner in which they affect the extremities 46
 — treatment of, considered 421
 Condiments should be used but very sparingly 613
 Conformation, particular bodily, predisposing to gout 63
 Conium, its medical properties discussed 231
 — succesful use of, in sciatica 707
 Constitution of gouty persons considered 63, 136, 464
Consumption apparently threatened, when a symptomatic state of complaint, how to be cured 559
 Convalescence, stage of, in gout, how to be conducted 260
 — from acute rheumatism, how to be treated 669
Conversions of disease, the subject referred to 21, 431
 Cooper, Mr. Astley, his experiments on digestion 632
 Cordial, gout, London 189
 Corpulence an usual character of the gouty constitution 63, 136
 — to be guarded against 63, 86
 Cough, occasionally a premonitory symptom of gout 17
 — gouty, nature of 335
 Country air, its importance 605
 Cradle for the bed, its use in gout 240
 Cramps, as premonitory of the paroxysm 20
 — frequent attendant upon the gouty paroxysm 39, 155, 306
 — occurring much in gouty persons 391, 467
 Cruickshank, Mr. his statement that the deposition of the pink sediment implies the existence of diseased liver, not sufficiently qualified 140
 — respecting the lateritious sediment 158
 — his pathology of the pink sediment 140
 — his erroneous grounds of prognosis in gout 170
 Crystals, urinous, commonly called gravel, their chemical nature 160
 — of the ammoniaco-magnesian phosphate, when present, the urine alkaline 590
 Cullen, Dr. his nosological arrangement examined 1
 — his remarks on the atonic state of gout 87
 — his definition of rheumatism 641
 — respecting bleeding as a remedy in acute rheumatism 659
 — his treatment of retrocedent gout, the propriety of discussed 437, 450
 Cupping, cases in which it afforded great benefit to the gouty patient 380, 383
 — when the most fit remedy 176, 182, 472
 Deafness increased by gout 17
 Debility, causes which induce, become predisposing to rheumatism 651
 Desault quoted as stating a case of the prophylactic effect of friction in gout 602
 Desquamation of the cuticle in gout considered 6, 24, 156
 — considered as a character in gout 6
 Diabetes more frequent in Scotland than in England 74
 Diagnosis of acute gout 167
 — of chronic gout 340
 — of retrocedent gout 435
 — of acute rheumatism 656
 — of chronic rheumatism 676

- Diaphragm, how spasmodically affected in a morbid state of the digestive organs 85
- occasionally affected by rheumatism 646
- Diathesis, gouty, its general influence 9
- Dict the most conducive to gout 68, 573, 610
- the most proper during a paroxysm of gout 236
- the most proper in the convalescence from gout 268
- the most conducive to gravel 574
- as to solids, the most proper for gouty persons 606
- as to fluids, the most proper for gouty persons 614
- excess or irregularity in, as a predisposing cause to rheumatism, considered 652
- and regimen the most suitable in acute rheumatism 666
- nutritive, may be occasionally suspended with advantage 617
- and regimen proper in acute rheumatism 666
- Digestion, experiments respecting 632
- Digestive organs, morbid state of, as predisposing to gout, considered 78
- morbid state of, as intimately connected with gout 85
- functions, disordered state of, variously and strongly marked 87
- unhealthy state of, how predisposing to rheumatism 651
- organs, a morbid state of, gives rise to pains which seem rheumatic 91, 677, 688
- morbid states of, generally considered as to the true method of treatment 541
- morbid state of, as occurring in gouty persons, how to be treated 545
- in connexion with a tendency to apoplexy, whether a gouty disposition exist or not, how to be treated 545
- in connexion with a tendency to inflammatory erysipelas, how to be treated 545
- in connexion with nervous symptoms, how to be treated 542, 553
- scrophula, how to be treated 559
- Dinner meal, respecting, for gouty persons 609
- Diseases incidental to the gouty — pathological explanation of 467
- acute, of gouty persons, how to be treated 468
- Disorganisation of parts, caused by gout 43
- Diuretics, their use with cathartics, important in gout 183, 186
- Draught, own favourite formula of, in gout 186
- Drinks, diluent, their free use important 237, 595
- Duodenum, considered as the seat of indigestion 79, 609
- Dura mater, occasionally affected by rheumatism 645
- Dyspepsia, nature of, in gouty persons 142, 333, 338
- in gouty persons, how modified 85, 333
- frequently precedes a fit of gout 87
- attendant on an obstructed state of liver 93
- East wind, as an exciting cause of gout 108
- Eau Medicinale, its sensible properties 216
- — its medicinal powers discussed at length 200
- — its injurious properties in gout 202, 327, 339, 360
- — its sediment 201
- Edinburgh alluded to in regard to gout 72
- Egan, Dr. upon gravel, referred to 577
- Elaterium with opium, recommendation of, by Dr. Sutton 184, 196
- administered in the case of a gouty patient with advantage 380
- Electricity, its successful use in a case of rheumatism 683
- Electuary of Sydenham 623
- Elliotson, Dr. referred to as having entertained a similar opinion respecting the seat of rheumatism with the author 642
- Emetics in gout, their propriety discussed 177, 182, 545
- good effect of, in a case of gout 182

- Emetics, much recommended by Dr. Small 183
- their use as a remedy in acute rheumatism 660
- Empirical treatment of gout, the bad effects shown, 360, 391, 400—and in the various cases, particularly of chronic and of retrocedent gout
- Enteritis in connexion with gout, how to be treated 440
- case of 432
- Epistaxis happens to some gouty persons, particularly in early life 136
- remarkable instance of, in an elderly gouty person 470
- Erysipelas, occasionally produced in connexion with the paroxysm of gout by improper diet 236
- sometimes takes place instead of the fit 467
- Erythema sometimes connected with the gouty paroxysm 95, 467
- Etymology of the word gout 11
- Evaporation, the principle of, as a remedy in gout, fully considered 252
- Exercise, its importance to gouty persons 603
- its importance in the treatment of chronic rheumatism 669, 685, 689
- carried to excess, injurious 112
- excessive, in a case of chronic rheumatism, effected a cure 685
- Exertion, bodily, in the paroxysm, its propriety discussed 238
- EXPERIMENTS on dogs with gout nostrums, and some other medicines—viz. by injection into the circulation.—With Eau Medicinale, 478—Wilson's Tincture, 480—Reynolds's Specific, 481—Hellebore and Laudanum, 482—Hellebore and Laudanum by injection, exp. repeated, 483—Elaterium and Laudanum, 484—Sir Everard Home's vinous infusion of colchicum, 485—Tincture of colchicum (prepared with proof spirit, as expressed at p. 495), 486—Expressed juice of the fresh roots of colchicum, 487—Administered by the mouth, with acetum colchici, 489—Expressed juice of colchicum followed by Eau Medicinale, clear and turbid, 491—Eau Medicinale in the transparent state, followed by Tincture of Colchicum, 494—Wilson's Tincture, 496—Reynolds's Specific, 497—Vinous tinctures of hellebore and opium, 499—Tincture of digitalis followed by laudanum and elaterium, 500—Want's Gout Remedy, 501—Acetic preparation of colchicum, 502—Clear and turbid aqueous decoction of colchicum, 503—Extract from Eau Medicinale, 504—Extract from wine of colchicum, 506—Extract from hellebore, followed by elaterium, and extract from a vinous preparation of colchicum, 507—Extract from acetum colchici, 509—Ditto, 510—Extract from spirituous tincture of colchicum, 510—The same extract inserted into a wound, 512—Sediment from wine of colchicum, 513—An extract procured from the macerated roots of the vinous infusion, according to Sir Everard Home, 513—Ditto from a weak vinous infusion, 515—Ditto from a weak proof spirit infusion, 515—Ditto from the ordinary acetic preparation, 516—Clear vinous infusion of colchicum, 516—Turbid ditto, 518—Elaterium by injection, 519—Resin of opium, followed by crude opium, administered by the mouth, 520—Observations on these Experiments, 522
- Experiments with urine, method explained 126
- as to the temperature of parts under gouty inflammation 148
- by Mr. Astley Cooper, with regard to digestion 632
- chemical, with the gout medicines 216
- with bile 556
- External applications, various kinds of, found injurious in gout 246, 433
- injuries as exciting gout 109

- Facces, respecting 37, 82, 85, 389
 Family susceptibility to gout considered 58
 Fatigue and anxiety jointly considered as exciting gout 113
 Feet, sometimes swell and become licated and dry just before the attack of gout 19
 Festina lenté, an important maxim in the treatment of gout 261
 Fever, sympathetic, an attendant on the gouty paroxysm 3
 Fever, not an essential character in gout 3, 23
 Fish, except of certain kinds, a food of easy digestion 612, 634
 Fit, first, of gout, its symptoms 22
 — its ordinary duration 23
 — sometimes very tedious *ibid.*
 — more mild and regular in men than women 25
 Flannel to be worn in this climate 600, 670
 Flannel bandage, when useful in gout 269
 — in rheumatism 692, 698, 699
 Flannel covering, when improper to a gouty part 245
 — — when proper 250
 Flatus in the stomach and bowels, producing tenderness to pressure 89
 Food, animal, as predisposing to gout 68
 — indigestible, as occasionally inducing retrocedent gout 435, 438
 Fomentations in gout improper 246
 Forbes, Mr. his treatise upon gravel and upon gout referred to 577
 French authors in regard to the treatment of gout 184, 220, 222, 242, 250
 Friction to weakened limbs, its important use 271, 370, 389, 403, 410, 602
 Fruits, subacid, when useful 239
 Functions, digestive, how affected in a fit of gout 36
 Gall, Dr. his peculiar idiosyncrasy in regard to mutton 607
 Galls, infusion of, its effects as a test for urine, not correctly stated by Mr. Cruickshank and Dr. Blackhall 281
 Gall-stone obstruction, severe case of 228
 Galvanism as a remedy 552
 Gibbes, Dr. on the Bath waters, referred to 352
 Glasgow alluded to in regard to the gout 74
 Gonorrhœa, sometimes leads to bursal inflammation 420
 Good, Mason, his nosology referred to 72
 Gout, regular and irregular, terms employed by Dr. Cullen 5
 — parts of the body most liable to be affected by 6, 25, 30
 — predisposing causes to 52
 — hereditary, considered 52
 — arising from a deranged state of the system 142
 — particular causes which call it into action 142
 — pathology of, particularly considered 136
 — own divisions of it into acute, chronic, and retrocedent 12
 — at what period of life it usually occurs 13, 54
 — its general character defined 13
 — acute, symptoms of *ibid.*
 — chronic, its characters defined 14
 — retrocedent, character of *ibid.*
 — acute, history of, and mode of attack 15
 — hereditary, influence of, considered 58
 — disposition to, accidental and hereditary, relative intensity of each 54
 — cause of being more frequent in males than in females 65
 — retrocedent, cases of, and treatment 7, 432
 — its regularity on many occasions as to the order of attack 28
 — sometimes attacks in the daytime, and sometimes most painful in 29
 — its disposition to increase in the constitution 41
 — case of, preceded by very singular premonitory symptoms 85
 — considered by some old authors as contagious 131
 — case of in W. W. in which daily examination was made of the urine, &c. 271

- Gout, case of in J. C. in which the abscess occurred, and in which the urine was albuminous 285
- case of in J. W. in which the urine was albuminous with the presence of pink sediment 292
- case of in J. M. also exemplifying albuminous urine with pink sediment 295
- case of in E. L. in which the urine was albuminous 295
- acute, case of in T. W. who had many uric concretions, and in whom some curious phenomena existed with respect to the urine, which was constantly albuminous 296
- case of in J. L. in which the causes of relapse are pointed out 301
- case of in D. S. related for a similar purpose as the foregoing 303
- case of in J. S. in which the necessity of radical treatment and caution in general management are still more strongly exemplified 305
- case of in B. M. very instructive in many important points 310
- case of in S. J. pointing out the example of an unhealthy state of liver as supporting the disposition to gout 318
- case of, in which an abscess furnishing the urate of soda was formed, one of the few instances in the author's experience 325
- case of, similar to the foregoing 328
- ditto, ditto 329
- some general conclusions respecting, both theoretical and practical 414
- etymology of, the term 11
- occasional suddenness of its attack 22, 378
- first fit sometimes very tedious 24
- first fit usually more mild in men than women 25
- if neglected, the consequences described 41
- as distinguished from rheumatism 144, 367
- how to be treated in the fit, in regard to position of the limbs 240
- radical treatment of 264
- case of in C. R. exemplifying in a striking manner the advantages of active treatment 320
- Gout, case of
- should it be invited? 9, 351
- case of, in V. R.; the constitution highly irritable 388
- case of in J. P. excited by accident, and becoming chronic from the influence of disorder of the digestive organs 386
- its power in suspending other morbid affections 428, 431
- suspected, but proved to be bursal disease 419
- case of, followed at a distant period by aneurism 418
- sometimes the only disease to which the patient seems liable 464
- Gout, chronic, history of 331
- — causes of 336
- — diagnosis of 340
- — theory of 337
- — treatment of 345
- chronic, case of in M. P. in which many urgent symptoms were soon removed 374
- chronic, case of, in A. L. in which much visceral disease was present 377
- chronic, case of, in a lady, in which much visceral disease was present 383
- chronic, case of, very remarkable, successfully treated 404
- chronic, case of, attended with puriform discharge from the urethra 407
- chronic, case of in A. B. a common example of lameness from neglect 409
- chronic, case of in C. D. shewing the intimate connexion of gout with disorder of the digestive organs 410
- chronic, case of, in G. B. affording a similar example 413
- chronic, case of, in J. R. in which much rheumatism was induced by the improper use of flannel 416
- chronic, as blended with rheumatism 361
- chronic, case of in A. B. exemplifying the benefits of friction 371
- chronic, case of in J. W. favourably exemplifying the advantages of regular treatment 390

- Gout, chronic, case of, in C. L. complicated with cough and other distressing symptoms 394
- Gout, chronic, case of in L. M. singular, from being long preceded by a palpitation of the heart 393
- chronic, case of, joined with frequent headach, successfully treated 396
- chronic, case of, in a person highly of the nervous temperament 397
- chronic, case of in C. R. remarkable from the frequency of relapse, and successfully treated 398
- chronic, case of, very complicated, in which friction of the limbs was very useful 401
- chronic, case of, in which the lameness was remarkable 403
- Gout, retrocedent, history of 427
- — causes of 432
- — treatment of 437
- — various cases of, concisely stated 432
- — case of, in S. R. in which the disuse of wine was hurtful 441
- — case of, in R. C. in which a distended state of stomach produced the symptoms 448
- — case of, in C. A. the symptoms depending on bilious obstruction 444
- — case of, shewing the immediate effects of cold applications externally 445
- — case of, in L. S. shewing the true nature of pure gouty spasm 446
- — case of, in R. G. in which bleeding was useful 447
- — case of, in R. B. exemplifying the nature both of spasm and of internal inflammatory action 448
- — case of in T. N. in which general bleeding relieved, after the failure of other means 449
- — case of, or in which the symptoms assumed, in great part, such a character, in J. G. 451
- — case of, chronic in its character, and attended with internal spasm 457
- seemingly retrocedent, very anomalous in the symptoms 460
- — chronic, case of 457
- Gouty affections suddenly transferred from one place to another 29
- diathesis, necessary existence of, in order to the production of the paroxysm by the influence of the remote causes 111
- Gratiola, or hedge hyssop, its inert action in gout 193
- tincture of, its sensible properties 217
- Gravel, as a symptom of faulty digestion 98, 584, 585
- external characters of 99
- occurs to gouty persons chiefly early in life 99, 467, 571
- theory of 571
- and gout considered pathologically in a comparative point of view 573
- treatment of, generally considered 583
- case of, successfully treated by alterative aperients 584
- giving rise to inflammatory action of the kidneys, how to be treated 595
- how distinguished from lumbago 677
- the regimen proper for persons subject to 619
- Gravity, sp. of urine, in health 96
- Greeks, their various terms for gout 10
- Greek MSS. on gout 193
- Greek motto applied to gouty persons 76
- Gregory, Professor, his condemnation of Eau Medicinale 203
- — gout scarcely known among his clinical patients 72
- — his observation with regard to the east wind 108
- Grosvenor, Mr. of Oxford, his treatment by friction recommended in certain states of the limbs induced by gout 370
- his peculiar practice when useful in chronic rheumatism 689
- Guaiacum, its use in rheumatism considered 685
- sometimes an useful addition to purgative medicines in gout 349
- Guilbert, M. his pathology of gout 132
- Habit in disease, explained 359, 360

- Hæmorrhage, spontaneous, to a certain extent salutary 470
- Hæmorrhoids in gout 40, 88, 136, 334, 377, 466
- Hæmorrhoidal discharge sometimes injurious 552
- Haller, his opinion that the nerves are the seat of disease in gout 154
- Hamilton, Dr. Lynn Regis, his divisions of gout into acute and chronic 12
- his opinion that in gout a fluid is formed *sui generis*, which he calls the gouty fluid 171
- Hamilton, Dr. Edin. his method of practice in a morbid state of the alimentary canal 543
- Harrowgate water an useful alterative 358
- Haygarth, Dr. his statement of the rarity of the acute rheumatism compared with cases of the chronic 340
- his diagnostic distinctions objected to 341
- Haygarth, Dr. his strong recommendation of the Peruvian bark, as a remedy in acute rheumatism 659, 664
- Head, injury of, disturbs the digestive organs remarkably 553
- to be washed with cold water 603
- Headach, connected with gout, successfully treated 395
- Health, the true method of preserving 631
- Heart, diseases of, as stated to be in connexion with rheumatism by Sir David Dundas 646
- palpitation of 16, 92, 334
- Heart-burn, sometimes a warning symptom of gout 177
- Heat of the fire in the apartment, to be guarded against in the paroxysm 246
- Heat of weather, predisposing to gout 77
- an exciting cause of gout 27, 78
- Heberden, Dr. his opinion that children do not have gout 61
- his caution on the use of the Bath waters for the gouty 351
- his forcible description of the calamitous consequences of gout 465
- Held, Dr. in praise of bark as a remedy in gout 219
- Hellebore, tincture of, its very injurious qualities as a remedy in gout 191, 304, 305
- Hellebore, tincture of, with laudanum 196
- its sensible properties 217
- Helmont, Van, that the gout is contagious 131
- Henry, Dr. Manchester, referred to respecting urea 557
- Hereditary predisposition to gout, the question considered 3, 52, 135
- and acquired gout, comparative tabular view of 53
- structure, as a remote cause of rheumatism, considered 650
- Hermodactyl of the ancients 193
- Hernia, example of, causing great irregularity in the action of the bowels 545
- Herring, the praise of, as a remedy in gout, by Dr. Clerk 612
- Hippocrates, his aphorism as to the adult age in connexion with gout 60
- as to the period of life when gout attacks females 66
- advised the burning of gouty parts with raw flax 244
- Home, Sir Everard, on the irritable state of the urethra in gout 18
- his recommendation of the use of magnesia in gout and gravel, referred to 116, 582
- his opinion of the Colchicum Autumnale and Eau Medicinale discussed 197, 215
- Home, Professor, Edin. his case of retrocedent gout, which proved speedily fatal 432
- Horace, his poetical pathology 114
- Horse exercise, a valuable remedy in most chronic diseases 604
- when useful in chronic rheumatism 687, 689
- Hot climate in some instances favourable 600
- Hour of dining 608
- Howship, Mr. his case of gouty dissection 49
- Humoral pathology, the doctrines 173, 623
- Humours, The, according to Hippocrates 64
- Humulus lupulus, an inert medicine 232
- Hunter, Mr. his opinion that gout is not always an act of the constitution, discussed 143
- his description of the sensations of gout 147

- Hunter, Mr. his question how far the inflammation of gout may affect the brain and stomach 429
- Hunterian Museum at Glasgow, various specimens of disease from gouty concretions, in the 47
- specimen of a stomach gangrenous from gouty inflammation, in the 436
- Hydrothorax an occasional remote consequence of gout 466
- produced by the Eau Medicinale 203
- Hyoscyamus niger, its medicinal properties discussed 231
- Hypochondriasis, commonly connected with a faulty state of the digestive organs 87
- frequently results from dyspepsia 87
- why prevailing in gouty persons 142
- in gout 333, 379
- Icthyocosis, as a consequence of gout 45
- Indies, East, a residence in, has proved a cure to some gouty persons 600
- Indigestion, when prevailing in the stomach 80
- various symptoms of *ibid.*
- when prevailing below the stomach 81
- Indolence predisposing to gout 74, 603
- Inflammation of gout, its peculiar character 143, 146
- Inflammation, as being constitutional or only local, rationale of treating 471
- Ingesta, excess of, its hurtful effects 138
- Injection of solvents into the bladder condemned 581
- Injuries, external, an exciting cause of the paroxysm 109, 143, 272
- Insurance on the lives of gouty persons requires an increased premium 465
- Intemperance, excessive, as an exciting cause of gout 101
- Interval in the paroxysms of gout 40
- Irritable, term defined 65
- Issues, very inefficient as a prophylactic remedy in gout 630
- James's Powder, the most useful preparation of antimony 185
- Jaundice as attendant on disordered digestive functions 84, 318, 452
- Johnson, Dr. his translation of M. Guilbert's work on gout 133
- Joints, the, in what manner occasionally affected by acute rheumatic inflammation 673
- — by chronic rheumatic inflammation 675
- Juvenal, his supplication for health of body with a sound mind 605
- Kidneys, the importance of their function 95, 139, 555
- inflammation of the, how to be treated 470, 595
- function of, hypothesis respecting 555
- Kinglake, Dr. his cold water practice in gout condemned 252, 310, 433
- Lactucarium*, its medicinal properties discussed 232
- its use in conjunction with stramonium in rheumatism 688
- Lameness consequent to gout, how to be treated 369
- Latham, Dr. on the best nosological divisions of gout 12
- Leamington water useful to gouty persons 358
- bath, its good effects 368
- Leather waistcoat sometimes useful 601
- Leeches, their use as a remedy in gouty inflammation, discussed 241
- their use in chronic rheumatism discussed 689
- Lewis Cornaro, his treatise referred to 598
- Ligaments, their morbid changes of structure, from gout 44
- — from rheumatism 675
- Ligaments, weakened state of, in gout 309
- Liger, his theory of hot and cold gout explained 132
- Lime water, as a prophylactic remedy for gouty persons, considered 628
- Liniment, stimulating to parts in a languid state of action, formula of 271
- Linnaeus, his opinion that drinkers of malt liquors only do not have gout 71
- Liquors, the comparative influence of with respect to gout, considered 69, 618

- Liquors, malt, especially when in conjunction with spirits, may excite the paroxysm of gout 104
 — malt, their use considered with respect to gravel 619
 — subacid, made from our own fruits, improper both for persons subject to gout and gravel 69, 619
 Liquor opii sedativus 228
 Liver, signs of diseased state of 90
 — diseased state of, as causing a relapse of gout 302, 307, 314, 378, 413
 — disordered action of, inducing a relapse of gout 415
 — diseased state of, causing a curious modification of gout 451, (Case ix.)
 — disordered state of, producing many anomalous symptoms in a case unconnected with gout 91
 — when chiefly the seat of complaint, how to be treated 549
 — the disordered functions of, a cause of protracting a fit of gout 41, 415
 — a diseased state of, connected with gout 85, 318, 377
 — disordered state of, produces pains which are to be distinguished from rheumatism 677
 — disease of, as consequent to gout 42, 264
 Local appearances in gout 30
 — treatment in the paroxysm 241
 — — in acute rheumatism 667
 London, prevalence of gout in 72
 — gout cordial 189
 Lotion, evaporating, own formula of 254
 — its good effects in gout related 257
 — its rationale considered 259
 — its use in rheumatic inflammation considered 667
 — when useful in chronic rheumatism 689
 — for chalk-stone concretion 283
 Lucian, the Greek poet, his amusing list of remedies in gout quoted 244
 — his poetical description of the miserable consequences of gout 174
 Luncheon, kind of, proper for gouty persons 608
 Lumbago, description of 676
 — proper treatment of 690
 — with disorder of the nervous system, case of 699
 Lumbago, with much general debility affecting several joints, case of 700
 — and sciatica, interesting case of 708
 Lungs, disordered state of, in connexion with a morbid state of the digestive organs, and the treatment required 559
 Magendie's Essay on Gravel, referred to, and its doctrines discussed 574
 Magnesia, recommended by Sir Everard Home and Mr. Brande, in gravel and in gout 116, 582
 — in what manner useful in gout 166, 583
 — its inadequate powers as a remedy in gout 312, 318
 — its virtues as a remedy in gravel critically examined 583
 — its virtues as an habitual medicine for the use of gouty persons critically examined 625
 — its peculiar mode of producing serious occasional injury 627
 Malt liquors, strong, as predisposing to gout 72, 618
 — — as exciting gout 105
 Marcet, Dr. referred to respecting stramonium 234
 Materies morbi, as to gout 115
 Meat, should be eaten only once a day 610
 Medicine, the prophylactic use of in regard to gout 622
 Melancholic, the term considered 64
 Merchant, anecdote of with respect to gravel 588
 Mercury, the imprudent use of, produces a severe form of chronic rheumatism 678, 703
 — how this form of complaint is to be treated 690, 703
 — to be used with caution in gout 190, 385, 405
 — to be used cautiously with scrophulous patients 559
 Mercurial medicine, formulæ of 266
 — preparations, their propriety in gout considered 189, 267
 Milk and vegetable diet, for gouty persons, and for others under certain states of repletion, considered 617
 — case in which its temporary adoption proved useful to a gouty person 321

- Mind, state of, its influence on the functions of the animal economy 68, 113, 554
 — state of, as predisposing to gout 68
 — state of, as exciting gout 113, 340
- Misplaced gout, the propriety of the term considered 7
- Monro, Dr. jun. on the effects produced upon the bones of the extremities by gout 45
- Moore, Mr. of his preparation of tincture of hellebore and laudanum 191, 196
 — — on the surgical treatment of ulceration from gouty concretions, referred to 423
- Morbific matter, doctrine of, combated by Cullen 9
- Morgagni, his description of his own case, in which ophthalmia yielded to an attack of gout 21
 — his case of dissection of a gouty patient, whose kidneys contained calculi in remarkable quantity 43
- Morphium*, newly announced principle of opium 230
- Moxa, burning with 243, 338
- Mucous membranes become disordered by the influence of gout 17
- Mucus, excess of, in the bowels, in long standing indigestion 82
 — of the stomach, in great excess in a case 384
 — excess of in the urinary bladder 37, 97, 100
 — pathological explanation of 162
- Muriatic acid bath, as a remedy in gout, considered 247
- Murray, Dr. his original and instructive opinions respecting the natural composition of waters, referred to 621
- Musgrave, his distinction of anomalous gout disapproved 8
 — his recommendation of blisters to parts left diseased from gout 374
- Mutton of easy digestion, and most so when roasted 611
- Nails, occasional brittleness of, 46
- Nature often not a good physician in the gout 175
- Narcotics, as a remedy in acute gout 221
- Narcotics, general conclusion respecting their use in gout 236
 — their use in chronic gout 346
- Neck, stiff, described 673
- Neckcloth, tight, to be guarded against by the corpulent and plethoric 475
- Nephritis, case of, in a gouty person, successfully treated by venesection, &c. 470
 — calculosa, how distinguished from lumbago 677
 — — Morgagni's case of 43
 — — treatment of 595
- Nerves, extreme sensibility of, in gout 17, 38, 85, 142, 148, 287
 — affected by rheumatism, cases of, and how cured 681, 705
- Nervous medicines, when admissible in gout 351
 — pains to be distinguished from rheumatism 678
 — term defined 65
- Nettles, stinging with, spoken of by Cullen as a hazardous remedy in gout 243
- Nitric acid, the most delicate test for albumen 293, 300
- Nodosity of the joints, Dr. Haysgarth's character of 343
- Nosology of gout, Cullen's 1
- Nostrums, gout, various, compared 196
- Numbness in the limbs occasioned by a disordered state of the digestive functions 88
- Observations on the experiments with the gout and other medicines administered to dogs 522
- Oedema in gout described 14, 24
- Onions, dressed, usually agree with gouty persons 613
- Opium, its valuable properties as a remedy in gout considered 221
 — with James's powders, formula of, in pills 224, 277
 — its effects remarkably modified by severe pain 224
 — to be used in complaints of the kidney, with much limitation 596
 — its use in acute rheumatism considered 662
 — its useful influence in gout 133
 — plaster, useful in local chronic rheumatism 699, 701
- Ophthalmia, case of, yielding to the supervention of gout 21

- Ophthalmia, in connection with gout 17
 — rheumatic 674
 Orfila, M. his opinion respecting opium 227
 — in regard to the preparing of vegetable medicines 232
 Oxalate of lime in urinary sediment 166
 Oxalic acid, the most delicate test of urea 297, 557
 Oxy muriate of mercury, its effects as a test for urine, not correctly stated by Mr. Cruickshank 281
 Oysters wholesome and very nutritious 612
- Pain of gout, its intensity described 32, 34, 154
 — except where arising from common inflammation, almost invariably demands and justifies the *free* use of opium 223, 225, 277, 596
 — of gout varies according to situation 32
 Palpitation of heart, in gouty persons 16, 92, 334, 394, 397
 Paralytic appearance of parts from gout 32
 — appearance of parts in protracted gout 306
 Paroxysm of gout, treatment of the 178
 — of gout described in detail 21
 Parr, Dr. his definition of the term specifics 197
 Parry, Dr. on the conversion of diseases, referred to 21
 — his pathology on gout referred to 134
 — his opinion respecting the use of the Bath waters for gouty persons 352
 — his cases of injury from Kinglake's treatment 433
 — his opinion respecting Eau Medicinale 203
 Parts of the body usually affected by a first fit of gout 25
 — — most painfully affected by gout 32
 — — most generally affected by gout 28
 Passions of the mind as exciting gout 113
 — — as curative in gout 240
 — — predisposing to gout 68
 Pathology of Cullen 9
 — own, of gout 136
- Pathology of rheumatism 3, 654
Patience and flannel, futility of 245
 Pediluvium improper in gout 246
 Pemberton, Dr. reference to his treatise on various diseases of the abdominal viscera 544
 Pericardium, in what manner secondarily affected by rheumatism 645
 Periodical nature of gout, 27, 143
 Periosteum sometimes the seat of rheumatism 644
 Perry, an unfit liquor for the gouty 618
 Perspiration produced in excess, not approved by the author in the treatment of gout 220, 245
 Perspiration, as occurring naturally in a hot climate, much counteracts the formation of gout 599
 — excessive, as a cause predisposing to rheumatism, considered 652
 — sudden cessation of in the feet sometimes precursory to the gout 19
 — gouty persons in general not deficient in 600
 Phenomena, some very curious, related by gouty patients 154
 Phosphoric acid, question of, with relation to gout 119
 Phosphates, earthy, their chemical composition, how ascertained 163
 — — when in excess, what practical indications affording 98, 555
 — — in excess in gout. See tabular view 120, 139
 — — when appearing suspended, not soluble in the urine, even in its natural temperature 162, 580
 — — when in excess, the treatment considered 589
 — — sediment of, most prevails in children 593
 Pickles unwholesome, unless old and well prepared, and then to be taken only in moderation 613
 Pills, alterative 266
 — opiate 224, 277
 — purgative, formulæ of 283, 548, 623
 Pink sediment 37, 97
 — pathology of 571
 Pitt, Mr. anecdote of, and father, with respect to gout 68
 Plethora, as predisposing to gout 76, 137
 — the true manner in which to be counteracted 629

- Plummer's pill, an efficacious alterative 266, 302, 624
- Pneumonia, in two cases yielded to the supervention of gout 20
- in another case, pursued its course with the paroxysm 21
- how to be treated 470
- Podagra, an objectionable term 4, 26
- Pork usually proves very indigestible 611
- broth said to suit a weak stomach 612
- Port wine, instance of its always disagreeing in a remarkable manner 617
- Porter, its free use assists the predisposition to gout 72
- Portland powder, account of 630
- Poultices in gout as a mode of evaporation considered 249
- useful form of one, if at all employed in gout 249
- of Pradier 250
- Precipitation, instances of its chemical misapplication as a term 293
- Predisposing causes to rheumatism 650
- Premonitory symptoms of gout 15
- sometimes wholly absent 15, 377
- — treatment of 176
- Pringle, Sir John, his treatment of acute rheumatism 658
- Probart, Mr. referred to as preparing genuine *lactucarium* 233
- Prognosis in acute gout 170
- Prophylaxis in regard to rheumatism, considered 669
- Prophylactic course of medicine occasionally useful 624
- regimen in regard to gout 597
- treatment of gout successful 328
- Prostrate gland affected in gout 430
- Prout, Dr. on a specimen of albuminous urine 297
- referred to respecting gravel 585
- Proust, his opinion concerning the pink sediment 156
- Proximate cause of gout 115
- — of rheumatism 653
- — definition of 130
- Pulsation of the aorta in certain morbid states of the digestive organs 91
- Pulvis ipeacuanhæ comp. when useful in gout 346, 362
- — in rheumatism 663, 688
- Pumping, dry, useful in sciatica 689
- — term explained 353
- Punch, a favourite beverage at Glasgow 73
- Purgatives, when to be particularly preferred in the form of pills 546
- Purgative treatment in gout 153
- Purpura simplex, remarkable case of, after the paroxysm, caused by too low a regimen 268
- Pus, how to be distinguished from mucus 384
- Quantity of food, an important object of attention 608
- Quincy, quotation from, respecting colchicum 194
- Ratio symptomatum in regard to gout 135
- — in regard to rheumatism 656
- Regimen, prophylactic, in gout 597
- and diet, the most suitable in acute gout 236
- proper for persons subject to gravel 619
- and diet proper in acute rheumatism 666
- Relapse in gout, causes of 40, 301
- — tendency to, aggravated by empirical treatment 360
- Residence, particular situation of, as to soil, &c. most to be preferred for the gouty 599
- Respiration frequently difficult in a morbid state of the digestive organs 86
- Rest in bed, the hours of, most favourable to gouty persons considered 605
- Retrocedent gout, the history of 427
- — causes of 432
- — cases of, concisely stated 432
- — treatment of 437
- Reviewer, mistake of, concerning the urinary sediment 37
- Reynolds's Specific 196, 200, 360
- — its sensible properties 217
- Rheumatic gout, the term 342
- Rheumatism, how distinguished from gout 167
- occurring in union with gout 36, 416
- introductory observations to the short treatise on 639
- etymology of 641
- pathology of 8, 532

- Rheumatism, its nature defined 655
 — as produced by the imprudent use of mercury 690
 — how far dependant on a morbid state of the digestive organs 651
 — how to be treated 688, 693, 697
 — acute, definition of 655
 — — diagnosis of 656
 — — seat of 642
 — — symptoms of 646
 — — sequelæ of 649
 — — predisposing causes to 650
 — — exciting cause of 652
 — — proximate cause of 653
 — — treatment, constitutional, of 657
 — — — local, of 667
 — chronic, its general history and particular symptoms 671
 — — — diagnosis of 676
 — — — sequelæ of 674
 — — — treatment of 679
 — — — of the nerves, treatment of 681
 — — — cases of 688
 — chronic, case, with error of the secretions 692
 — — case, ditto, and also kept up by fatiguing bodily exertions 694
 — — case of, chiefly supported by error of the secretions 697
 — — — of the shoulder joint, case of 700
 — — — of the knee, succeeded by lumbago and sciatica, case of 704
 — — — produced by cold after mercury 703
 — — sub-acute attack of, excited by cold 695
 Richerand, his opinions respecting the phosphoric ingredients of the urine 119
 Ring, Mr. his related case of fatal result from Eau Medicinale 203
 Robinson, Bryan, his calculations of disease from supposed qualities of the blood 130
 Rollers, when useful in gout 269
 Rosacic acid, theory of, by Pronst, objected to 156
 — — by M. Vogel, disproved 158
 Rose hue, concerning the 157
 Rose, Mr. his statement respecting urea discussed 557
 Rubefacients, useful in some states of chronic rheumatism 682, 688, 691
 Rush, Dr. in the author's opinion advised bleeding as a remedy in gout too indiscriminately 182
 Rush, his observations on the diseases of the North American Indians 70
 Salivation, occasioned in gout by the improper use of mercury 190
 Salt water, how artificially made 601
 Sarsaparillæ decoct. comp. concerning it as a medicine 268, 624
 Saunders, Dr. his opinion that as an external remedy, the influence of Bath and Buxton waters depends on temperature only 365
 Sauvages, his complicated nosology in gout 10
 Scheele referred to as having first discovered uric acid 579
 Sciatica, theory of 644, 671
 — how to be treated if inflammatory symptoms present 681
 — very remarkable case of, and how cured 685
 — case of, attended with general nervous relaxation 696
 — with constitutional debility, case of 706
 — sedative treatment in, highly useful 709, 710
 — produced by local injury, case of 710
 — to be distinguished from diseases of the hip joint 711
 Scotland, gout less frequent in, than in England 72
 Scott, Dr. his practice with nitromuriatic acid considered 563
 Seasons of the year predisposing to gout 27, 77
 — — as predisposing to rheumatism 652
 Sea, those who live upon the, as much affected by gout as when on shore 78
 — bathing, cold, recommended by Dr. Heberden as occasionally proper for gouty persons 363
 — — its propriety discussed *ibid.*
 — — warm, its particular effects in a case 367
 — — warm, useful in chronic rheumatism 706
 Secretions, derangement of, in the paroxysm of gout 37
 Sedatives, as a remedy in acute gout, considered 221

- Sedatives, in acute rheumatism considered 662
- Sediments of the urine 37
- — pathology of 118, 139
- — chemical composition of 156
- — a knowledge of, very important in a practical point of view 139, 555, 592
- — pink and lateritious 37, 97, 156
- — when blackish green in colour, may be suspected to contain oxalate of lime, how proved 166
- Sensations, intensity of, in acute gout, compared in different situations 32, 154
- — in gout, the peculiar nature of 33, 154
- — changes of, according to the alteration in the symptoms 147
- Serous membrances, in what manner secondarily affected by rheumatism 645
- Sequelæ of acute gout 41
- of acute rheumatism 649
- of chronic rheumatism 674
- Sex, male, as predisposing to gout 65
- the, considered with relation to rheumatism 651
- Shell fish not well suited to a weak stomach 612
- Skin to be kept cool in the paroxysm of gout 221
- Sleep, in a peculiar manner, occasionally not disturbed in a first attack of gout 22
- profound, through the night, in a case of the actual invasion of a severe fit ib.
- Small, Dr. his mode of treating the gout 183, 219
- — recommended bodily exertion in the paroxysm 239
- Smith, Dr. Carmichael, his definition of the seat of rheumatism called in question 643
- Soap plaster, when useful 251, 283, 373, 376
- Soda, subcarbonate of, given very largely as a prophylactic against gout, without success 626
- water in moderation useful to the dyspeptic stomach 609
- how rendered an agreeable drink 608
- Soup to be taken sparingly by gouty persons 610
- Spasm, internal, in gout, 434, 440, and in the cases of retrocedent gout
- Specifics, pretended, the most notorious of modern date considered 191
- Spirits, in what manner pernicious 70, 72
- in every way in which they can be used, more or less pernicious 74, 418, 619
- Spleen, disease of, occasionally consequent on gout 42, 338, 263
- Sponging the feet or other parts weakened by gout, every morning, strongly recommended by the author 270, 601
- Stanley, Mr. his interesting case of dissection 675
- Station of life and occupation most predisposing to gout 66
- Steel, the best preparations of, considered 262
- Stimulant treatment in chronic rheumatism defined 682
- Stimulant medicine, formula of 351
- Stomach, state of in indigestion 80
- liable to be affected by gout 4, several of the cases of retrocedent gout
- when most the seat of complaint, how to be treated 545
- the medium through which the gout is created 138
- Stone of the bladder of rare occurrence in gouty persons 43, 467
- by what habits of living produced 576
- various statements respecting 581
- Stone, as a sequel to gravel 99
- Stooping, its occasional effects in producing spasms 85
- Strain sometimes mistaken for gout 23
- Stramonium datura, its medicinal properties 233
- Stramonium and lactucarium, when useful in gout 235, 346, 362
- use of in rheumatism 688, 697, 699
- Stricture of the urethra increased by gout 18
- Structure, diseased changes of from gout 43
- hereditary, as predisposing to rheumatism 650
- Study, severe, predisposing to gout 68
- — exciting gout 606

- Sudorifics to be used with caution as remedies in gout 220
- Sudorifics as a remedy in acute rheumatism considered 661
- Sugar administered in a case of gravel 586
- Suppuration very rare from gouty inflammation 146
- Suicide committed by the friend of Pliny 155
- Sutton, Dr. his pathology of gout 133
- — his treatment of gout 183, 224
- Swelling takes place early from gouty inflammation 6
- Swieten, Van, his statement that the Dutch knew little of the gout, till they changed their beverage of beer for wine 71
- his idea that gout is a contagious disease 131
- his amusing account of the gouty dancing master 239
- his account of gouty persons cured by removal to a warm climate, referred to 599
- his narration of some gouty persons, by being reduced to adversity, losing their disease, referred to 605
- his idea respecting chalk-stones 422
- Sydenham, the first author who wrote with perspicuity on gout 11
- his description of the bursal swellings 30
- his imperfect description of the sense of heat in gouty inflammation 33
- his description of the sensations of gout 38
- his mistake with regard to the gout in females 61
- his opinion respecting the cause of gout 79
- his prohibitions to the medical treatment of gout condemned and answered 173
- on the radical method of treatment necessary to alter the gouty constitution 264
- his advice for the prevention and cure of chalk-stones 426
- his great praise of horse exercise 604
- his account of severe relapse in his own case from immoderate study 606
- Sydenham, his Essay on Rheumatism referred to 640
- his general opinions respecting gout 40, 60, 79, 173, 221, 264, 332, 422
- Sympathetic pains often mistaken for rheumatism 91
- Symptoms, premonitory, of gout 15
- some peculiar, which occasionally precede a fit of gout 17
- of a first fit of gout 22
- constitutional, of the paroxysm 36
- some remarkable, occasionally attendant on a fit of gout 39
- of a morbid state of the digestive organs, detailed 80
- Symptoms of seeming internal gout, very anomalous 460
- Synovial membranes inflamed during the active state of the rheumatic diathesis, how to be treated 690
- Syphilitic pains resembling rheumatism, case of 701
- Ditto, in which the pain affected the forehead, case of 702
- produced by cold after mercury, case of 703
- Tabular view of the relative proportions of phosphoric acid in urine 120
- Table, an analytical, exemplifying the author's plan in investigating the phenomena of gout 476
- Tactus eruditus*, in examining the state of the abdominal viscera, by pressure, to be used with cautious deductions 89
- Temper, how affected by an approaching fit of gout 114
- Temperament of gouty persons considered 63
- Temperaments defined 203
- the sanguineous, stated by Dr. Cullen, as most predisposing to rheumatism 651
- distinction of 64, 203
- Temperance and exercise indispensable to health 631
- Temperature, vicissitude of, an exciting cause of gout 27, 600
- of parts inflamed in gout 148
- in rheumatism 152
- in a whitloe and in a burn 153

- Temperature, vicissitude of the, exciting cause of rheumatism 652
- Temple, Sir Win. his successful employment of the *moxa* in a paroxysm 243
- — his high praise of the effects of regular friction 602
- Tendons, morbid changes of structure in, from gout 44
- — from rheumatism 649, 675
- subject to severe spasmodic action in gout 332
- Textures the most affected by gout 146
- affected by rheumatism 642
- Thermometer for animal heat, description of the mode of using 156
- Tincture of colchicum, a hurtful preparation in gout 198, 525
- — its sensible properties 217
- Tin tube described, and its use in affording the means of heated air, as a remedy in gout 247
- Tongue, appearances of, which indicate an unhealthy state 80
- Tonics, their propriety in gout diseussed 220, 265
- corrective, formulæ of 265, 266, 430, 551
- stimulant, formula of 350
- their prophylactic merit in gout considered 631
- Travers, Mr. referred to in connexion with an inquiry into the cause of gravel and stone 576
- Treatment of gout generally considered 173
- local 241
- general, in acute rheumatism 657
- local, in acute rheumatism 667
- of chronic rheumatism 682, and see the cases
- Tunbridge Wells water, its use in certain states of dyspepsia 547
- — chemical statement respecting *ibid.*
- — excites gout 357
- Urate of soda, how secreted 51
- — enters into the composition of the pink and lateritious sediments 159
- — concretions, cases of 273, 296, 423
- Urate of soda found in the matter of the gouty abscess 291, 325, 328, 329
- Urea, excess of, in gout 138, 284, 378
- — always found by the author in excess in a congested state of the vena portarum system 555
- the most delicate test for, considered 557
- found by the author to be in excess, in several cases of chronic hepatitis 558
- Urethra often remarkably irritable in gout 18, 404, 407
- Uric acid, supposed by some authors to be the proximate cause of gout 115, 144
- — does not appear to be always secreted in excess, in those cases where a crystalline precipitate is formed in the urine 117, 578
- — and soda, form the composition of gouty concretions 146
- — sought for in the matter of transpiration, without success 577
- — sediment in the urine, its practical indications 592
- — in the cases of gouty concretions, conceived by the author to be secreted *vicariously*, 145, 329, 587
- Uric concretions, the chief source of the doctrine of a morbid matter 115
- — of rare occurrence in gout 116
- Urine, its alkaline or acid state considered, Exp. 38, p. 125, 577, 589
- how affected during a paroxysm of gout 37, 95
- its average specific gravity in a state of health 96
- ditto, under a diversity of diet 96
- its high specific gravity in gout 37
- the morning, to be chosen for medical examination 126
- when containing the earthy phosphates suspended, known by its appearance 162
- albuminous 288, 293, 295, 296
- — its pathological indications discussed 298, 554
- the changes of, as premonitory of a fit of gout 95
- when loaded with sediment of high specific gravity 117, 162

- Urine, morbid state of, connected with disorder of the digestive organs 118
 — tabular view respecting 120
 Urticaria sometimes precursory to the paroxysm of gout 95
 Vapour bath, portable 323
 — — successful use of the, in a case of mercurial rheumatism 703
 Vapour, local, as a remedy in chronic rheumatism 682, 691
 Veal most easily digested when boiled, and why 611, 635
 Vegetable medicines, their most active state of preparation 231
 Vegetables, their use for gouty persons and the dyspeptic, considered 613
 Veins of the leg, how affected by continued gout 31, 288, 377
 — hæmorrhoidal, fulness of, a very usual complaint among the gouty 40, 88, 136, 334, 377, 466
 — varicose, as a consequence of gout 45, 377
 Vena portarum system, in a state of congestion, intimately connected with the production of gout 138, 337
 — — — congestion of, indicating obstruction in the circulation of the liver 552
 — — — when existing, how to be treated 184, 555
 Vinum Colchici, trusted to as a specific, why objectionable 360, 537
Venus nimia, as predisposing to gout 76
 Vesicatories and irritants, the employment of in gout condemned 243, 374
 Vibices in the lower extremities of gouty persons, occasionally found 45, 317
 Virgil, his praise of the life of the rustic 67
Vis medicatrix naturæ, its powers considered 137, 174, 345
 Vogel, M. his recently published opinion concerning the rosacic acid confuted 158
 Vomit, black, example of, referred to 436
 — cum as a remedy in gout too extravagant 194
 Wardrop, Mr. on rheumatic inflammation of the eye 674
 Warmth of covering to the inflamed parts in the paroxysm condemned 244
 — — even in the time of health, the excess of should be guarded against, in clothing the feet 601
 Warner, Rev. Mr. his strong recommendation of opium to relieve pain in the paroxysm, referred to 221
 Water, cold, its application as a remedy in gout condemned 252, 433, 310, 433
 — as a diluent beverage, the most favourable to health 618.
 — its qualities as an article of diet, chemically and generally considered 620
 Watson, Mr. his curious case of the dissection of a gouty patient 45
 Wells, Dr. on albuminous urine 293, 299
 Whey, rennet, an useful nutritive drink in the paroxysm of gout 237
 — — its use as a nutritive drink in the acute rheumatism 658, 666
 Whiskey, the too common beverage among the lower orders in Scotland 74
 Whytt, Dr. his instance of the prophylactic powers of lime water in gout, referred to 628
 Willan, Dr. referred to respecting the use of chlorine 560
 Williams, Dr. his publication on Wilson's Tincture 200
 Wilson's Tincture 196, 199, 360
 — — its sensible properties 217
 Wine of colchicum 195, 323
 — — its sensible properties 217
 Wine, use of, when admissible in the paroxysm 237
 — disuse of, discussed in regard to the question of preventing gout 322, 441, 616
 — quantity of, proper for gouty persons 615
 — kinds of, the most proper for gouty persons 617
 — excess in, as exciting gout 101
 — of good quality, and under certain restrictions, allowable, and even useful, to gouty persons 315, 322, 401, 614
 Want, Mr. his praise of colchi-

Wollaston, Dr. his opinion that in gouty subjects there exists a redundance of uric acid	116	Woodville, Dr. his account of the destructive effects of hellebore on the stomach, incautiously given as a medicine	192
— — his experiment on the composition of gouty concre- tions referred to	146	Wool, combed, as an application to the gouty part, much con- demned	245
— — his paper on the various modes of gestation for invalids, referred to	604	Young, Dr. referred to as having pointed out a test to distinguish pus from mucus	384
— — his recommendation of the use of alkalies in gout	625	Youth not subject to gout	59
Women much less liable to gout than men	65		

